


MAP 14A – SIMPLIFIED GEOLOGY AND AREAS OF HIGHEST MINERAL EXPLORATION AND MINING POTENTIAL



- QUATERNARY** 0 – 1.8 million years before present
 - Gravel, sand, clay, mud and minor limestone.
- TERTIARY** 1.8 – 65 million years before present
 - Unconsolidated or cemented gravel, sand, silt and clay, minor limestone and brown coal.
 - Basalt and related sediments.
- CRETACEOUS** 65 – 141 million years before present
 - Intrusive and volcanic igneous rocks of various compositions (Cygnet and Cape Portland areas only).
- JURASSIC** 141 – 205 million years before present
 - Dolerites.
- TRIASSIC** 205 – 251 million years before present
 - Sandstone and mudstone, minor black coal, basalt and volcanic sediments.
- PERMIAN – LATE CARBONIFEROUS** 251 – 314 million years before present
 - Mudstone, pabbly mudstone, sandstone and conglomerates, minor limestone, black coal and of shale.
- EARLY CARBONIFEROUS – ORDOVICIAN** 340 – 490 million years before present
 - Granite and related intrusive and minor volcanic igneous rocks of various compositions.
 - Folded and locally deformed sandstone, quartzite, siltstone, shale and slate, minor quartz veining.
- ORDOVICIAN** 434 – 490 million years before present
 - Limestone and minor sandstone, siltstone and shale.
- MIDDLE ORDOVICIAN – LATE CAMBRIAN** 460 – 500 million years before present
 - Folded and locally deformed conglomerate, sandstone, quartzite, siltstone and shale.

- LATE – MIDDLE CAMBRIAN** 490 – 510 million years before present
 - Folded, deformed and altered volcanic rocks of various compositions, and related intrusive igneous rocks, sandstone, siltstone and conglomerate (includes Mt. Read Volcanics).
- MIDDLE – EARLY CAMBRIAN** 510 – 545 million years before present
 - Serpentine and greenstones. Deformed and metamorphosed volcanic and intrusive igneous rocks of ultra mafic and basaltic compositions, and associated sandstone, mudstone and chert.
- NEOPROTEROZOIC** 545 – 1000 million years before present
 - Deformed and metamorphosed basaltic volcanic rocks and associated sandstone, siltstone, shale, dolomite, chert and schist.
- MESOPROTEROZOIC** 1000 – 1600 million years before present
 - Granite (King Island only).
 - Deformed and metamorphosed quartzite, siltstone, conglomerate, schist and dolomite.

Scale: 1:100000
ACZB2 – AMG Zone 55
Contour Interval: 100 metres

The simplified geology for this map is derived from the 1:250000 digital geology of Tasmania.

Mineral potential data compiled by K. Morrison M. Econ. Geol. (based on Weighted Composite Mineral Potential information compiled for the Tasmanian Regional Forest Agreement).

Mineral deposit information derived from Mineral Resources Tasmania DEPOSITs data base. Data point position has not been verified in every case.

Digital base information from Land Information Services Division, Department of Primary Industries, Water and Environment.

Map produced by the Data Management Branch, Mineral Resources Tasmania using G.I.S. software.

While every care has been taken in the preparation of this data no warranty is given as to the correctness of the information and no liability is accepted for any statement or opinion or for any error or omission. No reader should act or fail to act on the basis of any material contained hereon. Readers should consult professional advisers. As a result the Crown in right of the State of Tasmania is not liable to any person in respect of any negligence or wrongful act or omission in any person whatsoever in respect of any data or content in this data by any such person in reliance whether in total or in part upon any of the material in this data. Crown copyright reserved.

THIS MAP SHOULD BE USED IN CONJUNCTION WITH MAP B – MINING LEASES AND ACTIVE MINES, PITS AND QUARRIES.

INDEX TO ADJOINING MAP SHEETS

**MUNICIPAL PLANNING INFORMATION SERIES
MAP 14A**