


GEOLOGY OF MACQUARIE ISLAND

TASMANIAN GEOLOGICAL SURVEY



LEGEND

- ALLUVIAL, LACUSTRINE AND SWAMP DEPOSITS**
- Alluvium, including deposits at the margin of lakes.
 - Older alluvium of terraces.
 - Colluvium of wash slopes.
 - Bogs, swamps and "featherbed" (includes significant areas of peat without bedrock outcrops; on the north-west coast these peat deposits cover palaeo-beach deposits on an elevated wave-cut platform 6-9m above sea level).
- SLOPE DEPOSITS**
- Scree slopes.
 - Alluvial fans, with slopes of less than 22 degrees.
 - Deposits of large (2-20m) boulders along coasts, typically in bays on west coast.
- BEACH AND AEOLIAN DEPOSITS**
- Pebbly to cobbly beach deposits.
 - Beach deposits with sand.
 - Coastal wind blown and dune sand.
 - Aeolian silt and sand deposits, often with gravel lag; approximate thickness of deposit in metres indicated where known.
- GLACIOGENE DEPOSITS**
- Deposits of grey clay with striated, faceted and polished cobbles and pebbles.
 - Lag deposits of striated, faceted and polished cobbles and boulders.
- SEDIMENTARY ROCKS**
- Mudstone and siltstone, usually laminated and red (grey at 490400mE 3952900mN).
 - Sandstone and pebbly sandstone with muddy to silty matrix; usually grey, greenish-grey or red.
 - Conglomerate, usually clast supported, consisting of sub-rounded to sub-angular cobbles to boulder-sized clasts of basalt and dolerite, in a mudstone to sandstone matrix. (Gabbro clasts in conglomerate at 490900mE 3956050mN).
- VOLCANICLASTIC ROCKS**
- Hyaloclastite breccia consisting of angular to sub-rounded blocks of usually aphyric basalt in a glass matrix.
 - Volcaniclastic breccia, matrix-supported, with blocks of usually aphyric basalt.
 - Volcaniclastic breccia, clast-supported, with blocks of usually aphyric basalt.
- LAVAS**
- Pillow basalt, aphyric to very sparsely phytic (<5% plagioclase phenocrysts), usually amygdaloidal.
 - Pillow basalt, sparsely to moderately phytic (5-30% plagioclase phenocrysts), usually amygdaloidal.
 - Pillow basalt, densely to very densely phytic (>30% plagioclase phenocrysts), usually amygdaloidal.
 - Tabular basalt flows, medium- to fine-grained, usually aphyric; rarely sparsely plagioclase-phyric or densely plagioclase-phyric. Rarely with zones of pillows.
 - Tabular basalt flows, medium- to coarse-grained with mesoscopically visible plagioclase laths, usually aphyric; rarely sparsely plagioclase-phyric. Rarely with zones of pillows. Autobrecciation locally present.
 - Hornblende-phyric massive tabular basalt flows, medium-grained. Rarely sparsely plagioclase-phyric. Rarely with mesoscopically visible plagioclase laths.
 - Small plugs of picrite. Picritic pillow basalts at (490400mE 3945700mN).
 - Olivine-bearing, matrix supported breccia at (490500mE 3945700mN).
- INTRUSIVE ROCKS**
- Sheeted dolerite dykes, with screens of massive, coarse-grained gabbro absent, minor (<5% or abundant (5-30%)) in Sandell Bay - Sellick Bay area, upper level sheeted dolerite dykes with rare screens of pillow basalt.
 - Sheeted dolerite dykes with minor to abundant screens of massive fine-grained gabbro. Proportion of sheeted dolerite dykes typically 50-80%.
 - Transition zone of subequal proportions of massive coarse-grained gabbro screens (30-70%) and sheeted dolerite dykes (30-70%).
 - Gabbro and olivine gabbro, usually massive; may contain dolerite dykes (0-30%) and rare ultramafic screens.
 - Gabbro and olivine gabbro with compositional layering, (Handspike Point, 492250mE 3961250mN).
 - Troctolite, usually with compositional layering; intruded by dolerite dykes.
 - Dominantly plagioclase-bearing massive whiteite and minor dunite, intruded by dolerite dykes.
 - Massive whiteite intrusion; plagioclase absent to very rare; without dolerite dykes (North Mawson, 494800mE 3955800mN).
 - Dunite and plagioclase-bearing dunite; massive; partly or completely serpentinized.
 - Dominantly harzburgite with subordinate dunite and plagioclase-bearing whiteite, intruded by dolerite dykes.
 - Harzburgite; usually massive or occasionally with mineral layering; partly or completely serpentinized. Proportion of dolerite dykes typically <10% but locally up to 30%.
- Geological boundary - position approximate.
Geological boundary - position inferred.
Fault.

GN

MN

Magnetic declination
29.6 deg (1991)

Scale 1:50 000



Projection: Universal Transverse Mercator - Zone 57
Horizontal datum - WGS1984

Geology by B.D.Goscombe, BSc (Hons), PhD and J.L.Everard, BSc (Hons), December 1994 - May 1995; September 1995 - January 1996. Project initiated and supervised by A.V.Brown, BSc (Hons), PhD, Director, Mineral Resources Tasmania, with funds provided by the Australian Antarctic Foundation, and logistical support provided by the Australian Antarctic Division.

Base map drawn from several sources:
The Spot multispectral satellite mosaic produced by the Australian Centre for Remote Sensing (ACRES) 1994.
Division of National Mapping Macquarie Island 1:50,000 topographic map (1971) warped to conform with the satellite mosaic along coastline and lakes.
Incomplete aerial photography flown in 1976 (mainly in the north of the island).
GPS positions and field observations.

Map produced by the Data Management Group, Mineral Resources Tasmania, using GIS software. Original map produced March 1998.
Absolute position with respect to horizontal datum and topographic features is approximate.

While every care has been taken in the preparation of this data, no warranty is given as to the correctness of the information and no liability is accepted for any statement or opinion or for any error or omission. No reader should act or fail to act on the basis of any material contained herein. Readers should consult professional advisers. As a result the Crown in Right of the State of Tasmania and its employees, contractors and agents expressly disclaim all and any liability (including all liability from or attributable to any negligent or wrongful act or omission) to any persons whatsoever in reliance whether in whole or in part upon any of the material in this data.
Crown copyright reserved.

LOCATION DIAGRAM

