

- 6 -

- B. Marine influence is restricted to the late Paleocene to Late Eocene part of the section but is never great until the top of the Demons Bluff Formation, as is usual within the basin.
- C. The cuttings are fairly clean, but significantly caved into the early to middle Eocene. Relatively few sidewall cores have been examined, and therefore some picks (especially base middle asperus, base asperopolus, base middle diversus and base upper balmei) may have been picked too low.
- D. Maturity shows a good overall trend, but the section is only mature for oil (not yet peak mature) and not yet mature for gas/condensate. Near 2080m, a local igneous event has produced high maturities over a limited part of the section.

REFERENCES

- Harris, W.K. (1985) Middle to Late Eocene Depositional Cycles and Dinoflagellate Zones in Southern Australia Spec. Publ., S. Aust. Dept. Mines and Energy 5 : 133-144
- Helby, R.J., Morgan, R.P., and Partridge, A.D., (in press) A palynological zonation of the Australia Mesozoic Australas. Assoc. Palaeontol. Mem. 3.
- Partridge, A.D. (1973) Revision of the Spore-pollen Zonation in the Bass Basin Esso unpubl. palaeo. rept. 1973/74
- Partridge, A.D. (1976) The geological Expression of Eustacy in the Early Tertiary of the Gippsland Basin Aust. Pet. Explor. Assoc. J., 16 : 73-79.
- Stover, L.E. and Evans, P.R. (1973) Upper Cretaceous - Eocene Spore Pollen Zonation, offshore Gippsland Basin, Australia Spec. Publ. geol. soc. Aust., 4 : 55-72
- Stover, L.E. and Partridge, A.D. (1973) Tertiary and Late Cretaceous Spores and Pollen from the Gippsland Basin, South-eastern Australia Proc. R. Soc. Vict., 85 : 237-286.