

REFERENCES

- HOPKINS, B., 1966. Geological Interpretation - Bass Basin. 8th. Commonwealth Mining & Metal Congress, vol.5; 217-221.
- HORNIBROOK, N. deB., 1961. Tertiary Foraminifera from the Oamaru District (N.Z.) - Part I - Systematics and distribution. N.Z. Geol. Surv. Paleont. Bull. 34 (1).
- JENKINS, D.G., 1965. Planktonic foraminiferal zones and new taxa from the Danian to lower Miocene of New Zealand. N.Z. J. Geol. & Geophys., 8 (6); 1088 - 1126.
- RAGGATT, H.G., & CRESPIAN, Irene, 1955. Stratigraphy of the Tertiary rocks between Torquay and Eastern View, Victoria. Proc. Roy. Soc. Vict., 67; 75-142.
- TAYLOR, D.J., 1965a. The mid-Tertiary Foraminiferal sequence Esso Bass-1 Well, Tasmania. Appendix to completion report - Esso Bass No.1 Well.
- TAYLOR, D.J., 1965b. Preservation, composition and significance of Victorian Lower Tertiary "Cyclammina faunas" Proc. Roy. Soc. Vict., 78(2); 143-160.
- TAYLOR, D.J., 1966. The mid-Tertiary foraminiferal sequence, Esso Gippsland Shelf No.1 Well. Appendix 1 in Petroleum Search Subsidy Acts Publications (in press).
- WADE, Mary, 1964. Application of the lineage concept to biostratigraphic zoning based on planktonic foraminifera. Micropaleontology, 10 (3); 273-290.

....