

SIN

BASS

DATE

March 30, 1973

334081

WELL NAME

BASS-2

ELEVATION

+31 feet

AGE	PALYNOLOGIC ZONES	HIGHEST DATA				LOWEST DATA					
		Preferred Depth	Rtg.	Alternate Depth	Rtg.	2 way time	Preferred Depth	Rtg.	Alternate Depth	Rtg.	2 way time
OLIG-MIO.	<u>P. tuberculatus</u>	3050	1				3050	1			
	<u>U. N. asperus</u>										
EOCENE	<u>M. N. asperus</u>	3800	1				3822	1			
	<u>L. N. asperus</u>	3992	1				4142				
	<u>P. asperopolus</u>	4308	2	4463	1		4463	1			
	<u>U. M. diversus</u>										
	<u>M. M. diversus</u>	4573	1				4648	1			
	<u>L. M. diversus</u>	4740	1				4861	2	4740	1	
	<u>U. L. balmei</u>	4939	1				4939	1			
PALEOCENE	<u>L. L. balmei</u>	5071	1				5508	1			
	<u>T. longus</u>										
LATE CRETACEOUS	<u>T. lilliei</u>										
	<u>N. senectus</u>										
	<u>C. trip./T.pach.</u>										
	<u>C. distocarin.</u>										
	<u>T. pannosus</u>										
EARLY CRETACEOUS											
PRE-CRETACEOUS											

COMMENTS: The Upper M. diversus Zone is absent in this well owing to a disconformity at about 4500 feet.

- RATINGS: 0; SWC or CORE, EXCELLENT CONFIDENCE, assemblage with zone species of spores, pollen and microplankton.
 1; SWC or CORE, GOOD CONFIDENCE, assemblage with zone species of spores and pollen or microplankton.
 2; SWC or CORE, POOR CONFIDENCE, assemblage with non-diagnostic spores, pollen and/or microplankton.
 3; CUTTINGS, FAIR CONFIDENCE, assemblage with zone species of either spore and pollen or microplankton, or both.
 4; CUTTINGS, NO CONFIDENCE, assemblage with non-diagnostic spores, pollen and/or microplankton.

NOTE: If a sample cannot be assigned to one particular zone, then no entry should be made. Also, if an entry is given a 3 or 4 confidence rating, an alternate depth with a better confidence rating should be entered, if possible.

DATA RECORDED BY: L.E. Stover & A.D. Partridge DATE June 1971

DATA REVISED BY: A.D. Partridge DATE March, 1973