

BASIN BASS DATE June 1971
 WELL NAME CORMORANT -1 ELEVATION +100'

AGE	PALYNOLOGIC ZONES	HIGHEST DATA				LOWEST DATA					
		Preferred Depth	Rtg.	Alternate Depth	Rtg.	2 way time	Preferred Depth	Rtg.	Alternate Depth	Rtg.	2 way time
O ^o -MIOC.	<u>T. bellus</u>										
	<u>P. tuberculatus</u>										
EOCENE	<u>U. N. asperus</u>										
	<u>L. N. asperus</u>	4304	1				4305	1			
	<u>P. asperopolus</u>	4978	1				4999	1			
	<u>U. M. diversus</u>	5356	1				7365	1			
	<u>L. M. diversus</u>	7635	1				9845	1			
EOCENE	<u>L. balmei</u>										
	<u>T. longus</u>										
LATE CRETACEOUS	<u>T. lilliei</u>										
	<u>N. senectus</u>										
	<u>C. trip./T.pach.</u>										
	<u>C. distocarin.</u>										
	<u>T. pannosus</u>										
	<u>C. paradoxa</u>										
EARLY CRETACEOUS	<u>C. striatus</u>										
	<u>U. C. hughesii</u>										
	<u>L. C. hughesii</u>										
	<u>C. stylosus</u>										
Pre-Cretaceous											

COMMENTS: Sample from 3815 feet definitely in N. asperus Zone, but spore-pollen assemblage too sparse to determine whether in upper or lower N. asperus

- RATINGS:
- 0; SWC or CORE, EXCELLENT CONFIDENCE, assemblage with zone species of spores, pollen and microplankton.
 - 1; SWC or CORE, GOOD CONFIDENCE, assemblage with zone species of spores and pollen or microplankton.
 - 2; SWC or CORE, POOR CONFIDENCE, assemblage with non-diagnostic spores, pollen and/or microplankton.
 - 3; CUTTINGS, FAIR CONFIDENCE, assemblage with zone species of either spores and pollen or microplankton, or both.
 - 4; CUTTINGS, NO CONFIDENCE, assemblage with non-diagnostic spores, pollen and/or microplankton.

NOTE: If a sample cannot be assigned to one particular zone, then no entry should be made. Also, if an entry is given a 3 or 4 confidence rating, an alternate depth with a better confidence rating should be entered, if possible.

DATE RECORDED BY: L.E. Stover / A.J. Partridge DATE June 1971
 DATA REVISED BY: _____ DATE _____