

BASIN BassDATE August 1973WELL NAME Dondu-1

ELEVATION _____

AGE	PALYNOLOGIC ZONES	HIGHEST DATA					LOWEST DATA				
		Preferred Depth	Rtg.	Alternate Depth	Rtg.	2 way time	Preferred Depth	Rtg.	Alternate Depth	Rtg.	2 way time
OLIG-MIO.	<u>P. tuberculatus</u>										
	<u>U. N. asperus</u>										
EOCENE	<u>M. N. asperus</u>	4840	0				5600	1			
	<u>L. N. asperus</u>	5700	1				6208	1			
	<u>P. asperopolus</u>	6530	1				6608	1			
	<u>U. M. diversus</u>	6700	2				6700	2			
	<u>M. M. diversus</u>	7075	2				7075	2			
	<u>L. M. diversus</u>	7378	1				7688	2	7510'	1	
PALEOCENE	<u>U. L. balmei</u>	7980	1	7674	2						
	<u>L. L. balmei</u>						9590	2			
	<u>T. longus</u>										
LATE CRETACEOUS	<u>T. lilliei</u>										
	<u>N. senectus</u>										
	<u>C. trip./T.pach.</u>										
	<u>C. distocarin.</u>										
	<u>T. pannosus</u>										
EARLY CRETACEOUS											
PRE-CRETACEOUS											

COMMENTS:

Samples from deeper than 7000 feet generally contain poorly preserved assemblages with very few diagnostic species.

- RATINGS: 0; SWC or CORE, EXCELLENT CONFIDENCE, assemblage with zone species of spores, pollen and microplankton.
 1; SWC or CORE, GOOD CONFIDENCE, assemblage with zone species of spores and pollen or microplankton.
 2; SWC or CORE, POOR CONFIDENCE, assemblage with non-diagnostic spores, pollen and/or microplankton.
 3; CUTTINGS, FAIR CONFIDENCE, assemblage with zone species of either spore and pollen or microplankton, or both.
 4; CUTTINGS, NO CONFIDENCE, assemblage with non-diagnostic spores, pollen and/or microplankton.

NOTE: If a sample cannot be assigned to one particular zone, then no entry should be made. Also, if an entry is given a 3 or 4 confidence rating, an alternate depth with a better confidence rating should be entered, if possible.

DATA RECORDED BY: L. E. StoverDATE August 1973