

1470-1480	50% <u>coquina</u> as above. 50% <u>calcarenite</u> ; as above, slight trace glauconite.
1480-1490	30% <u>coquina</u> ; as above. 70% <u>calcarenite</u> ; as above.
1490-1500	10% <u>coquina</u> ; as above. 90% <u>calcarenite</u> ; as above.
1500-1520	As above.
1520-1530	100% skeletal, silty, micritic, very fine grained, non porous, <u>calcarenite</u> ; light grey, medium soft-medium hard.
1530-1550	As above.
1550-1570	80% <u>calcarenite</u> ; as above, with trace glauconite. 20% skeletal, granular - micritic <u>limestone</u> ; light grey, hard, non porous.
1570-1580	80% <u>calcarenite</u> ; as above. 20% skeletal, granular-micritic <u>limestone</u> ; yellow-white, hard, abundant loose fossils in tray mainly bryozoans and foraminifera.
1580-1590	70% <u>calcarenite</u> ; as above. 20% <u>loose fossil</u> fragments as above. 10% <u>limestone</u> ; as above.
1590-1600	40% <u>calcarenite</u> ; as above, no glauconite, no pyrite. 10% <u>limestone</u> ; as above. 50% <u>clayey marl</u> ; medium light grey, very fossiliferous (as above.)
1600-1610	As above.
1610-1620	30% <u>calcarenite</u> ; as above. 10% skeletal, granular-micritic <u>limestone</u> as above. 60% <u>marl</u> ; as above.
1620-1630	40% <u>limestone</u> ; as above. 60% <u>marl</u> ; as above.
1630-1660	As above.
1660-1670	20% <u>limestone</u> ; as above. 80% <u>marl</u> ; as above, frequent loose unconsolidated fossils.
1670-1680	As above (amount of fossils decreasing).
1680-1770	As above.
1770-1780	10% <u>limestone</u> ; medium light grey, granular skeletal, very fine grain, hard. 90% <u>marl</u> ; as above.
1780-1880	As above.
1880-1900	20% <u>limestone</u> ; light grey and yellow-white, granular-micritic, and micritic, hard, non porous. 80% <u>marlstone</u> ; as above.
1900-1910	As above, abundant loose fossils, as above.
1910-1920	As above, marlstone, soft-firm.
1920-1950	As above.
1950-1960	30% <u>limestone</u> ; as above. 70% <u>marl</u> ; as above, loose fossils, as above, abundant.