

COMPANY: Golden Triangle NL
PROJECT: Main Creek
HOLE NUMBER: MC 40

Page No: 3

Description		Core Recovery			RQD			Assays											
From	To	From	To	%	From	To	%	From	To	MgO	CaO	SiO ₂	Fe ₂ O ₃						
119.0	130.5	MAGNESITE, pyritic (not assayed): gray, mottled largely crystalline magnesite, dolomitic and siliceous in part; upper and lower sections contain irregular areas of dark gray talcose schist (ie) gradational with units above and below; 119.0-123.0 m: pyritic, 3-5% medium-coarse euhedral pyrite as individual grains, thin veinlets and irregular semi massive patches; 123.0-130.0 m: 1-3% pyrite as disseminated grains and fine veinlets; lower contact marked by semi massive pyrite seam; core very competent; wide spaced joint sets 45, 60 CA;	119.0	130.5	100	119.0	130.5	100	84.0	85.0	46.24	1.24	<0.05	0.70					
										85.0	86.0	46.13	1.28	0.22	0.70				
											86.0	87.0	45.69	2.14	<0.05	0.67			
											87.0	88.0	45.06	2.80	<0.05	0.73			
											88.0	89.0	45.35	1.83	0.87	0.76			
											89.0	90.0	44.99	2.39	0.27	0.78			
											90.0	91.0	44.91	2.64	0.28	0.75			
											91.0	92.0	45.34	2.35	<0.05	0.77			
											92.0	93.0	45.14	2.31	<0.05	0.75			
											93.0	94.0	46.65	1.22	<0.05	0.69			
											94.0	95.0	45.82	1.49	<0.05	0.66			
											95.0	96.0	46.10	1.54	<0.05	0.73			
											96.0	97.0	45.89	1.70	<0.05	0.68			
											97.0	98.0	45.48	1.87	0.24	0.76			
130.5	147.8	INTER BEDDED SCHIST and MAGNESITE: interbedded dark gray pyritic, talcose schist and dolomitic, siliceous (?) magnesite; schistose margins of unit highly deformed and strongly pyritic; 130.5-132.0 m: dark gray strongly deformed schist with abundant pyrite and quartz veining; SCA 10-30; 132.0-141.0 m: more massive, less sheared dark gray, speckled schist (?volcanic); generally talcose with 1-3% pyrite as individual grains and concentrated in thin seams; SCA 30; joint set with slickensided surfaces 30 CA but striking 90 schistosity; wide spaced 2-5 mm quartz and quartz carbonate veins; 141.0-146.2 m: magnesite, light gray and dolomitic; mottled patches of coarse crystalline carbonate; band talcose schist near base; minor patches talc and disseminated 1-2% pyrite; 146.2-147.8 m: schist, pyritic and talcose; inclusions of magnesite; slumped and strongly sheared with abundant quartz veins and pyrite; unit very soft and broken; sharp contact with unit below 40 CA;	130.5	147.8	100	130.5	134.0	50	99.0	100.0	45.88	1.53	<0.05	0.64					
											100.0	101.0	46.30	1.32	<0.05	0.70			
												101.0	102.0	46.20	1.03	0.32	0.74		
												102.0	103.0	46.33	1.16	<0.05	0.77		
												103.0	104.0	46.15	1.36	<0.05	0.76		
												104.0	105.0	46.12	1.35	<0.05	0.68		
												105.0	106.0	44.95	2.68	<0.05	0.68		
												106.0	107.0	45.58	1.83	0.11	0.71		
												107.0	108.0	45.60	1.91	<0.05	0.72		
												108.0	109.0	46.40	1.27	0.12	0.82		
												109.0	110.0	45.68	1.59	0.13	0.89		
												110.0	111.0	45.56	1.77	0.21	0.94		
												111.0	112.0	45.90	1.39	0.11	1.01		
												112.0	113.0	45.71	1.54	0.20	1.07		
									113.0	114.0	45.78	1.59	0.36	1.06					
									114.0	115.0	45.47	1.65	0.98	1.21					
									115.0	116.0	44.67	2.22	0.71	1.45					
									116.0	117.6	33.33	14.95	2.14	1.91					