



McKIMMIE CREEK EL 34/2004

**ANNUAL REPORT
FOR THE PERIOD ENDING 30th JULY 2005**

Author: A. W. McNeill

Date: 7 September 2005

Submitted To: MTS Superintendent – Zinifex Rosebery Mine

Copies To: Mineral Resources Tasmania
Zinifex Rosebery Mine

Submitted By:

Accepted By:

Rosebery Report No: R83

CONTENTS

3404_200508_02_MainReport

1. SUMMARY	1
2. INTRODUCTION	2
2.1 Attribution	2
3. LAND TENURE	3
4. GEOLOGY	3
5. PREVIOUS EXPLORATION.....	4
6. WORK COMPLETED 2004-2005 REPORTING PERIOD	6
6.1 Previous Exploration Data.....	6
7. CONCLUSIONS & RECOMMENDATIONS.....	6
8. EXPENDITURE.....	7
9. KEYWORDS & LOCALITY	8
10. REFERENCES	8

LIST OF TABLES

Table 1	Previous exploration over the area of AEM anomaly D1
Table 2	Previous exploration over the area of AEM anomaly D2

LIST OF FIGURES

Figure No.	Title	Scale
<i>3404_200508_03_fig1.pdf</i>	Tenement Location Diagram	1:500,000
<i>3404_200508_04_fig2.pdf</i>	Regional Geology	1:35,000

LIST OF APPENDICES

<i>3404_200508_05_App1a.txt</i>	Previous exploration data (digital copy only)
---------------------------------	---

1. SUMMARY

This report details exploration work undertaken during the first year of EL 34/2004 McKimmie Creek.

Work was restricted to commencing a review of previous exploration. It is recommended that gridding and soil sampling of AEM anomalies D1&D2 be completed in year 2 of the licence.

2. INTRODUCTION

This report details exploration work undertaken on McKimmie Creek EL 34/2004 during the first year of this tenement.

Zinifex's main target on EL 34/2004 is Devonian Pb-Zn vein mineralisation, with Cambrian VHMS systems as a more 'speculative' target, of sufficient tonnage and grade that can be used as mill feed to supplement current underground resources at the Rosebery Mine.

Zinifex plans to systematically explore the EL using a combination of geological mapping (where possible), partial leach soil geochemistry and infill time-domain ground EM, where there is no existing coverage, or the work that has been done is considered to have been ineffective.

Although close to the town of Rosebery, access to the tenement is not as easy as one would think. The sole vehicular access to the tenement is an 8.1 km long 4WD track (opened –up by Comstaff in the late 1970's) that runs roughly through the centre of the tenement and originates from the Pieman Road (and associated High Voltage powerline access tracks) west of the Bastyan Dam. Several Grids were established by Comstaff between 1973 and 1985 but are now too overgrown to be of any practical use.

2.1 Attribution

The following personnel were responsible for the work carried out by Zinifex Rosebery Mine on the EL 34/2004 McKimmie Creek licence area during the reporting period:

Senior Geologist: Andrew McNeill – Zinifex Rosebery Mine

3. LAND TENURE

EL 34/2004 McKimmie Creek (17 sq km) was granted to Zinifex on 30 July 2004 for a period of 5 years. The location of the tenement is shown on Figure 1. EL 34/2004 covers ground that fell vacant on the relinquishment of EL 21/96 (Pasmaico) in February 2001, EL 29/91 (Golden Reef Enterprises) in 1996 and EL 12/194 (Bruce Resources NL) in 1995.

Land covered by EL 34/2004 is a mixture of crown land designated as Multiple Use State Forest, Unallocated Crown Land, private property, part of the Mt Kershaw Regional Reserve, MDC informal reserve and some HEC land all of which are available for exploration under the Mineral Resources Development Act 1995. The licence area excludes approximately 3.5 sq km of ML 28M/1993, which impinges on the eastern boundary of McKimmie Creek.

4. GEOLOGY

The regional geology of the tenement area is shown on Figure 2, which is largely derived from Corbett and McNeill (1986 and 1987) and Corbett (2002).

Much of the bedrock geology of EL 34/2004 McKimmie creek is obscured by a variable thickness of Quaternary glacial cover (approximately 75% of the area of the licence). The bedrock geology of the area can be divided into three domains, separated by major north-south regional faults:

1. The Mount Read Volcanics (MRV)
2. The “Rosebery Group” or Dundas Group correlates (as defined by Corbett and Lees, 1987)
3. Correlates of the Crimson Creek Formation or the ‘Cleveland-Waratah association’

The Mount Read Volcanics, pumice breccias and intrusives forming part of the CVC, occur in the far southeastern part of the licence area, lying above of the Rosebery fault, a moderately east dipping thrust. The MRV fall in the excluded part of the licence, within the Rosebery Mine Lease ML 28M/93.

West of the Rosebery Fault lies a predominantly sedimentary sequence that has been termed the ‘Rosebery Group’. The understanding and correlation of the ‘Rosebery Group’ has undergone considerable re-interpretation since the early work by Hills in 1914 and this evolution is summarised in Green (1983) and Corbett and Lees (1987).

It is now generally agreed that the Chamberlain shale, largely outcropping in the eastward bulge of the Rosebery Fault, immediately west of the Rosebery ore body, is a west facing correlate of the White Spur Formation (i.e., the Rosebery Hangingwall sequence). The Chamberlain shale is conformably overlain by the Stitt Quartzite (including the Munro Creek formation of Green [1983]), a prominent marker unit that has fossils indicating correlation with the Cambro-Ordovician Owen Group (Corbett, 2003). The Stitt Quartzite is in turn overlain by the Westcott Argillite, a more carbonate-rich unit. Several other units, in particular the Natone Volcanics, calc-

alkaline ‘Mount Read’-type volcanics, the Salisbury Conglomerate and gabbroic bodies, have unclear stratigraphic relations to the units described above; Corbett (2002, 2003) concludes that they may be older than the Owen Group correlates (Westcott Argillite, Stitt Quartzite) due to ‘structural complexities’, whilst Parfrey (1993) argues that the Salisbury Conglomerate is part of the Westcott Argillite and that as the Salisbury Conglomerate overlies and interfingers with the Natone Volcanics (in drill hole Rosebery 1) they are also equivalent to the Westcott Argillite (implying a Owen Group age if Corbett’s correlations are correct).

The ‘Rosebery Group’ is bounded to the west by a major north-south fault, the ‘Marion Oak Fault’. To the west of this structure is a sequence of mafic greywacke and siltstone with minor tholeiitic basalt lavas that extend from south of Colebrook Hill north to the Huskisson River. These rocks were mapped, following previous workers, as Crimson Creek Formation by Corbett and McNeill (1986), however, these lithologies are now correlated with the allochthonous early-Cambrian Cleveland-Waratah Association (Corbett, 2002 and 2004).

Two mineralised zones are known from the licence area; Salmons Lode, described by A. McIntosh Reid (1918), and minor vein style Pb-Zn mineralisation at Shell’s Bastyan Dam grid (Smyth, 1983), both of which occur in lithologies mapped as Westcott Argillite by MRT (a third mineral occurrence is recorded by Green and Bamford (1986), in the Pieman River downstream of the Bastyan Dam, but little information is available on this prospect).

5. PREVIOUS EXPLORATION

The area of EL 34/2004 McKimmie Creek has a long history of ‘modern’ exploration commencing in the 1970s, as part of EL 5/1963. In the first year of tenure previous exploration over airborne EM anomalies D1 and D2, located by Pasmaenco (Briggs and McNeill, 2001), has been reviewed and is summarised in Tables 1 and 2, respectively.

Table 1 Previous exploration over the area of Airborne EM anomaly D1

Reporting Period	Work Completed
1972 (Piggott, 1972)	Area covered by –80# and panned concentrate stream sediment sampling and geological mapping of creeks. Work established a geological framework for the area and located low-order stream sediment anomalies.
1973-1974 (Orr, 1974)	Stream sediment sampling was considered to have been ‘inadequate’ and soil sampling on a large grid was completed; Anomaly D1 lies just east of the sampled area. Recommend an airborne EM survey as the best way to test the area.
1975 (Butt, et al., 1975)	An Input EM survey was completed. Data was interpreted and several anomalies located, including anomaly CS10 a ‘fair’ bedrock conductor that is roughly coincident with the Pasmaenco D1 anomaly. No follow-up work appears to have been completed.
1983 (Dvorak, Z.)	A DIGHEM III survey was flown over the entire area of interest. Anomaly 190H was located in the area of interest and was recommended for ground follow-up.
1984 (Trussell, 1984)	The DIGHEM data were further reviewed and the 190H area recommended for ground EM follow-up, as it appeared to be a ‘thick conductor’. This work does not appear to have been done.

Table 1 Previous exploration over the area of Airborne EM anomaly D1, cont..

Reporting Period	Work Completed
1998-1999 Parfrey and Simpson (1999)	Identification of priority prospect areas through the completion of an airborne EM Survey. Several discrete anomalous responses, including anomaly D1 were identified - these were considered worthy of further investigation.
2000-2001 Briggs and McNeill (2001)	Detailed interpretation of 1999 Airborne EM survey; – 6 anomalies, including D1 were recommended for follow-up, however this work was not completed.

Table 2 Previous exploration over the area of airborne EM anomaly D2

Reporting Period	Work Completed
1972 (Piggott, 1972)	Area covered by –80# and panned concentrate stream sediment sampling and geological mapping of creeks. Work established a geological framework for the area and located low-order stream sediment anomalies.
1973-1974 (Orr, 1974)	Stream sediment sampling was considered to have been ‘inadequate’ and soil sampling on a large grid was completed; this grid did not cover the ‘marshy’ area overlying anomaly D2. Recommend an airborne EM survey as the best way to test the area.
1975 (Butt, et al., 1975)	An Input EM survey was completed over the entire East Renison Block. Data was interpreted and several anomalies located, including the GAO anomaly that is roughly coincident with the Pasmenco D2 anomaly.
1979 (Hall, 1979)	The GAO anomaly area was gridded (3.97km), geologically mapped, soil sampled and ground magnetic and EM (Crone shootback) surveys completed. It was concluded that there were no positive results and no further work was warranted.
1983 (Dvorak, Z.)	A DIGHEM III survey was flown over the entire area of interest. Anomalies 170F-180H and 190K were located in the area of interest and were recommended for ground follow-up.
1984 (Trussell, 1984)	The DIGHEM data were further reviewed and the 170F-180H, 190K area (Pasmenco anomaly D2) recommended for ground EM follow-up.
1984-1985 (Shaw and Everett, 1985; Everett, 1985)	A 5 line km grid was established, soil sampled on the southern two lines (as most grid overlaps GAO), ground magnetic and GENIE EM surveys completed and two diamond drill holes (180H/1 and 180H/2) completed for a total of 218.5m. Both holes intersected black carbonaceous shales with little geochemical encouragement and no further work was recommended.
1998-1999 Parfrey and Simpson (1999)	Identification of priority prospect areas through the completion of an airborne EM Survey. Several discrete anomalous responses, including anomaly D1 were identified - these were considered worthy of further investigation.
2000-2001 Briggs and McNeill (2001)	Detailed interpretation of 1999 Airborne EM survey; – 6 anomalies, were recommended for follow-up. Anomaly D2 was not recommended for follow-up on the basis of the lack of geochemical response in DDH 180H/1 & 2.

6. WORK COMPLETED 2004-2005 REPORTING PERIOD

Work completed in the current reporting period was limited to a review of previous exploration over the two known EM anomalies.

6.1 Previous Exploration Data

Previous exploration data from the GAO/180H grids (covering the D2 anomaly) was compiled. A0 horizon soil sampling for the GAO grid and results from diamond drill holes 180H1 and 180H2 were digitally compiled and are included in Appendix 1 (digital copy only). Only line profiles, and not primary data, were available for the C Horizon soil sampling on the GAO grid as is the case for the two lines of the 180H grid sampled in 1984/1985 (Shaw and Everett, 1985; however, refer to a report by Thynne [1984] which may include the raw data. Unfortunately this report does not appear to be in the MRT library).

Compilation of data from Shell's Bastyan Dam grid commenced and drill hole data are included in Appendix 1 (digital copy only).

Data compilation is still in process and all outstanding data will be included as part of the next annual report.

7. CONCLUSIONS & RECOMMENDATIONS

Due to resource constraints, work during the first year of EL 34/2004 McKimmie Creek was limited to the compilation of previous exploration data. During the second year of the licence it is recommended that:

- Data compilation, particularly from the northern part of the tenement, be completed.
- The old Comstaff access road be re-opened and small grids be established over Pasmenco AEM anomalies D1 and D2. These grids should be partial leach soil sampled, and, should there be sufficient encouragement, be covered by ground EM.

8. EXPENDITURE

Expenditure on EL 34/2004 McKimmie Creek during the 12 month period ending 30 July 2005 was **\$2,328**. A detailed breakdown of this expenditure is presented below.

Personnel	\$
Travel & Accommodation	\$
Consultants & Contractors	\$
Geological Consultants	\$
Geochemical Consultants & Assays	\$
Geophysical Surveys & Contractors	\$
Drilling	\$0
Stores & Supplies	\$188
Vehicles Plant & Equipment	\$
Land	\$233
Computing	\$1,907
Office	\$
Administration Fee	\$
Total Tenement Expenditure	\$2,328

9. KEYWORDS & LOCALITY

Keywords

Previous Exploration, GAO grid, 180H grid, geophysics – Airborne EM, Renison East.

Locality

1:250,000	QUEENSTOWN SK55-5
1:100,000	SOPHIA 8014; PIEMAN 7914
1:25,000	ROSEBERY 3637

10. REFERENCES

- Briggs, T., and McNeill, A.W., 2001. Dundas EL 21/96 Final Report for the period 9/10/00 to 8/10/01. Unpub. Pasminco Exploration Report D19 (**TCR01-4607**).
- Butt, G.R., Finney, W., Hansen, J., 1975. Interpretation Report Airborne Electromagnetic (INPUT) Survey of the Comstaff South Leases Tasmania, Australia. Unpub. Geoterrex report 83-134 (**TCR85-2397**).
- Corbett, K.D., 2002. Updating the geology of the Mount Read Volcanics Belt; WTRMP Mt. Read Volcanics Compilation. Tasmania Geological Survey Record 2002/19, Mineral Resources Tasmania, Hobart.
- Corbett, K.D., 2004. Updating and revision of the 1:25 000 scale series geological maps covering the Mt Read Volcanics belt in western and northwestern Tasmania. Tasmania Geological Survey Record 2004/03, Mineral Resources Tasmania, Hobart.
- Corbett, K.D., and Lees, T.J., 1987. Stratigraphic and structural relationships and evidence for Cambrian deformation at the western margin of the Mount Read Volcanics, Tasmania. *Aust. J. Earth Sci.*, 34:45-68.
- Corbett, K.D., and McNeill, A.W., 1986. Geological compilation map of the Mount Read Volcanics and associated rocks, Hellyer to South Darwin Peak. MRVP Map 6. Mineral Resources Tasmania.
- Corbett, K.D., and McNeill, A.W., 1987. Geology of the Rosebery – Mt Block area. MRVP Map 2. Mineral Resources Tasmania.
- Dvorak, Z., 1983. DIGHEM III Survey of the East Chester North Pieman and Arthur River Areas, Tasmania. Unpub. Comstaff Pty Ltd. Report (**TCR83-2129**).
- Everett, M.P., 1985. East Renison (EL 5/63), Part 6, Interim Report. Unpub. Comstaff Pty. Ltd. Report (**TCR85-2419**).

- Green, G.R., 1983. The Geological setting and formation of the Rosebery volcanic-hosted massive sulphide orebody, Tasmania. Unpub. PhD. Thesis, University of Tasmania, Hobart.
- Green, G.R., and Bamford, A., 1986. Tullah, Metallic Mineral Deposits Map Series Sheet 8014-IV, Geological Survey of Tasmania.
- Hall, D.B., 1979. Pieman Area (Input Anomaly GAO), 1979 Report on Work Completed. Unpub. Comstaff Pty Ltd. Report 9709 (**TCR79-1363**).
- Lees, T.C., 1987. Geology and mineralisation of the Rosebery-Hercules area, Tasmania. Unpub. MSc. Thesis, University of Tasmania, Hobart.
- McIntosh Reid, A., 1918. The North Pieman and Huskisson and Sterling Valley mining fields. Geological Survey Bulletin No. 28, Tasmania Department of Mines, Hobart.
- Parfrey, O., 1993. Mesoscopic and microscopic structural evolution in the Dundas Group at Rosebery, western Tasmania. Unpub. BSc (Hons.) Thesis, University of Melbourne.
- Parfrey, O.C., and Simpson, K.L., 1999. Dundas EL 21/96 Annual report for the period November 1998 to November 1999. Unpub. Pasmenco Exploration report VC284 (**TCR00-4413**).
- Shaw, R.W.L., and Everett, M.P., 1985. Final Report on Areas surrendered to the Department of Mines, Tasmania, Exploration Licence 5/63 Area 6 East Renison. Unpub. Comstaff Pty Ltd. Report (**TCR 85-2418**).
- Smyth, W.D., 1983. EL 22/74 Marionoak, annual report on exploration, June 1982-June 1983. Unpub. Shell Company of Australia Limited report 08.2063 (**TCR83-2044**).
- Trussell, D.B., 1984. Interpretation of the Renison East area DIGHEM survey. Unpub. Comstaff Pty Ltd. Report (**TCR84-2134**).
- Orr, D.B., 1974. Pieman South Summer Field Season 1973/1974. Unpub. Comstaff Pty Ltd. Report (**TCR74-1023**).
- Piggott, G.F., 1972. Exploration Licence 5/63, 1971/1972 Summer Field Season Report, Pieman Area. Unpub. Comstaff Pty Ltd Report (**TCR72-0850**).