



EXPLORATION LICENCE  
EL 35/2008  
MANGANA, NORTH EAST TASMANIA

ANNUAL REPORT  
FOR THE YEAR ENDED  
18 NOVEMBER 2009

Prepared by:  
John Richardson  
Consulting Geologist  
Macquarie Harbour Mining Limited

Submitted by:  
Frank Rogers  
Managing Director  
Macquarie Harbour Mining Limited

Macquarie Harbour Mining Limited ABN 124 212 175  
52 Channel Highway, Kingston, Tasmania, 7050  
TELEPHONE: 03 6229 9955 FACSIMILE: 03 6229 8430  
EMAIL: [info@mhml.com.au](mailto:info@mhml.com.au) WEBSITE: [www.mhml.com.au](http://www.mhml.com.au)

## 1 EXECUTIVE SUMMARY

As the result of a successful tender application for ERA 699 the company was granted EL35/2008 which covers the Mangana Goldfield. Exploration is directed towards economic gold mineralisation forming extensions of, or proximal to, areas of previous historical production.

Work conducted in the first year of tenure has mainly comprised literature research and database compilation. Several field trips were made to the tenement in order to gain an understanding of terrain and access limitations, and to investigate the nature, size and distribution of the old workings.

## CONTENTS

	Page
1 EXECUTIVE SUMMARY	1
2 INTRODUCTION	3
3 REVIEW OF PREVIOUS MINING AND EXPLORATION	6
4 EXPLORATION COMPLETED DURING THE REPORT PERIOD	9
5 REFERENCES	11
6 EXPENDITURE	13

## FIGURES

Figure 1	Tenement location map	4
Figure 2	Geology of EL35/2008	5
Figure 3	Historic prospects near Mangana	10

## 2 INTRODUCTION

This is the first annual report on EL35/2008 which is held in the name of Goldstock Mining Pty Ltd, a wholly owned subsidiary of Macquarie Harbour Mining ("MHM"). The tenement (see Fig.1) covers an area of 178 sq km and lies immediately northwest of the town of Fingal in NE Tasmania. The old gold mining centre of Mangana, which is at the southern end of the Mangana – Waterhouse "gold corridor", falls in the southern part of the licence. RL14/1988 (17 sq km), which is held by The Cornwall Coal Mining Company Pty Ltd, covers Tower Hill, a prominent landmark north of Mangana. This retention licence, and a small mining lease at Bucklands Hill (1769P/M) are excluded from EL35/2008.

Most of the tenement lies in strongly dissected hill country designated as State Forest. Access in the eastern half is facilitated by a network of logging roads which are less widespread in the west. Private ownership is mainly confined to the lower lying areas in the south and east.

Surface geology (see Fig. 2) is dominated by the Mathinna Group turbidite sequence of sandstone and siltstone which hosts the widespread gold mineralisation in NE Tasmania. The topography is dominated by two landforms in excess of 1000 metres, Tower Hill and Byatt's Razorback, both of which are capped by Jurassic dolerite with associated talus. Underlying are sub-horizontal Triassic sandstones and Permian sandstones, siltstones and mudstones. Quaternary alluvium fills the valley floors.

Three styles of gold mineralisation exist in the Mangana region; sheeted vein systems, stockworks and alluvial deposits. The sheeted veins tend to be sub-vertical and occur parallel to bedding within slaty sequences of the Mathinna Beds. They are narrow (0.05 – 1.5m) and have no surrounding alteration haloes. Almost all the hard rock production has been of this style. Quartz stockworks are hosted in the Mathinna Beds that have undergone brittle deformation. They generally appear to be associated with larger sheeted vein systems and can be seen at Bucklands Hill and the Tower Hill Goldfield. Alluvial gold at Mangana is hosted in a clayey cobble wash in Major's Gully, Sharkeys Gully, Richardson's Creek and the creek flats surrounding Tower Hill Rivulet.

### 3 REVIEW OF PREVIOUS MINING AND EXPLORATION

The Mangana goldfield was the first discovery of payable gold in Tasmania. Most production occurred between 1852 and 1910 with the first lode mining commencing at the Sovereign Mine in 1859 while some alluvial mining was still taking place at Majors Gully in the 1990's. The total recorded production from lode deposits in the area is 208kg but actual production was probably much greater. Between 5,000 and 15,000oz (160 – 470kg) of gold was also produced from alluvial deposits. The area is highly mineralised with more than 30 known deposits close to the township itself. It forms a NNW trending corridor approximately 6km long and 2km wide where the gold occurs in structurally (fault) controlled veins and breccia zones within sandstone and siltstone and slate. The veins are typically small and erratic in grade although it is considered that the Golden Entrance reef may extend for 2km. This reef was the largest single producer with 91kg (2940oz) at an average grade of 127g/t Au. This is small in comparison to Golden Gate (253,856oz) which lies immediately north of EL35/2008.

At Tower Hill, 8km to the north of Mangana and considered a separate goldfield, the only recorded production was from the Sunbeam Mine (12t of ore for 0.12kg Au).

#### Recent Exploration

Like other small goldfields in NE Tasmania Mangana and Tower Hill experienced a number of decades of inactivity following the initial production period. The last 25 years have seen modern exploration by a number of companies over wide areas; the work carried out on EL55/1983 (Tasmanian Alluvials, Alcaston Mining NL and Pegasus Gold Australia Ltd), EL18/1991 (Resolute Samantha Ltd, Defiance Mining NL) and EL22/1992 (Newcrest Mining Ltd) is the most relevant.

EL 55/1983:

. 1984-85 (CR 85-2510) – Seventeen backhoe holes were dug in Majors Gully, with visible gold in 10 of the 17 pan concentrates, and the slate bottom found to be less than 3m deep. Three 20 litre drums of tailings from the Mangana reef averaged 1.69 g/t Au (note that in 1994 Mancala Pty Ltd estimated the Mangana tailings to contain 20,000t at 0.83 g/t Au).

.1985-86 (CR 86-2607) – Bulk testing of alluvials in Majors Gully delineated a resource of 300,632 cu metres at 0.307 g/cu.m Au.

.1987-88 (CR 88-2883) – Work concentrated on the Argyle Reef, a south-eastern extension of the Golden Entrance Reef about 1.5 km SE of Mangana. Access to the reef was through three drives, two from the NW end and one from the SE, and Alcaston stated in this report that limited mining took place in only one of the drives (Argyle No1 South) with the ore mined being less than 500 tonnes. Earlier sampling (Tasminex NL, 1981) in Argyle No1 South produced grades from 17 rock chip samples ranging from 4-133 g/t (mean 31.2 g/t) with the vein width over the interval sampled ranging from 10-90 cm (average 44 cm).

In 1988 Alcaston tested the Argyle Reef, adjacent wall rocks and two minor parallel quartz reefs, with eight shallow (31m) angled open hole percussion holes and 38 composite rock chip samples taken from inside the old drives. Although significant (0.1 – 0.4ppm) vein quartz was encountered in all holes except one there were no ore grade

intersections. Background arsenic values in the drill holes varied from 10 to 200ppm but higher arsenic did not necessarily correlate with high gold values.

Four anomalies were found in the rock chip sampling, two from Argyle No.1 South adit (53 metres in) recorded 10.02 and 9.03 g/t, and two from the Argyle No. 2 South adit (21 metres in) were 4.93 and 3.15 g/t. The remaining 34 samples ranged from 0.026 – 0.759 g/t, again with no direct arsenic correlation.

.1988-89 (CR 89 – 3052) Argyle: Resampling and assay of the bulk samples from the previous year's drilling came up with similar results. It is of interest that AA7 is the only hole drilled below the water table which returned anomalous values (10x background). These occurred over a true thickness of approximately 5 metres within a pyritic black shale. On the other hand the samples from the oxide profile have narrower, often high grade sections.

Tower Hill: The quartz vein stockwork target was assessed in two outcropping sub-parallel sandstone units. This consisted of cutting two grids over the prospective sandstones, mapping the geology and old workings at 1:1000, composite rock chip sampling in prospective locations and soil sampling using a hand auger. A ground magnetic survey was also conducted.

Results from the rock chip sampling show that only 17 of the 68 rock chip samples produced grades of >0.1 g/t Au and 16 (8 of which were > 1g/t Au) of these came from dumps around the mine shafts and old exploration trenches. The soil sampling indicates a group of anomalous gold values (>20ppb) trending approximately magnetic E-W to the south of the Tower Hill – Sunbeam workings.

RC drilling was recommended to follow up at Tower Hill.

.1989-90 (CR 90-3197) A helicopter aeromagnetic survey was flown with 150m spacing and an 80m terrain clearance. This indicated "systematic but subtle variations" in the magnetic field across the Mathinna Beds. It was also found to correlate with the previous year's ground survey at Tower Hill.

A programme of RC drilling was completed in the Tower Hill – Sunbeam area (13 holes totalling 357m). These holes were aimed at a combination of geochemical, structural and magnetic targets and encountered minor gold mineralisation. Best intersections were 1m @ 3.11g/t Au from 43-44m and 3m @ 2.87g/t Au, both from Sunbeam.

EL 18/1991:

.1993-94 Alec White (CR 94-3564) Thirty rock chip samples from the Golden Entrance/Mangana area analysed for gold with best result being 1.37 ppm. Alluvial testing at Sharkey's Gully, recoverable grade of 0.096 g/cu.yd reported.

.1995-96 Resolute Samantha Ltd. (CR 96-3842) Work consisted of gridded soil sampling in the Golden Entrance/Mangana Gold Reefs/Fingal area east of the town, at Buckland Hill and at Blackboy Ridge. The area of old workings at Mangana was also geologically mapped and 64 rock samples collected. Useful historical production information is provided in a set of plans and sections. Results of the rock sampling indicate strong variability in grades and the mapping confirmed a general narrow reef width (< 0.5 metres the norm). At Buckland Hill a zone of stockworked quartz gave poor results in

rock sampling with a single 1.52 g/t Au sample not supported by further sampling. A significant soil anomaly (peak value 195 ppb Au, 570 ppm As) was delineated on Blackboy Ridge. Enhancement of aeromagnetics highlighted a major NNW trending structure passing through the Fingal Mine and continuing through the zone of Au and As anomalism on Blackboy Ridge.

It was concluded that further work should be aimed at drill testing the reef systems east of the town with one targeting the north end of No 4 adit at Mangana Gold Reefs and follow-up at Blackboy Ridge.

.1997-98 (CR 98-4212) Incorporated into EL 3/97. No field work by new operators Connemara GM Pty Ltd. Proposals for drilling at Buckland (two 100m RC holes), Mangana (one 200m diamond hole) and Blackboy Ridge (four fences of 80m RC holes).

.1998-1999 (CR 99-4321) Operator now Defiance Mining NL. Compilation of geochemical data into a GIS, processing of existing aeromagnetic data. Drilling proposed for Argyle/Golden Entrance and Blackboy Ridge areas.

.1999-2000 (CR 00-4438) Work comprised two declined percussion drill holes (total 180m) into the Golden Entrance Reef and follow-up soil sampling at Blackboy Ridge. While a number of quartz veins were intersected in both holes at Golden Entrance the sulphide content was very low and assays were disappointing, with the best intersection being 1m from 43m at 0.5g/t Au. The work at Blackboy Ridge downgraded the anomaly.

.2000-01 (CR 01-4573) No field work carried out during this year. It was proposed that 6-8 RC holes should be drilled to test the southern end of the Golden Entrance Reef/Argyle Reef between Sailor's Gully and Irvine's Gully.

There are no further reports on EL18/91 so presumably the licence was relinquished.

EL 22/1992:

.1992-1993 (CR 93-3498) Newcrest's targets were large volumes of quartz stockwork gold with grades > 1.5 g/t Au. Their EL excluded the main part of the Mangana and Golden Entrance reefs.

Soil and outcrop sampling at Tower Hill failed to identify significant gold mineralisation. Work then concentrated on the Golden Gully – Blackboy Ridge area some 3 km to the north and along strike from the Mangana/Golden entrance mines and upstream from the Majors Creek alluvials. Initial drainage results (up to 143ppb Au from Golden Gully) led to ridge and spur sampling which showed quartz stockworked sandstone occurs extensively throughout the catchment. The only significant value (1.1ppm) was sourced from a narrow (<0.5m quartz matrix breccia vein) of limited extent.

Newcrest concluded that although quartz vein stockwork zones that been demonstrated it was apparent that much of it was unmineralised.

#### 4 EXPLORATION COMPLETED DURING THE REPORT PERIOD

A literature review was conducted of the area covered by the Exploration Licence, with emphasis on “open file” reports compiled by exploration companies.

Two visits were made to EL35/2008 during the year with the aim of traversing as much of the tenement as possible to examine ease of access for drilling and gain a good understanding of surface geology and the nature of mineralisation in the old workings. The most pertinent observations are summarised below.

- The area is steep and forested meaning access, particularly for drilling equipment, will be difficult.
- Numerous open shafts, difficult terrain, fire in summer, and falling trees in high wind all present significant safety risks.
- A number of land titles, particularly close to Mangana, exist in the area and some exploration activities may prove problematic.
- Lack of phone coverage makes setting up a remote office/caravan a more difficult prospect.
- All hardrock mines have worked isolated narrow vein or reef systems. With the exception of Bucklands Hill no rock was observed with quartz stockwork.
- Alluvial deposits in the area are very unlikely to be economic.

Conclusions drawn from the data review and field reconnaissance indicate that alluvial deposits are very unlikely to be economic and very little potential exists for disseminated bulk mineable mineralisation. The area however is highly prospective for high grade concentrated vein systems of a similar nature to the Golden Gate mine 10km to the north of Mangana. Of particular interest is a narrow northwest trending zone that extends from Argyle in the south through the Golden Entrance, Underlay, Fingal and Golden Gully Mines to the Tower Hill Freehold in the north. These old workings are shown in Figure 3.

## REFERENCES

- Akerman, T.E. 1997. Partial Surrender Mangana-Alberton – EL23/92. Mancala Pty Ltd. MRT Ref. 97\_3996
- Akerman, T.E. 1998. Annual Report Sep 1996 to Sep 1997 – EL23/92. Mancala Pty Ltd. MRT 98\_4110
- Akerman, T.E. 1998. Annual Report Sep 1997 to Sep 1998 – EL23/92. Hercules Resources Pty Ltd. MRT Ref. 98\_4221
- Anon. 2006. Report on Alluvial Exploration Program. Tasmanian Alluvials on EL55/1983 at Mangana for Sept Oct 1986. MRT Ref. 06\_5362
- Anon. 1986. Annual Report on EL55/83. Tasmanian Alluvials. MRT Ref. 86\_2607
- Ashley, J. 1995. Project Interpretation of Aeromagnetic Data – EL27/94 Mathinna. MRT Ref. 95\_3791A
- Bottrill, R.S. 1992. The Mangana Goldfield and adjacent Gold mining areas. Tasmania Dept of Mines. MRT Report 1992/29.
- Charlton, E.N. 1980. Report for the six month period ending 30<sup>th</sup> Apr 1980 – EL17/78. Tasminex NL. MRT Ref. 80\_1428
- Charlton, E.N. 1980. Summary of Completed Work Programme for the period 1<sup>st</sup> May 1980 to Oct 31<sup>st</sup>, 1980 and Proposed Work Programme for the period Nov 1<sup>st</sup> 1980 to Apr 30<sup>th</sup> 1981 – EL 17/78. Tasminex NL. MRT Ref 80\_1502
- Charlton, E.N. 1981. Report covering the period from Nov 1980 to May 1981 – EL17/78. Tasminex NL. MRT Ref 81\_1551
- Charlton, E.N., Daly, A.B. 1981. Report covering the period from 1<sup>st</sup> May 1981 to 30<sup>th</sup> Oct 1981 – EL17/78. Tasminex NL. MRT Ref. 81\_1642
- Charlton, E.N., Daly, A.B. 1982. Report for the six month period 1<sup>st</sup> Nov 1981 – 30<sup>th</sup> Apr 1982. Tasminex NL. MRT Ref. 82\_1760
- Colville, R. 1998. Annual Report Mathinna Gold Project 12 mths ending Sept 1998 – EL3/97. Connemara Gold Mines Pty Ltd. MRT Ref. 98\_4212
- Findlay, R.H. 1993. Report on visit to Victoria, Long Struggle and Linton Mines, and to White's Alluvial Workings, Mangana 30 Jun to 1 Jul 1993. MRT Report 1993/19
- Gresham, J.J. 2004. Exploration Potential in North East Tasmania – EL 17/1991. Cala Resources Pty Ltd. MRT Ref. 04\_5034
- Iloff, G.D. 1994. Investigation of Gold Mine Tailings, NE Tasmania Gold Corridor 1994. MRT Ref. 94\_3632
- Iloff, G.D. 1996. Investigation of Gold Mine Tailings Summary Report 1994 – EL22/92. Mancala Pty Ltd. MRT Ref. 96\_3932
- Jackson, D.G. 2000. Annual Report for the period 30 May 1999 to 29 May 2000 – EL18/91 Mangana. Defiance Mining NL. MRT Ref. 00\_44438
- Jackson, D.G. 1999. Annual Report Mangana – EL18/91. Defiance Mining NL, Connemara Gold Mines Pty Ltd. MRT Ref. 99\_4321

Leaman, D.E. 1989. Ground Magnetic Survey – EL55/83 Mangana. Alcaston Mining NL. MRT Ref. 89\_3052A

Leaman, D.E. 1990. Aeromagnetic Survey Mangana Acquisition Report (including preliminary interpretation) – EL55/83 Mangana. Alcaston Mining NL. MRT Ref. 90\_3197A

McDonald, G. 1996. Annual Report 1995 – EL18/91 Mangana. Resolute Samantha Ltd. MRT Ref. 96\_3842

Morrison, K.C. 1987. Assessment of Alluvial Gold Prospects: Mathinna, Mangana and Lisle. Alcaston Mining NL. MRT Ref. 87\_2750A

Morrison, K.C. 1988. Partial Relinquishment Report – EL 55/83. Alcaston Mining NL. MRT Ref. 88\_2873

Morrison, K.C. 1988. Annual Report – Year 5 24<sup>th</sup> Nov 1987 to 23<sup>rd</sup> Nov 1988 – EL55/83 Mangana. Alcaston Mining NL. MRT Ref. 88\_2883

Morrison, K.C. 1989. Annual Report – Year 6 24<sup>th</sup> Nov 1988 to 23<sup>rd</sup> Nov 1989 – EL55/83 Mangana. Alcaston Mining NL. MRT Ref. 89\_3052

Morrison, K.C. 1990. Annual Report – Year 7 24<sup>th</sup> Nov 1989 to 23<sup>rd</sup> Nov 1990 – EL55/83 Mangana. Alcaston Mining NL. MRT Ref. 90\_3197

Murdoch, R. 1986. A Report on Metallurgical Testwork to Nov 1985 Mathinna Gold Tailings Dumps Tasmania. MRT Ref. 86\_2546

Newnham, L.A. 1993. Review of previous exploration information – EL22/92 and 23/92 North-East Tasmania. Resolute Samantha Ltd. MRT Ref. 93\_3498A

Pearson, D.F. 1993. Annual Report to Sept 1993. Newcrest Pty Ltd. MRT Ref. 93\_3498

Steward, W. 2001. Annual Report for the period 30 May 2000 to 29 May 2001 – EL18/1991 Mangana Defiance Mining NL. MRT Ref. 01\_4573

Summons, T.G. 1985. Hard Rock Gold Potential of EL55/83 North East Tasmania. MRT Ref 85\_2510A

Twelvetrees, W.H. 1907. The Mangana Goldfield Geological Survey Bulletin No. 1. MRT Ref. GSB\_100

White, K.A. 1985. Annual Report Year 2 24<sup>th</sup> Nov 1984 to 23 Nov 1985 – EL55/83. MRT Ref. 85\_2510

White, K.A. 1994. Yearly Report – EL18/91. Mangana for 1993/94. MRT Ref. 94\_3564

Wolff, G.C. 1978. EL 17/78 Mathinna Goldfield. Tasminex NL. MRT Ref. 78\_1318

Wolff, G.C. 1979. Exploration within Mathinna, Tasmania, 30<sup>th</sup> Oct 1978 to 30<sup>th</sup> Apr 1979 – EL17/78. Tasminex NL. MRT Ref. 79\_1344

6 EXPENDITURE

Total annual expenditure for EL35/2008.

<b>Geoscientific Costs</b>	
Geology	8,465.61
Geochemistry	
Geophysics	
Remote sensing	
<b>Drilling &amp; Gridding Costs</b>	
Gridding	
Drilling	
Earthmoving	
<b>Land Access Costs</b>	
<b>Feasibility Costs</b>	
Other Costs	354.25
Rental fees	
Vehicular track Construction	
Surveying, contract drafting etc	
Capital equipment purchase	
<b>Administration Costs</b>	
(note: not to exceed 10% of annual expend)	
Legal	
Office & Admin	
<b>Total</b>	<b>8,819.86</b>