

SHREE MINERALS LIMITED

ACN 130 618 683



TENEMENT SURRENDER REPORT

MOUNT BERTHA - EL42/2004

February 2014

Author : M. Pal
Shree Minerals Limited
ACN 130 618 683
Unit 4, Pine Business Centre, 86 Forrest Street
Cottesloe WA 6011

TABLE OF CONTENTS

SUMMARY

1. INTRODUCTION	2
2. AIM	2
3. LOCATION AND ACCESS	2
4. TENEMENT STATUS	3
Schedule	4
Land Tenure	4
Exclusion	4
5. PHYSIOGRAPHY & VEGETATION	6
6. GEOLOGICAL SETTING	7
Regional Geology	7
Local Geology	9
Mineralisation	10
7. PREVIOUS EXPLORATION	11
Pickands Mather 1960 to 1966	12
ESSO Minerals Limited - 1974	12
Mineral Holdings Australia – 1978	12
Comstaff Pty – 1981	12
BHP – 1982	12
CRAE Pty Limited – 1983	12
Petrecon Australia – 1988	12
Geopeko - 1992	12
Allstate – 1996	12
Titan Goldstream – 1998	13

Pacific Nevada – 1999	13
Zinico NL. 2005 – 2008	13
Shree Minerals – 2008 – 2009	13
Shree Minerals – 2009 – 2010	14
Shree Minerals – 2010 – 2011	15
Shree Minerals - 2011-2012	16
Shree Minerals - 2012-2013	16
8. WORK PERFORMED	16
9. RECOMMENDATIONS	16
10. REFERENCES	17

Item No.	LIST OF FIGURES	Page No.
Figure 1	Tenement (EL42/2004) location and access map	3
Figure 2	Mt Bertha (EL42/2004) Land Tenure map	5
Figure 3	Mt Bertha (EL42/2004) and environs map showing excluded areas	6
Figure 4	Location of “Stratotectonic Elements” and tenements	9
Figure 5	Tenement (EL42/2004) geology map	11
Figure 6	Mt Bertha location plan showing parts retained /surrendered	14
Figure 7	Geophysically identified exploration targets – EL42/2004	15
	LIST OF APPENDICES	
Appendix I	List of appended digital data files	18

SUMMARY

The Mt Bertha tenement (EL42/2004) of Shree Minerals Limited (the Company, or Shree) is located 20 km northeast of the Savage River Iron Ore Mine and about 50 km southwest of Burnie, in North West Tasmania. The tenement covers an area of 134 km².

The licence geology comprises a variable volcano-sedimentary package of Neoproterozoic-aged rocks including part of the Arthur Metamorphic Complex (AMC). The Arthur Metamorphic Complex (ACM) occupies a tectonic feature also known as the Arthur Lineament and lies between the Rocky Cape and Dundas stratotectonic elements. The AMC is a strongly deformed blueschist and greenschist grade metamorphic belt of 110 km northeast-southwest strike and 10 km width in the remote NW of Tasmania. The rock sequences within the complex are considered prospective for industrial mineral deposits such as silica sand at Corinna, magnesite at Keith River and base metals (copper-lead-zinc-silver), as well as gold, etc.

The region due to rugged terrain, thick vegetation, lack of good access, etc., has not been explored systematically in detail even by large companies like ESSO, CRAE, BHP, Geopeko, etc.,. Prior to Shree's exploration, work done was mainly geophysical study of available airborne magnetic-radiometric data with some water and sediment sampling. The studies have identified a number of potential exploration targets. Unfortunately none of the targets have been examined in detail. Since 2005 a number of desktop studies (Tear 2005/6, Hungerford 2005 and Cowan 2010) were carried out.

During the report period, a reconnaissance exploration field work over two weeks period in March 2012 was undertaken.

Exploration mainly targeted significant magnetic highs, some being identified within or adjacent to windows through the Tertiary and Permian cover sequences located in the vicinity of Savage River Slurry Pipe Line Road identified in the 2010 geophysical study for the tenement and environs. The areas visited during reconnaissance, in addition to magnetite were explored for base metal and other commodities such as magnesite.

Magnetic susceptibility meter was utilised to test as many rocks as possible in the field. This generated 166 data points to enable more effective modelling of the area; in particular allowing characterisation of the variably magnetic Tertiary Basalts, which are widespread and mask potential subsurface magnetite mineralisation within the Arthur Metamorphic Complex (AMC) schists.

A total of 17 rock chip samples were collected, mostly in the form of composites, some with additional reference samples. A road side schist (AMC) outcrop was composite sampled returned an indicative 35m @ 242ppmCu(max401ppmCu).

Details on work conducted are given in Appendices III-V.

1. INTRODUCTION

The Mt Bertha tenement (EL42/2004) geology comprises of a variable volcano-sedimentary package of Neoproterozoic-aged rocks including part of the Arthur Metamorphic Complex (AMC). The AMC is considered potential for copper-lead-zinc-silver, gold and may be some near surface goethitic-hematite mineralisation, magnetite, and magnesite resources.

This report summaries the work performed from 1 March 2013 to 28 February 2014.

2. AIM

To explore for iron, magnetite, magnesite, base metals and gold resources.

3. LOCATION AND ACCESS

The centre of the Mt Bertha tenement is located 20km northeast of the Savage River Iron Ore Mine and about 50km south-west of Burnie, in North West Tasmania.

Access to the tenement is restricted, with the only access being via the Savage River Slurry Pipe Line road, which runs through more or less 2/3rd the length of the licence area (Figure 1).

Off-road access is difficult; requiring track cutting and possibly helicopter-supported access. Previous explorers have created some 4WD tracks, which may require restoration for access to areas of interest.

Access to the north of the tenement is by unsealed roads constructed for previous magnesite exploration.



Figure 1: Tenement (EL42/2004) location and access map

4. TENEMENT STATUS

The tenement EL42/2004 (Figures 1) was granted to Zinico NL on 1 March 2005 for 5 years with expiry on 28 February 2010 for exploring Category 1 and 5(a) Minerals. On 22 November 2005 Zinico NL changed its name to Zelos Resources NL (Zelos), and on 23 November 2006, to reflect the major shareholding, the Zelos name was changed to Gujarat NRE Resources NL. In May 2008 the tenement was acquired by the Indo Australian Consulting Group Pty Ltd (IACG). The tenement covers an area of 134 km².

Pursuant to a Farmin Agreement with IACG, the terms of which were varied under a Deed of Variation entered into on 10 November 2009, the Company was granted the right to acquire 75% interest in EL42/2004. Now the ownership of Mt Bertha tenement is 75% (Shree) and 25% (IACG). However, the operational part is 100% controlled by Shree. Moreover, under the terms of the Deed of Variation, Shree is responsible to sole fund all expenditure in relation to the tenement, including pre-feasibility study.

Schedule

Land district: Russell vicinity of Mt Bertha
Municipality: Waratah/Wynyard and Circular Head
Exploration Licence: 42/2004
Area: 134 km²
Ownership: Shree Minerals 75% and IACG 25%
Operator: Shree Minerals Ltd.

The coordinate datum for the licence is based on GDA 1994, MGA Zone 55. Tenement boundary is shown in Figures 1 to 3.

Land Tenure


The area comprises (Figure 2):

- State/Multiple use Forest
- Donalson River Nature Recreation Area
- Savage River Regional Reserve
- Arthur River Forest Reserve; and
- MDC Informal Reserve


Exclusion

The exclusion areas are list below and shown in Figures 3 and 6:

- Any land owned or leased by the Commonwealth of Australia.
- Mining Leases amounting to 357 ha (more or less) which were applied for or in force prior to the date of application for this licence.
- Crown reservations or other land set apart or dedicated for any public purposes such as public reserves, municipal reserves or roadways unless such areas have been brought under the provisions of the *Mineral Resources Development Act 1995*.
- Areas of private land which either have been, or are in the process of being, purchased by the Crown under the Regional Forest Agreement - Private Forests Reserves Program and / or private land over which the landowners have agreed, or are in the process of agreeing, to place a covenant or management agreement for conservation purposes under the Regional Forest Agreement - Private Forests Reserves Program.



Source: MRT
Figure 2: Mt Bertha (EL42/2004) Land Tenure map



Source: MRT

Figure 3: Mt Bertha (EL42/2004) and environs map showing excluded areas

5. PHYSIOGRAPHY & VEGETATION

The licence area has considerable relief, being higher on average to the south east. This raised area occurs throughout the SE half of the licence and presents a considerable challenge to exploration.

Vegetation comprises dense forest (temperate rainforest) making access very difficult, necessitating substantial track cutting to reach areas of interest. Climate is temperate with substantial annual rainfall, typical of Western Tasmania. Temperature ranges from just above freezing in winter to a likely maximum of 30°C in summer.

6. GEOLOGICAL SETTING

Regional Geology

Tasmania has been geologically divided by MRT into seven Proterozoic-Lower Palaeozoic regions or "Stratotectonic Elements", each with a different geological history and economic mineral associations (Figure 4). Because of multiple subduction episodes these elements or terranes were welded together during geological history, which has produced the current geological framework for Tasmania. An abbreviated stratotectonic history of Tasmania is detailed below:

- Formation of basement as Early Neoproterozoic-aged shelf clastic sedimentation with an age range of 900-1000 million years ago (ma) followed by a major orogenic event at 760ma, which included granite intrusions". This produced the Rocky Cape Element.
- A failed rift episode then followed with its associated clastic sedimentation and volcanic inputs ensued by a second, successful rift event that happened in the Late Neoproterozoic to Early Cambrian. This added an assortment of units including mafic lavas to the Rocky Cape Element i.e. the Togari Group (and its equivalent Ahrberg Group).
- An island arc-continent collision east or northeast of Tasmania occurred in the late Early Cambrian and the emplacement of a series of allochthonous slices across Tasmania, including oceanic assemblages (ultramafics and associated mafic lavas) and other units. This formed the Dundas, Sheffield, Tyennan, and Adamsfield-Jubilee Elements.
- A series of Mid to Late Cambrian clastic basins developed post-collision and were concomitant with major calc-alkaline volcanism - the Mt Read Volcanics that contain an excellent volcanogenic-hosted massive sulphide (VHMS) province.
- This was followed by Late Cambrian orogenesis comprising fold belt-style tectonics at 500-510ma and includes some thrust stacking of units.
- The establishment of a statewide clastic basin began in Late Cambrian times with initial basal conglomerates overlain by limestone lithologies followed by a gradually deepening marine clastic sequence up to Mid Devonian times. At the same time, the Northeast Tasmanian Element developed as a turbiditic basin quite distinct from the other elements and lies east of an inferred subduction suture zone.
- Cessation of sedimentation was caused by uplift and erosion associated with the Tabberabberan Orogeny (Mid-Devonian) and with a subsequent Late Devonian to Early Carboniferous phase of major granitic intrusions. This included the Heemskirk, Meredith, and the Northeast Tasmanian Granites, with the first two causing modifications to the Cambrian morphology via structural overprints and hydrothermal alteration effects. These granite intrusions resulted in the formation of many skarn and vein deposits for tin, nickel, lead/zinc etc. The tectonism also resulted in the

structurally controlled Henty gold deposit. In Northeast Tasmania, the Devonian-aged intrusions and deformation are associated with gold mineralisation.

- Minor sedimentation including glacial deposits occurred in the post-Devonian Tasmania Basin. Substantial amounts of dolerite and basalt were formed because of continental break up associated with Jurassic and Tertiary global events. Continental extension and rifting began in Mid Jurassic times with separation occurring in the Mid Cretaceous. Major Jurassic dolerites related to a Gondwana event occur as sills across Tasmania and are similar to the Karoo series in Africa.

The Arthur Metamorphic Complex (AMC), or the Arthur Lineament as it is also known, is an elongate (100 km by 10 km), NE/SW striking, high metamorphic grade geological belt/tectonic boundary that occurs in NW Tasmania. It lies between the Rocky Cape and Dundas/Sheffield stratotectonic elements and has a metamorphic grade of upper greenschist to amphibolite with localised blueschist facies. The eastern boundaries of the complex are transitional into less deformed and less metamorphosed rocks whereas the western boundary is thought to be an east-dipping thrust contact. The rock sequences within the complex are rich in industrial mineral deposits e.g. iron ore at Savage River, silica sand at Corinna, magnesite at Keith River etc.

Current theory for the AMC suggests that it is an allochthonous unit of strongly metamorphosed Rocky Cape Group sediments i.e. the Neoproterozoic Togari Group. These sediments include mafic volcanics that have been strongly deformed across a major fault suture zone i.e. a klippe. Offshore seismic data across the AMC indicates a relatively shallow level of penetration for the unit implying that the rock sequences became detached from source, were substantially folded by thrust tectonics, and placed into their current position by an east-dipping thrust system. The west margin of the AMC and further west are believed to have been emplaced by west dipping thrusts.




Figure 4: Location of “Stratotectonic Elements” and tenements

Local Geology

The geology of the Mt Bertha licence comprises a variable volcano-sedimentary package of Neoproterozoic-aged rocks including part of the Arthur Metamorphic Complex, as described above (Figure 5).

In detail, the Proterozoic group strikes generally northeast southwest and steeply dipping and young from west to east across the licence. The oldest units are siltstones and pyritic mudstones of the Early Neoproterozoic Cowrie Siltstone. These are overlain by a mixed siliciclastic package of siltstones, quartzites and sandstones with minor pelitic shales (Detention Quartzite, Jacobs Quartzite, Irby Siltstone etc). Subsequent units in the southern part of the property comprise carbonates, clastics, volcanic turbidites, and tholeiitic basalts of the Neoproterozoic Forest Conglomerate, Togari and Ahrberg Groups. These in turn are overlain by chert, shale, conglomerate, and dolomite of the Black River Dolomite and associates. To the north of the property Neoproterozoic phyllites occupy the Togari, Ahrberg, and Black River Dolomite positions. The remaining Neoproterozoic sequence consists of a chloritic schist unit, the Bowry Formation, with dolomites and magnesite deposits, The youngest unit in the complex is the Keith Schist which comprises quartz mica schists, quartzite and phyllite and is thought to be a more deformed version of the

east bounding Burnie and Oonah turbiditic siltstone packages (both Late Neoproterozoic in age).


Permian sequences that run along the centre of the lease appear to be fault bound in a graben-like structure against the various Neoproterozoic sequences, comprising a lower glaciomarine clastic sequence with limestones and the Tasminite Oil Shale. Overlying these rocks are coal measures followed by an upper glaciomarine sequence. At the northeast corner of the tenement it appears that, the Permian unconformably lies on Neoproterozoic schists.

A subsequent Tertiary basalt eruptive phase resulted in extensive coverage of the tenement (about 60%) masking the underlying Proterozoic units. A review of recently flown airborne magnetic data indicates that the basalt cover may be quite thin in several instances, as demonstrated by the continuity of the Neoproterozoic-related magnetic signatures underneath the basalt cover.

The airborne magnetic data indicates a substantial structural complexity with several major structures transecting the licence. There are likely to be some differences between new geologic deductions from this air magnetic data and the published geology, which may create exploration opportunities.

Mineralisation

Reported mineral occurrences at the property are restricted to magnesite on the periphery of the main magnesite leases that occur just beyond the northeast corner of the tenement and two small gold occurrences at the extreme north-east end of the licence. There are minor copper occurrences outside the tenement in the northeast area. The nearby Savage River Iron Ore Mine (15km south west of the southern boundary) consists of concordant massive pyrite-magnetite hosted by greenschist grade tholeiitic metabasalts of the Bowry Formation. In general, the gold occurs as numerous small scale hard rock and alluvial deposits which were mainly worked in the 19th Century. A few small base metal deposits, mostly of copper are also known around the general area (MRT data source). With a strong structural overprint, fundamental geological faults and favourable mineral hosting units, there is strong potential for a variety of styles of economic gold/ copper mineralisation within the licence.



Source: MRT

Figure 5: Tenement (EL42/2004) geology map

7. PREVIOUS EXPLORATION

North-Western Tasmania is subject to severe weather, lacks infrastructure and has dense vegetation cover.. These factors along with extensive Tertiary Basalt cover, difficult terrain and lack of good outcrops have inhibited exploration in the area.

Airborne surveys have led the way in the past with a series of anomalies being identified, some of which have been followed up. No drilling has been undertaken in the licence area but in the past, pre 1985, one or two holes were drilled peripheral to the licence area, e.g. Comstaff in the 1980's.

Below is a very brief out line of previous explorer activities:

Pickands Mather 1960 to 1966

Modern exploration can be considered to have commenced in the mid-1960's with regional stream geochemical sampling by Pickands Mather International throughout western Tasmania. In 1966 they developed and commenced mining at the Savage River Iron (magnetite) Ore Deposit.

ESSO Minerals Limited - 1974

Esso's exploration target was base metal (massive sulphides). For that it conducted an airborne magnetic and EM INPUT survey over a wider area, including EL 2/73 and identified 63 EM anomalies from a survey much larger than EL42/2004 of which 36 were not explained due to access problems. Some of these anomalies are within the current licence. Most of these anomalies appear to be formational features associated with the Cowrie Siltstone and associates from the Rocky Cape Group.

Mineral Holdings Australia – 1978

Mineral Holdings Australia carried out exploration to the north east of the tenement and outlined the magnesite deposits in the Lyons and Arthur Rivers areas (EL43/70).

Comstaff Pty – 1981

Comstaff Pty Ltd partially covered the licence with DIGHEM and did some drilling south of the current licence. The search included tin, platinum group and chromite as well as base metals (EL1/68).

BHP – 1982

BHP Minerals in 1982 carried out photogeology, re evaluated of past surveys, stream sediments, and heavy minerals search (EL18/80).

CRAE Pty Limited – 1983

CRAE in 1983 carry out airborne magnetic and radiometric survey and identified a series of magnetic anomalies within the general area, some of which were followed up with drilling with negative results i. e. the Rapid and Rapend series of prospects located near the Rapid River (EL1/79).

Petrecon Australia – 1988

Petrecon Australia completed a new geology map of the area including EL24/87

Geopeko - 1992

Geopeko carried out water and stream sediment sampling over lands of ELs 40/89, 41/89, and 42/89 for stratabound tungsten of the Mittershill-type. Results were disappointing.

Allstate – 1996

Allstate in 1996 carried out exploration work for low grade, fine-grained gold in close association with ironstone-carbonate-mafic volcanics over tenements EL35/94 and EL36/94. The work also included Besshi-type targets. The company completed a geophysical study of airborne data procured by AGSO and identified a variety of anomalies based on inferred structural intersections coinciding with magnetic features. There appears to

be AMG coordinate discrepancies between maps presented by the geophysicist in his report and CRAE's airborne magnetic maps. The CRAE targets are common to both maps and indicate a coordinate shift of some 6 km on a 190° bearing.

Titan Goldstream – 1998

Titan-Goldstream Joint Venture, using Homestake-style Proterozoic ironstone associated gold exploration model, explored EL37/96 and EL38/96 by stream sediment sampling in 1999. The results were disappointing.

Pacific Nevada – 1999

Pacific Nevada explored EL24/97 by reprocessing of available AGSO airborne geophysical data and some stream sediment sampling.

Zinico NL. 2005 – 2008

From 2005 to 2008 no field work was carried out. The only work performed was review of “open file” information from the MRT data base, literature search and geological and geophysical interpretations of information collected. Geological interpretation was undertaken by SMG Consultants and geophysical interpretation using airborne magnetic-radiometric data collected as a part of the [Western Tasmania Regional Minerals Program](#) (WTRMP) by Hungerford Consulting.

From these studies, a geological map of the area was generated. The map was generated from the identification of geological units with distinctive magnetic signatures.

Geophysical study identified 6 targets (4 priority 1 and 2 for consideration).

Table 1: Hungerford identified geophysical targets

Target No	Location (m)		Target Category
	Easting	Northing	
			Primary
A	360 400	5 428 500	Primary
B	357 600	5 425 300	Primary
C	354 900	5 425 500	Primary
D	364 600	5 433 900	Primary
E	357 500	5 427 400	Secondary
F	360 000	5 430 400	Secondary

The work was reported in Harder 2006

Shree Minerals – 2008 – 2009

During the report period the Company made several visits using Aircomander light aircraft and helicopter to the area. These flights were aimed to assess the topography and possible alternate access methods such as helicopter landing sites. Later on using helicopter three out of four geophysically suggested targets were visited.


Shree Minerals – 2009 – 2010

No field work was undertaken, however preparations for exploring “Hungerford suggested geophysical Target C (354, 930 m E/ 5, 425, 430 m N)” were made. **Target C** is the highest intensity anomaly within the tenement and occurs on a very long magnetic trend obscured by Tertiary basalts.

The modeled profile indicates that the magnetic source is from the Bowry Formation sub-cropping at the base of the overlying basalts about 70 m below the natural surface. Bowry Formation rocks outcrop about 1.5 km to the south along strike so that mapping and geochemical sampling here may indicate the presence of alteration or mineralisation.

The tenement Mt Bertha (EL42/2004) is in the interior of the north- west quarter of Tasmania. The tenement area is large and based on information from past exploration is considered prospective for base metals (copper-lead-zinc-silver), magnesite, as well as gold.

The Company based on available information considered the interior of the licence (Figure 6) prospective and accordingly 90 km² of tenement area as shown in Figure 6 was surrendered.



Source: MRT


Figure 6: Mt Bertha location plan showing parts retained /surrendered

Shree Minerals – 2010 – 2011

During the report period, a preliminary geological examination of one target (Target C) identified by earlier studies and a detailed study of the 2001 Tasmanian Government acquired airborne magnetic-radiometric data were undertaken. The geological examination of Target C due to thick vegetation and lack outcrops did not deliver much information.

The geophysical study has identified 12 exploration targets (6 Tabular type and 6 Plug type Figure 7). In general the study concluded that the structural setting along part of the Arthur River Metamorphic Complex (AMC) is favourable for base metal and other commodities such as magnesite.

Unfortunately, there is no evidence of high amplitude magnetic anomalies similar to Savage River, however there could be potential for goethitic -hematite type discovered at the Company's Nelson Bay River Iron Project.



Source: Cowan Geodata

Figure 7: Geophysically identified exploration targets – EL42/2004

Shree Minerals - 2011-2012

Due to heavy work commitment at its advance Nelson Bay Iron Ore Project the Company could not carry out field work during the reporting period at the tenement. The Company applied to MRT for approval of an exploration program to commence in mid to late January 2012. However, due to rugged terrain, poor access, thick vegetation cover, and non availability of suitable technical personnel the program could only be undertaken in March 2012.

Exploration mainly targeted significant magnetic highs, some being identified within or adjacent to windows through the Tertiary and Permian cover sequences in the vicinity of Savage River Pipeline.

Shree Minerals - 2012-2013

During March 2012 two weeks reconnaissance field work at the tenement was carried out and results reported in in the 2012-13 Annual report..

8. WORK PERFORMED

During the report period all the Company resources were devoted in attending court proceedings relating to injunction brought by the Tarkine Coalition and seeking reapproval from the Commonwealth Government for commencing mining at the Nelson Bay Iron Ore Project, as a result no exploration work was carried out in the Exploration Licence (EL42/2004) area.

9. RECOMMENDATIONS

To date work carried out at the tenement although is geologically encouraging, but in view of rugged terrain, poor access, thick vegetation cover, limited reliable information on geology, especially stratigraphy, it is recommended that the resources be directed towards the development of the Company's Flagship Nelson Bay River Iron Project and tenement is surrendered.

10. REFERENCES

1. Cowan, D., 2010. Mt Berth Aeromagnetic Survey Reprocessing and Preliminary Interpretation
2. Funnell, F. R. , 1987. Rapid River EL 1/79 NW Tasmania, CRAEL47/53 Final Report 1987 (MRT Report No 87-2723)
3. Harder, W. M., 2006., EL42/2004 Mt Bertha Year 1 Annual Report, For the period 1 July 2005 – 1 March 2006. (Zelos Resources NL)
4. Harder, W. M., 2007. EL 42/2004 Mt Bertha Year 2 Annual Report, 1 March 2006 – 1 March 2007. (Gujrat NRE Resources NL)
5. Harder, W. M., 2008. EL 42/2004 Mt Bertha Year 3 Annual Report, For the period 1 March 2007 – 1 March 2008 (Gujrat NRE Resources NL)
6. Harder, W. M., 2008. EL 42/2004 Mt Bertha Year 4 Annual Report, 1 March 2008 – 1 March 2009
7. Harder, W. M., 2010. EL 42/2004 Mt Bertha Year 5 Annual Report, 1 March 2009 – 1 March 2010
8. Hungerford., N., 2005., Geophysical Interpretation Report., December 20
9. Matison, I., EL 41/89 Mt Bertha, Geopecko Annual Report 1991 (MRT Report No 92-3329)
10. Matison, I., EL 42/89 Rapid River Geopecko Annual Report 1991 (MRT Report No 92-33320)
11. Pal, M., 2011. Annual Report for the period 1.03.2010 to 28.02.2011 Mt Bertha - EL42/2004
12. Pal, M., 2011. Annual Report for the period 1.03.2011 to 28.02.2012 Mt Bertha - EL42/2004
13. Pal, M., 2012. Annual Report for the period 1.03.2012 to 28.02.2013 Mt Bertha - EL42/2004
14. Ridge, K. J., 1996. Savage River Mt Bertha , N. W. Tasmania. EL35-36/94., Annual Report 1995 Allstate Exploration NL. (MRT Report No 96-3876)
15. Reid, R., 2010., Report on Field Work at Mt Bertha (EL42/2004) 2010 Drilling – Davie Prospect, Sulphide Creek, EL42/2004
16. Tear., S., 2005., Zinico Resources NL: Prospectus., August 2005
17. Tear., S., 2006., Literature Study and Geological Interpretation, August 2005
18. Turner., N. J., 1998. EL 37/96 Rapid EL38/96 Savage River EL46/96 Flowerdale Combined Annual Report Goldstream Mining NL

APPENDIX I

List of appended digital data files

1. EL422004_201402_01_Digital_Files.txt
2. EL422004_201402_02_Annual_Report.pdf