



STELLAR RESOURCES LIMITED
Rubicon MinTech Ventures Pty. Ltd.

EL 1/2004 RAMSAY RIVER

FINAL & ANNUAL REPORT FOR THE PERIOD
10 FEBRUARY 2017 – 9 FEBRUARY 2018

by: **A. M. Rigg**

DATE: 9 February 2018

DISTRIBUTION:
Mineral Resources Tasmania, a Division of the
Department of Infrastructure, Energy and Resources - Hobart
Stellar Resources Limited - Melbourne

ABSTRACT

This Annual Report for EL1/2004 Ramsay River covers the period from 10 February 2017 to 9 February 2018. This is the final report for the licence.

From 2011 to 2015 Stellar conducted detailed geological mapping and geochemical sampling of the Butler's Road tin anomaly area, with broadly encouraging results, but has not been able to clearly define effective drilling targets. Partial relinquishments reduced the licence area in 2009 and 2012. The remaining licence area is being relinquished, as a high grade target cannot be defined.

Expenditure on EL1/2004 for 2017 totalled \$8,106.

Expenditure for the licence from 2005 to 2017 totalled \$941,788.

TABLE OF CONTENTS

ABSTRACT

1. INTRODUCTION	4
1.1. EXPLORATION RATIONALE & GEOLOGICAL SETTING	4
1.1.1. <i>Geological Setting</i>	4
1.2. LICENCE	4
LAND TENURE	7
2. REVIEW OF PREVIOUS & STELLAR WORK	9
3. EXPLORATION COMPLETED DURING THE REPORTING PERIOD	10
4. CONCLUSIONS	14
5. ENVIRONMENT	14
6. EXPENDITURE	15
7. REFERENCES	16

LIST OF FIGURES

Figure 1. EL1/2004, Licence Location and Relinquishment Area	5
Figure 2. EL1/2004, MRT Geology Plan	6
Figure 3. EL1/2004, Public Land Classification, LIST	8
Figure 4. EL1/2004, Pre-Stellar Exploration, Drilling, Geology, Geochemistry, Ground Geophysics. ...	11
Figure 5. EL1/2004, Pre-Stellar Exploration, Airborne Geophysics	12
Figure 6. EL1/2004, Stellar Exploration, Drilling, Geology, Geochemistry, 2005 - 2017.	13

1. INTRODUCTION

1.1. EXPLORATION RATIONALE & GEOLOGICAL SETTING

As a consequence of exploration results from previous Comstaff work and Stellar Resources work on EL1/2004, exploration aims for the 2012 - 2017 period were focussed on the low sulphide tin potential of fractionated facies and apophyses of the Meredith Granite and its contact zones, within the remaining portion of the original EL (Figure 6).

There are several small tin and base metal occurrences within the licence area, with the Mt Bischoff (Sn) and Cleveland (Sn-Cu) mines lying within 3km to the east and west.

1.1.1. Geological Setting

The licence covers part of the Late Carboniferous tin-prospective NE portion of the Meredith Granite, which has intruded an association of Early Cambrian Luina Group marine sediments and basaltic rocks, and Early Cambrian allocthanous slivers of mafic-ultramafic rocks (Figure 2). The northern end of the prospective area is partly covered with Tertiary basalt.

The north-east corner of the Meredith Granite is known to extend at shallow depth and underlie the historic Mt Bischoff porphyry and dolomite replacement tin deposit. The historic Magnet Mine is located on a northern boundary of the Ramsay licence. It is a lode-style base metal and silver deposit (0.64Mt @ 7.3%Zn, 7.3%Pb and 427 g/t Ag) hosted by a structurally emplaced mafic/ultramafic body known as the Magnet Dyke. The lower levels of the old mine (below 8 level) are within EL1/2004 while the postulated feeder structure trends south-west into the EL. Tin may occur below the old workings in a zone closer to the underlying granite. Anomalous tin soil geochemistry is evident in the Butler's Road area, where previous explorers generated currently unexplained EM anomalies, highlighting the area for possible significant greisen style mineralisation within a zone of quartz-tourmaline alteration (schorl rock) of a Devonian feldspar-quartz porphyritic biotite granitoid between the cupola and the Early Cambrian Luina Group host sedimentary rocks.

1.2. LICENCE

TENEMENT NUMBER: 1/2004

TENEMENT NAME: Ramsay River

TENEMENT LOCATION: Located 60km south-west of Burnie, with main road access from the Corinna-Waratah Road approximately 10km west of the Murchison Highway (Figure 1). The licence covers 41km² from the Magnet Mine area west of Waratah township, south to 5405000mN (GDA94) which is 1km south of the Corinna-Waratah Road in the vicinity of the Mt Ramsay Track. Most of the EL area is Crown Land, covered by patches of rainforest, forest, tea-tree scrub and button grass plain. Access is provided by the Corinna Road, numerous logging and old exploration tracks, and walking tracks. Much of the area is accessible only by foot.

TENEMENT YEAR: 10 February 2017 to 9 February 2018.

TENEMENT HOLDER: Rubicon MinTech Ventures Pty Ltd., a wholly owned subsidiary of Stellar Resources Ltd.

REPORTING PERIOD: 10 February 2017 to 9 February 2018.




Figure 1. EL1/2004, Licence Location and Relinquishment Area.




Figure 2. EL1/2004, MRT Geology Plan

LAND TENURE

SCHEDULE

LAND DISTRICT OF RUSSELL
VICINITY OF RAMSAY RIVER 8KM SW OF WARATAH
MUNICIPALITY OF WARATAH / WYNYARD
EXPLORATION LICENCE 1/2004 41km²
RUBICON MIN TECH VENTURES PTY. LTD.

Commencing at the northwest corner at grid coordinates 367 112 mE 5 412 184 mN, thence grid east to 370 112 mE, grid south to 5 410 184 mN, grid east to 375 112 mE, grid south to 5 406 000 mN, grid west to 371 000 mE, grid south to 5 405 000 mN, grid west to 369 000 mE, grid north to 5 406 000 mN, grid west to 367 112 mE and finally grid north to the point of commencement.

Coordinate datum – GDA94, MGA Zone 55.

EXCLUSIONS

- (a) Any land owned or leased by the Commonwealth of Australia.
- (b) Mining leases amounting to 70ha (more or less) which were applied for or in force prior to the date of application for this licence.
- (c) Crown reservations or other land set apart or dedicated for any public purposes such as public reserves, municipal reserves or roadways unless such areas have been brought under the provisions of the *Mineral Resources Development Act 1995*.
- (d) Land declared as a fossicking area under the *Mineral Resources Development Act 1995* as shown hereunder:
10ha Magnet Fossicking Area
- (e) Areas of private land which either have been, or are in the process of being, purchased by the Crown under the Regional Forest Agreement - Private Forests Reserves Program and / or private land over which the landowners have agreed, or are in the process of agreeing, to place a covenant or management agreement for conservation purposes under the Regional Forest Agreement - Private Forests Reserves Program.

LAND TENURE

The area (Figure 3) comprises:

Private Property
Multiple Use State Forest
Meredith Range Regional Reserve
Savage River Regional Reserve

The licence area contains areas, which are listed (including listed on an interim basis) on the Register of the National Estate kept under the *Australian Heritage Commission Act 1975*.




Figure 3. EL1/2004, Public Land Classification, LIST.

2. REVIEW OF PREVIOUS & STELLAR WORK

Data from MRT digital geology, geophysics and geochemical datasets and open-file company reports has been captured, summarised and reviewed. Over the term of the licence several targets have been reviewed and field tested with soil geochemistry programmes and/or drilling.

Comstaff Ltd was the most active explorer over the Butler's Rd area. In 1983 Comstaff conducted a Dighem survey, defining five anomalies. In 1984 they took geochemical samples over the Dighem anomalies and cut their Butler's Rd grid, upon which they undertook geological mapping, soil geochemical sampling, a Genie EM survey & ground magnetics. The geological mapping showed a zone of tourmalinisation/greisenisation associated with the northern 'nose' of the Meredith granite. The geochemical sampling indicated anomalous Sn, Pb & Ag with minor Cu & Zn. Pb & As coincided with tourmalinisation/greisenisation. The Genie EM survey produced five anomalies, coincident with the Dighem results, but more accurate. In 1987 BHP in an arrangement with Comstaff, drilled hole BR1 into the Butler's Hill EM/mag anomaly (SRZ RY01). The 32m vertical hole hit mineralisation from 1m below surface to 10m. Sub-vertical veins of calcite-siderite-quartz with sphalerite, minor galena and cassiterite including trace chalcopyrite in hornfelsed sediments were intersected. Granite was not intersected. Assay analyses: 8.5m @ 1.65% Zn, 0.27% Pb, 0.08% Sn, 29g/t Ag, incl 2.8m @ 4.27% Zn, 0.71% Pb, 0.18% Sn, 71g/t Ag, and 0.25m @ 4.42% Zn, 0.91% Pb, 0.17% Sn, 74g/t Ag, and 0.40m @ 2.89% Zn, 0.09% Pb, 0.15% Sn, 21g/t Ag. No further work was done. See Figures 4 & 5.

Pasminco Exploration Ltd held the whole licence area from 1996 to 1998. They flew the "Waratah" aeromagnetic survey (100m fls), and for the Butler's Rd area reviewed previous exploration work, followed by a ground magnetic traverse along Butler's Rd. No further work was done. See Figure 5.

MRT conducted two geophysical surveys that included the licence area, the 2001 WTRMP Area C 200m fls aeromagnetic survey, and the 2002 WTRMP Meredith granite 200m fls HEM survey. Stellar later reviewed the HEM survey data with Dr Jovan Silic and Dr Tom Whiting both (separately) selecting anomalies for testing. EM anomaly RY02 had been detected in previous survey work, but had not been tested.

During 2006 Stellar carried out a mapping and soil/rock chip programme aimed at identifying potential nickel targets in the ultramafic rocks that outcrop around the northern end of Betts Track and near Arthur Dam. The company also carried out a five hole (AD005 – AD009), 1200 m diamond drilling program that was primarily aimed at the further testing of known base metal targets near Arthur Dam. One drill hole tested a magnetic anomaly just west of the entrance to Betts Track. The conclusions were:

- Serpentinised pyroxenite bodies around the northern end of Betts Track and Arthur Dam are relatively small, structurally emplaced lenses with limited potential for nickel mineralisation;
- Hornfelsed, greywacke sandstone that contains substantial magnetite as disseminations and in veinlets is the likely source of strong aeromagnetic anomalies around Betts Track and Arthur Dam;
- Vein style copper mineralisation in the eastern part of the Arthur Dam prospect appears to be of sub-economic grade. However, there is potential for the further drill testing of vein style zinc, lead and silver mineralisation in the south western part of the Arthur Dam prospect.

During February 2009, 36 samples of soil or rock were collected from channels cut in the walls of a series of old costeans and adits south of the Magnet Mine, and assayed for Cu, Pb, Zn, Ag, Sn & Au. These workings were cut across the southern extension of the structure hosting the Magnet mineralisation. Only one sample, from the northernmost adit, returned any significant assays (0.4 %Pb, 1.9 %Zn & 17 g/t Ag).

In 2011 Stellar conducted soil geochemical surveys in five areas of the then licence area to test prioritised EM/mag/geochem anomalies defined through earlier analysis (Silic, 2006) of the 2002 WTRMP Meredith Granite airborne EM survey. Four EM anomalies lie within the current licence area, no's RY01, 2, 3 & 25, with three (RY01, 2 & 3) having been geochemically sampled.

RY01, an EM/mag/geochem anomaly on 'Butler's Hill' near Butler's Road, previously sampled and drill tested by Comstaff/BHP, was soil sampled and mapped. Results confirmed the Comstaff sampling,

showing anomalous Sn, Pb & Zn in a greisenised zone. There was no apparent relationship between the EM anomaly and the adjacent magnetic anomaly. A Sn zone of up to 170ppm in soil was defined over and to the east of the EM anomaly. No further work is considered warranted on RY01.

RY02, an EM/mag/geochem anomaly 900m north-east of RY01 near Butler's Road, also previously soil sampled by Comstaff was sampled and mapped. Again Comstaff sampling was confirmed, showing anomalous Sn, Pb & Zn in a greisenised zone. There was no apparent relationship between the EM anomaly and an adjacent magnetic anomaly. There was no coherently anomalous Sn over the EM anomaly and no geochemical anomalism over the magnetic anomaly. From the western flank of the RY02 EM anomaly, an anomalous Sn zone, larger than that of RY01, extends west for approximately 600m. Following four surface geochemical sampling programmes, the 600m x 300m area (as at 2015) appears to be closed. Rock chip assays range up to 5000ppm, soil assays range up to 3700ppm and pan concentrate assays range up to 11.9%. Phase 1 sampling was conducted in June 2011, Phase 2 in February 2013 and Phase 3 in March 2014, all extending west. See annual reports 2011, 2013 & 2014 for detailed results.

Phase 4, a westerly extension of Phase 3 work, was conducted in January / February 2015.

The Phase 4 soil sampling disclosed some highly anomalous results within the catchment zones of two north/north-east draining creeks where anomalous Phase 4 pan concentrate samples had been taken. The soil sampling showed 17 samples >300ppm Sn with 4 >1000ppm, to a maximum of 2260ppm. The 13 pan concentrate samples for the two drainage lines ranged from 2.6 to 11.9% Sn. The mean of two methods, semi quantitative XRD analysis and thin section point counting, indicates about 6% cassiterite in the pan concentrate and, more importantly, no indication of sulphides or tin minerals other than cassiterite. An anomalous Sn zone approximately 600 metres east-west by 300 metres north-south was generated through the sampling programmes, with the western end being the most compelling. The anomaly coincides with an oval shaped zone of quartz-tourmaline alteration (schorl rock) of a Devonian feldspar-quartz porphyritic biotite granite. The alteration appears to be greisen style and is focussed on the roof and wall rock intrusive contacts between the granitoid cupola and the Early Cambrian Luina Group host sedimentary rocks. It is also the area in which a NNW trending dyke (mapped by MRT) emanates from the granite apophysis and suggests potential for a different, more tin mineralised, alteration style associated with the western contact geology. Geological mapping gave sufficient confidence that the concentration of black tourmaline and quartz distributed around the granite-Luina Group contact represents a greisen-like alteration facies along the edge of the granite, with results consistent with the tourmaline-rich alteration sourcing the weak tin anomalism detected in soil and rock chips. There is however, no convincing evidence of a vector towards high grade zonation within the anomalous zone but rather a substantial zone of shallow, low grade, low sulphide, cassiterite mineralisation controlled by shoots, veins and fracture-fill breccia lodes of quartz-tourmaline greisen. See annual report 2015 for detailed results.

RY03, 1200m south-east of RY01, represented a weaker EM/mag target. The low-order geochemical response was reflective of the geology, rather than elevated mineralisation. No further work is considered warranted on RY03.

See Figure 6.

Note: A complete MapInfo database is supplied separately.

3. EXPLORATION COMPLETED DURING THE REPORTING PERIOD

Administrative work and a final Butler's Road (RY02 west) field inspection was the only work undertaken for the reporting period.




Figure 4. EL1/2004, Pre-Stellar Exploration, Drilling, Geology, Geochemistry, Ground Geophysics.




Figure 5. EL1/2004, Pre-Stellar Exploration, Airborne Geophysics.




Figure 6. EL1/2004, Stellar Exploration, Drilling, Geology, Geochemistry, 2005 - 2017.

4. CONCLUSIONS

For the Butler's Road (RY02) tin prospect, sufficient surface exploration has been done to infer anomaly closure in all directions and although some relatively high grade soil and pan concentrate results were achieved, there is no convincing evidence of a vector towards high grade zonation within the anomalous zone. Microscope and XRD mineralogy conducted on pan concentrates indicates that sand sized cassiterite is the only tin species present and that sulphides are either absent, or at least are rare enough to be undetected. The most plausible conclusion from the data at hand is that a substantial zone of shallow, low grade, low sulphide, cassiterite mineralisation is present, controlled by shoots, veins and fracture-fill breccia lodes of quartz-tourmaline greisen. The mineralisation may be discontinuous in detail but uniform in grade over the bulk scale.

There is no need for further mapping or geochemistry and due to the lack of sulphide and the abundance of tourmaline, it is unlikely that further geophysics would resolve internal detail within the anomaly. The Phase 5 geochemical sampling proposed in the ATR2014, is now regarded as unnecessary.

There is no convincing evidence of a vector towards a high grade zonation within the anomalous zone, and no indication of a trend leading towards a specific and large enough geochemical high worthy of a drill target. The drilling of one 200m hole may establish the tenor and distribution of the predicted greisen style mineralisation at the granite-wall rock contact and within the quartz-tourmaline aureole, however this would test the exploration concept of the mineralised zone only, rather than test a specific spot anomaly. Without a specific drill-worthy target no further work is warranted at Butler's Rd.

The historic Magnet Mine is a lode-style base metal and silver deposit hosted by a structurally emplaced mafic/ultramafic body known as the Magnet Dyke. The lower levels of the old mine, from 8 to 16 level (approx 160 to 365m below surface at licence boundary), are within EL1/2004 with a postulated feeder structure trending south-west into the EL. The periphery of the historic mine workings is prospective for residual Pb-Zn mineralisation while the host structure is prospective for repeat mineralisation but the recent channel sampling results provide no encouragement to continue prospecting to the south-west. Comstaff has commented that elevated surface tin geochemistry trending north-east from the mine might be due to leakage from depth up the fault structure. Mineral zonation may be present at Magnet with tin mineralisation underlying base metal mineralisation. No record has been found of tin assays having been taken within the mine levels. Whatever the case, tin mineralisation is likely to be at some depth, and difficult and expensive to evaluate. Funding is not available for deep exploratory work at the Magnet Mine.

5. ENVIRONMENT

For the period of the licence, all track cutting and sampling was in accordance with the MRT Exploration Code of Practice, the specific conditions of work programmes, and in consultation with the land manager, Forestry Tasmania Murchison District. No environmental issues remain outstanding from any work programmes.

6. EXPENDITURE

Rubicon Min Tech Ventures Pty Ltd	Printed At: 7/2/2018 2:28:31 PM	
EL 1/2004 Ramsay River		
Expenditure 10/02/2017 - 07/02/2018		
Rubicon Limited	Base Currency: AUD	Page: 1
Job Details	Department	Class
Posting Ref		
EL 1/2004 Ramsay River	D1	GROUP
105	STAFF COSTS	8207.43
150	Refund MRT rent Ramsay River	-100.96
	Total	8,106.47

7. REFERENCES

- Collins, P.L.F, 1983 Luina and Wombat Flat Exempt Areas: A review of previous exploration and a reconnaissance survey of an aeromagnetic anomaly. Tasmania Department of Mines Unpublished Report No 1983/35.
- Everard, J.L, 2003 Digital Geological Atlas 1:25,000 Series. Luina. Tasmanian Geological Survey.
- Gunston, J. & Turley, S.D. & Sumpton, J.D.H. 1984 A Report on the Exploration Activity for EL 6/83, Mount Cleveland. December 1983 to June 1984. MRT TCR 84-2156.
- Hazeldene, R.K. 2008 EL 1/2004, Ramsay River, Annual Report for the Period 10 January 2008 – 9 January 2009, Stellar Resources Limited.
- Hazeldene, R.K. & Rigg, A.M. 2009 EL 1/2004, Ramsay River, Annual Report for the Period 10 January 2009 – 9 January 2010, Stellar Resources Ltd.
- Hazeldene, R.K. & Rigg, A.M. 2010 EL 1/2004, Ramsay River, Annual Report for the Period 10 January 2010 – 9 January 2011, Stellar Resources Limited.
- Hazeldene, R.K. & Rigg, A.M. 2016 EL 1/2004, Ramsay River, Annual Report for the Period 10 January 2016 – 9 January 2017, Stellar Resources Limited.
- Morrison, K.C. & Rigg, A.M. 2011 EL 1/2004, Ramsay River, Annual Report for the Period 10 January 2011 – 9 January 2012, Stellar Resources Limited.
- McGunnigle, M.K. & Basford, P.W. 1997 Luina EL17/93 Joint Venture. Annual Report April 1996-March 1997, Pasmaenco Exploration, MPI Gold Pty Ltd. MRT TCR 98-4137.
- Morrison, K.C. & Rigg, A.M. 2014 EL 1/2004, Ramsay River, Annual Report for the Period 10 January 2014 – 9 January 2015, Stellar Resources Limited
- Morrison, K.C. & Rigg, A.M. 2015 EL 1/2004, Ramsay River, Annual Report for the Period 10 January 2015 – 9 January 2016, Stellar Resources Limited
- Palmer, K.G. 1975 Progress Report, Exploration Licence 1/63 for Six Months Ending August 11 1975, Cominco Ltd. MRT TCR 75-1115.
- Rigg, A.M. 2006 EL 1/2004, Ramsay River, Annual Report for the Period 10 January 2006 – 9 January 2007, Stellar Resources Limited.
- Rigg, A.M. 2007 EL 1/2004, Ramsay River, Annual Report for the Period 10 January 2007 – 9 January 2008, Stellar Resources Limited.
- Rigg, A.M. 2012 EL 1/2004, Ramsay River, Annual Report for the Period 10 January 2012 – 9 January 2013, Stellar Resources Limited.
- Rigg, A.M. 2013 EL 1/2004, Ramsay River, Annual Report for the Period 10 January 2014 – 9 January 2014, Stellar Resources Limited.
- Turner, N.J. 2006 EL 1/2004, Ramsay River, Report on 2006 Program, N.J. Turner Geological Services Pty Ltd. MRT TCR 07-5430A.

Keywords

Location:	Waratah – Luina, Butlers Road
Mineralisation environment:	stockwork veins, greisen,
Minerals:	cassiterite, galena, sphalerite, tourmaline, chromite, ilmenite
Exploration methods:	mapping, soil and rock chip geochemistry
Mine/prospect name:	Magnet Mine, Arthur Dam, Butlers Road, RY02, RY01
Stratigraphic name:	Luina Group, Meredith Granite
Lithologic name:	granite, schorl, basalt, volcanoclastic, siltstone, ultramafics
Geological Province:	Waratah-Luina, Tyenna Orogeny Stage 1
Geological age:	Cambrian, Devonian-Carboniferous, Tertiary