

# EXPLORATION LICENCE 46/2010

## Huskisson River, Tasmania

### NINTH ANNUAL PROGRESS REPORT

for the period 26 May 2019 – 25 May 2020



**Yunnan Tin Australia TDK Resources Pty Ltd**

15 Gubbuteh Road, Little Bay, NSW, Australia 2036

**Authors:** Jack Ma & Joe Xie

**Signed:**

A handwritten signature in black ink, appearing to be 'J Ma' followed by a flourish.

**Date:** May 2020

**Distribution:** Yunnan Tin Australia TDK Resources Pty Ltd  
Yunnan Tin Group, Yunnan, China  
Mineral Resources Tasmania

*Co-ordinate system used in maps and diagrams within this report is MGA55 (GDA94), unless otherwise specified.*

# Abstract

The licence area is considered prospective for Renison Style mineralisation, associated with Ordovician granite. The area is also being explored for Avebury-style nickel, PGM, chromite and to a lesser extent, gold. The most significant geological feature within the tenement is the Huskisson River Ultramafic Complex (HRUC) and a portion of Wilson River Ultramafic Complex (WRUC).

During the reporting period, an extensive review of drill data in neighbouring Renison Mine has revealed a potential extension of Renison Orebody into SW part of the tenement. This potential is also supported by VTEM data collected by Yunnan Tin Australia in 2013. Further review has led to planning a 1,050m angled diamond drill hole to verify the potential. An application for Tasmanian Government's EDGI drilling grant to support this drilling program has been made with Mineral Resources of Tasmania.

Petrological studies were carried out over two rock chip samples collected from south western tenement area. The results show they are not igneous rocks, and rather typical Chrisom Creek lithology of tuff.

Exploration expenditure on the licence was \$68,889 for the reporting period.

## Table of Contents

1. Introduction.....	3
2. Geology and Mineralisation.....	4
1.1. Regional Geology.....	4
1.2. Mineralisation and Exploration History.....	6
2. Exploration during the ninth year of tenure.....	9
2.1 Mineral potential assessment and drilling planning.....	9
2.2 Petrological studies of rock chip samples.....	15
3. Work Plan for Year Ten.....	16
4. Environment.....	17
5. Expenditure Statement.....	18
6. References.....	19

## Figures

Figure 1: EL46/2010 Huskisson River locality plan.....	3
Figure 2: Geology of EL46/2010 area (base map from MRT 25K geological mapping).....	4
Figure 3: Structure trend between Renison and Mt. Lindsay.....	9
Figure 4: Renison orebody mineralisation trend.....	10
Figure 5: Interpreted longitudinal section to NW of Renison Orebody.....	11
Figure 6: VTEM cvg_RTP magnetic contour over the area of interest.....	11
Figure 7: Various VTEM SFz frequency channels showing conductivity contours.....	12
Figure 8: Plan showing locations of cross sections and historical Renison drilling.....	13
Figure 9: Cross sections with interpreted geology and proposed drilling.....	14
Figure 10: Plan showing RN_01 location over MRT 25k geology.....	15

## Tables

Table 1: EL46/2010 Expenditure for current year.....	18
--	----

## Appendices

Appendix 1: Petrological report, by Dr. B. J. Barron (*in a separate digital file*)

## 1. Introduction

EL46/2010, Huskisson River, is located about 10 km west of Tullah, on the west coast of Tasmania (Figure 1). This tenement is found on Rosebery and Parsons 1:25,000 map sheets, with an area of 58 sq. kms.




Figure 1: EL46/2010 Huskisson River locality plan

Topography is moderately rugged, and notable topographic features include Lynch Hill in the northern part of the tenement and parts of the Huskisson River catchment. The bitumen HEC Pieman Road bisects the area, providing access to the central portion of the tenement. The northern part of the area can be accessed via forestry tracks connected to Boco Siding, whilst the southern part of the area has very limited access.

The principal land use includes State Forest, Regional Reserve, and Forest Reserve.

The area contains temperate rainforest, eucalypt woodland and relatively open button grass flats.

## 2. Geology and Mineralisation

### 1.1. Regional Geology

The major geological feature within the tenement is the Huskisson River Ultramafic Complex (HRUC), which is a part of similar ultramafic bodies scattered along the Dundas and Adamsfield in western and north-western Tasmania. One such ultramafic body is the Wilson River Ultramafic Complex (WRUC), located to the west of the HRUC, which is the largest exposed ultramafic body in the Dundas Trough with an area of approximate 25 sq. km. The Huskisson River Ultramafic Complex has a relatively smaller exposed area, about 3.5 sq. km (Figure 2).

These two ultramafic bodies may be continuous with each other beneath Silurian-Devonian sedimentary rocks exposed in the core of the Huskisson Syncline (Owen, 2005).

These two complexes have been collectively studied and explored in the past.




Figure 2: Geology of EL46/2010 area (base map from MRT 25K geological mapping)

Brown (1986) identified two petrogenetically distinct ultramafic successions within the WRUC and HRUC:

1) **Layered Dunite-Harzburgite succession (LDH)**, comprising fine to medium-grained, well-layered dunite, orthopyroxene-bearing dunite, and harzburgite composed of olivine, enstatite, chromite, and serpentine after the former silicate phases. Layering ranges from ca. 10 mm to 400 mm thick, with a primary bedding-parallel foliation defined by the primary alignment of enstatite and chromite crystals in the plane of bedding. There is also typically a later foliation defined by flattening of olivine crystals. Both olivine and orthopyroxene occur as cumulous phases, and chromite an accessory phase (1-5%) typically most abundant in the dunite layers. Discontinuous chromite laminations individually up to ca. 1-2 mm thick and 1-2 m long are locally present in the LDH.

Brown (1986) mentions the occurrence of PGE-rich chromite nodules in LDH of the Serpentine Ridge area. The western 100-150 m of the LDH in the Harman River area consists of interlayered dunite pyroxene-bearing dunite, and the eastern part layered harzburgite with minor thin dunite layers (Brown 1986).

2) **Layered Pyroxenite-Dunite succession (LPD)**, consisting of fine to medium-grained well-layered orthopyroxenite, olivine orthopyroxenite, and dunite. Layering is typically thinner than in the LHD, ranging up to 150 mm thick but mostly a few millimetres to 20 mm thick. Olivine and orthopyroxene dominate with accessory amounts of clinopyroxene (1-2%) and chromite (1-2%). Chromite is more common in the dunite layers. The layering sequence dunite-orthopyroxenite-dunite-orthopyroxenite is the most common, followed by dunite-orthopyroxenite-olivine orthopyroxenite-orthopyroxenite.

According to Brown (1986) serpentinite shears or faults separate the LDH and LPD everywhere and the original relationship of the two successions is unclear. The exposed parts of WRUC and HRUC are dominated by the LDH sequence. Two small, unfaulted blocks of LPD have been mapped by Brown (1986) at the north end of the WRUC (the Websterite Hill area), and the southern part of both complexes comprises LPD.

General interpretation is that the WRUC and HRUC are entirely fault bounded, the lower margins against Neoproterozoic and Early Cambrian volcanics and carbonates of the Crimson Creek Formation and correlatives, the upper margins against Devonian shallow-marine conglomerates, quartz arenites, siltstones and marls (Crotty Quartzite, Florence Quartzite, and Bell Shale), and locally slivers of the Cambrian Gordon Limestone. Radiometric dates are not available for the WRUC and HRUC and a broad Eocambrian to Cambrian age has been estimated according to stratigraphic constraints (e.g. Brown 1986). A major episode of folding during the Devonian formed

the north-west to north trending Huskisson Syncline, and contact metamorphism indicates emplacement of the WRUC and HRUC into the current stratigraphic position prior to the intrusion of the Meredith Granite around 370 Ma. Vein and replacement-style tin and tungsten mineralisation appears to be associated regionally with the intrusion of the Meredith Granite (Owen, 2005).

Although fault bounded, the prevalence of orthopyroxene over clinopyroxene, absence of protoclastic textures, and lack of stratigraphically associated sheeted dyke and pillow lava units suggests the WRUC-HRUC is not ophiolitic. Brown (1986) proposed intrusion of ultramafic bodies into the opening Dundas Trough during the Early Cambrian followed by tectonic reemplacment prior to the Devonian. The presence of serpentinite pebbles and abundant detrital chromite within Huskisson Group sedimentary rocks at Merton Hill and Red Lead Conglomerate of the correlative Dundas Group in the Mt Razorback area (Brown 1986) suggests exposure and partial erosion of the ultramafic complexes prior to the Middle Cambrian (Owen, 2005).

Quaternary fluvio-glacial sediments and Quaternary-Recent alluvial gravels cover much of the HRUC, and minor parts of the WRUC. Osmiridium, gold, and chromite are locally concentrated in the Quaternary-Recent alluvial gravels. Patches of laterite and saprolite are locally present over the WRUC in the Serpentine Ridge area, representing relicts of a more extensive lateritic cover developed during the Tertiary. Some lateritic nickel and cobalt mineralisation has been identified. Goethitic soils are widespread over Serpentine Ridge and the Websterite Hill area (Owen, 2005).

## **1.2. Mineralisation and Exploration History**

The licence area is considered prospective for nickel, Platinum Group Metals (PGMs), chromium and gold mineralisation, and has also been explored for tin, lead, zinc and silver.

Owen (2005) summarised mineralisation in the Wilson River and Huskisson River ultramafic complexes area in general and it was recited below.

Osmiridium, a rare naturally occurring alloy of the PGMs osmium and iridium, was first reported in Tasmania from the Wilson River valley in the 1876 by Surveyor-General Sprent (initially identified as palladium), and the Riley, Trinder, Three Mile, Lippy Jane, Fowler, Sweeney, Osmiridium and Gold creeks were later extensively worked for detrital osmiridium. An exact osmiridium production figure for the Wilson River area is not available, but of the total 31,100 oz produced from Tasmania between 1910 and 1968 (first and last reported production) around half came from the Adamsfield area ca. 120 km to the southeast and much of the rest from the Heazlewood-Bald Hill area near Waratah approx. 30 km to the north. Riley, Trinder, Three Mile, Lippy Jane, Fowler, Sweeney,

Osmiridium and Gold creeks were the most extensively worked for osmiridium in the Wilson River area. While there are some small test pits within serpentinite basement in the Riley Creek area the historic mining focussed on alluvial gravels in active creeks.

The detrital osmiridium typically occurs as flaky nuggets up to a few millimetres dimension, and petrographic work (Callina NL 1986-1990, Brown 1986) also indicates occurrence as inclusions within chromite grains from the ultramafic basement. Numerous workers have identified small chromite lenses up to 20-30 mm thick and 1-2 m long within the ultramafics, and analyses of some primary chromitites indicate highly anomalous PGM levels (Brown 1986).

There was additionally minor alluvial tin and gold production from the Wilson and Huskisson valleys and during the 1970's the area in the vicinity of the Meredith Granite was extensively explored for tin and tungsten mineralisation. Tin-bearing alluvials occur in many drainages on the north-eastern side of Serpentine Ridge, including Barnes, Sweeney and Tin creeks and Alfred River. Low-grade primary tin mineralisation occurs in the Harman River, Merton Hill, and Laurel Creek areas, and Reid (1932) makes reference to narrow dykes of tin-bearing quartz-feldspar porphyry cropping out in the vicinity of Tin Creek. Merton Hill was tested with 3 small adits by prospectors in the early 1900s, and later, 7 diamond drill holes (DDH MH1 to 7) by Renison Ltd (1980-1982). The drilling results were discouraging, the best intersection being 7.6m from 48.9m at 0.08% Sn, 0.76% Pb, 2% Zn and 36ppm Ag in MH1. The identified mineralisation was associated with veins and breccias within the Devonian Eldon Group (specifically, within the Crotty Quartzite and unnamed limestone member of the Amber Shale) associated with a north-east dipping fault zone adjacent to the contact with the Wilson River ultramafic body. Narrow granitic dykes with disseminated pyrrhotite were encountered in some of the drill holes at Merton Hill. Garnet skarns were identified in the Gordon Limestone around the confluence of Little Wilson and Wilson Rivers.

The source of the alluvial gold has not been thoroughly investigated but is in most cases probably reworked from glacial gravels. Significant gold mineralisation has not been reported from any of the identified tin prospects within the area, although it was not commonly assayed. Adit samples and some of the Renison drill core from the Merton Hill tin prospect (see above) was subsequently re-assayed for Au (Black Horse Mining, 1986-1987 and Cyprus Gold Australia Corp, 1987-1989) with a best result of 2m at 0.165ppm Au obtained in a magnetite skarn.

Lateritic nickel and cobalt mineralisation was identified in the southern Serpentine Ridge area by Aberfoyle in the late 1960s by a program that included hand auger drilling and man-portable coring (5 core holes) to a maximum depth of 30 ft. Grades of up to ca. 2% Ni and 1.5% Co were obtained from thin (<1-5m) patches of relict laterite and in the underlying saprolitic serpentinite assays of >0.5% Ni were commonly obtained. Sulfides were not observed. There was no systematic

investigation for Ni-sulphide mineralisation beyond the Serpentine Ridge – Riley Knob area (the Camp 30 area of Aberfoyle).

Callina NL (1985-1990) defined a detrital chromite resource in the Riley Creek area on the south western flank of Serpentine Ridge (the area that was also focus of the historic osmiridium workings). While the chromite is premium quality (>60% Cr<sub>2</sub>O<sub>3</sub>) the Callina resource was small (approx 1.7Mt at 1.9% chromite) and at the time not considered economic. The associated detrital PGM (Os and Ir, lesser Pt) and gold content were not assigned any economic value by Callina.

The last systematic exploration in the area was carried out by Adamus Resources Ltd during the period from 2003 to 2007, under EL18/2002. Work conducted during the period included stream sediment, soil and rock chip geochemistry. A heli-borne magnetic survey over Wilson River Ultramafic Complex was flown in 2005.

In 2007, Adamus carried out a substantial program of soil sampling with a 400m grid and a sample interval of 50m over the majority of the Wilson and Huskisson Ultramafic complexes. The grid was closed up to 200m line spacing over a previously identified electromagnetic anomaly on the north-western flank of Websterite Hill.

Fourteen lines were cut for approximately 267 samples over 22.5 km over the Huskisson River Ultramafic Complex, with a further 7 km of baseline also cut. Soil samples were assayed for Au, Ag, Cu, Pb, Zn, Ni, Co, Pt, Pd, Cr, Fe, Mg, Mo, Sb, Sn, W, Bi and S and pH was also determined for a number of samples. No significant results were reported (Grabham, 2007).

## 2. Exploration during the ninth year of tenure

Working during the reporting period consists of assessment of potential north-western extension of Renison Orebody into the tenement area, planning of drill test of Renison extension and petrological studies of rock chip samples from south-western area of the tenement.

### 2.1 Mineral potential assessment and drilling planning

#### Geological assessment

Following from field excursion in previous reporting period to the south-western area of the tenement, Yunnan Tin Australia has conducted analysis of Renison drill database and review of VTEM data flown in 2013, to assess the Renison style mineralisation in this part of tenement. Results so far indicates that Renison mineralizing system along the Federal Fault may extend north into south-western corner of EL46/2010.

The south-western part of EL46/2010 lies within the Federal-Bassett Faults Zone, which links Renison Mine and Mt. Lindsay deposit. This zone trends generally along contact of Crimson Creek Formation and Success Creek Group in a NW-SE direction (Fig. 3).




Fig. 3: Structure trend between Renison and Mt. Lindsay (from Venture Minerals Ltd.)

Plotting of up-to-date drill data from Renison Mine has demonstrated that the mineralisation trend has not followed the surface trace of Federal-Bassett Faults zone. It has however followed a straight line towards NW (Fig.4) and probably extended into area of interest within EL46/2010.




Fig. 4: Renison orebody mineralisation trend (Federal-Bassett Fault as red line)

Observations from Leatherwood area of Renison underground operations suggest that No. 2 dolomite on hanging wall seems to be associated with a flexure and/or folded structure and occurs away from the footwall of Federal Fault. With its shallow plunge updip, the approximate projection at surface would be around northern bank of Pieman River (personal communication 2015 with Francisco Navidad, Renison Mine Senior Resource Geologist then).

Based on the data from Renison mine and VTEM interpretation (discussed below), a longitudinal section NW of Renison has been proposed (Fig. 5).




Fig. 5: Interpreted longitudinal section to NW of Renison Orebody

## VTEM Review

Review of VTEM data flown by Yunnan Tin Australia in 2013 has also provided geophysical evidence along the predicated mineralization trend within the area of interest.

VTEM *cvg\_RTP* (*Calculated Vertical Derivative*) magnetic intensity (Fig. 6) exhibits a NW-SE variation trend, roughly following the projected Renison mineralisation trend (green line in Fig. 6). This VTEM data suggests considerable contrast of magnetic intensity along the trend and also a general increase of intensity with depth.




Fig. 6: VTEM *cvg\_RTP* magnetic contour over the area of interest

Conductivity based on various SFz frequencies (Fig. 7) suggests a broad NW-SW corridor of conductivity zone, spatially associated with the interpreted mineralization trend.




Fig. 7: Various VTEM SFz frequency channels showing conductivity contours  
(Channel number shown at the lower left corner of each contour; the higher the channel number, the deeper the conductivity it represents)

## Drilling Planning

Historically, resource exploration within Renison Mine has conducted a series of drillholes, outside SW corner of current area of interest, to evaluate the potential NW extension of known Renison mineralisation (Fig. 8). Those drillholes have been collared around intersection of mapped Federal-Bassett Fault and outcropping Renison Mine Sequence, with only minor mineralization intersected.

Yunnan Tin Australia believe that there might be a NW extension of Renison Orebody in a straight line, as discussed above.




Fig. 8: Plan showing locations of cross sections and historical Renison drilling

Based on discussion above, interpreted geology and mineralisation has been summarized in cross sections in Fig. 9 with Section S3-S3' showing proposed drill hole RN\_01.




Fig. 9: Cross sections with interpreted geology and proposed drilling

The details of the proposed drillhole, RN\_01, are listed below:

Hole No.	MGA55_E	MGA55_N	Elevation (m)	Dip	Azi-muth	Length(m)
RN_01	367,837	5,375,146	192	55	233	1,050

A plan view of proposed drillhole is plotted on top of MRT 1:25k geological map in Fig. 10 below.




Fig. 10: Plan showing RN\_01 location over MRT 25k geology

## 2.2 Petrological studies of rock chip samples

Two rock chip samples were collected from within the area of interest with their co-ordinates listed below during last reporting period. They were submitted to Dr. B. J. Barron in Sydney for petrological identification. The report is included in the report (Appendix 1).

Both rock samples are of tuff as part of typical Chrisom Creek lithology, and are not igneous as they appear.

Sample No.	MGA55_E	MGA55_N
837-S-02	368,141	5,375,695
837-S-01	367,916	5,375,683

### **3. Work Plan for Year Ten**

The proposed exploration work for Year 10 includes:

- Complete proposed drill hole RN\_01; and
- Central-east area of the tenement will be surveyed for Riley-type DSO iron ore resource. This work was planned for current reporting period and was not completed due to the COVID-19 travel restriction.

#### **4. Environment**

Yunnan Tin Australia TDK Resource Pty Ltd has environmental policies in place to always ensure minimisation of the impact that exploration activities have on the environment.

During the report year no surface disturbance occurred.

## 5. Expenditure Statement

A total expenditure of \$68,889 for the period 26/05/2019 to 25/05/2020 is detailed below:

<b>Expenditure</b>	<b>\$AUD</b>
Geology	\$58,269
Geochemistry	
Geophysics	
Remote Sensing	
Gridding	
Drilling	\$132
Land Access Costs	
Rehabilitation Costs	
Feasibility Study Cost	
Other Cost	\$4,226
Administration Cost	\$6,262
<b>TOTAL</b>	<b>\$68,889</b>

**Table 1:** EL46/2010 Expenditure for current year

## 6. References

- Black Horse Mining NL. Annual report to the Tasmanian Mines Department. Jordan, M., 1969. Camp 30 Report. EL2/1963. Aberfoyle Tin NL. Annual report for the Tasmanian Mines Dept.
- Bombardieri D. & Duffett M., 2018. Geophysics in the Surveys – News. Feb. 2018. AEGC 2018 Conference Handbook.
- Bombardieri D., Duffett M., Xie J. 2018. 3D models integrating geology and geophysics stimulate exploration in Tasmania. Australian Geoscience Council Convention, 14-18 Oct. 2018, Adelaide.
- Brown, A. V., 1986. Geology of the Dundas – Mt Lindsay – Mt Youngbuck Region. Tasmania Department of Mines. Geological Survey Bulletin 62.
- Callina NL 1986 Technical Report, Wilson River, NW Tasmania 1986. Annual report to the Tasmanian Mines Department 87-2633.
- Callina NL 1987 Annual Report 1/12/86 – 1/12/87 Wilson River Exploration Licence 24/85. Annual report to the Tasmanian Mines Department 87-2744.
- Callina NL 1990. Annual Report 1/12/1989 – 1/12/1990, Exploration Licence 24/85, Wilson River. Callina NL. Annual report to the Tasmanian Mines Dept.
- Everett, M.P., 1972. Huskisson Diamond Drill Hole No. 1, Exploration Licence 5/63. Comstaff Pty Ltd. Company Report to the Tasmanian Mines Dept 72\_0903.
- Creasy, N. W., 1987. John Lynch Creek Annual Report 1986/87. Annual report to the Tasmanian Mines Dept.
- Creasy, M. G., 1990. John Lynch Creek Annual Report 1988/89. Annual report to the Tasmanian Mines Dept.
- Davis, N., 1987. Annual Report EL14/86 Huskisson River Area Western Tasmania 1986-1987.
- Davis, N., 1988. Interim Report EL24/85 Wilson Annual Report EL14/86 Huskisson River Area Western Tasmania 1986-1987. Black Horse Mining NL. Annual report to the Tasmanian Mines Department 88-2879.
- Glasson, K. R., 1969. Report on the Trinder/Camp 30 area, Exploration Licence 2/63. Aberfoyle. Annual report to the Tasmanian Mines Dept.
- Grabham, G. 2007. Exploration report 2007 EL18/2002 Serpentine Ridge, NW Tasmania. Adamus Resources Ltd.
- Jessup, A., & Chenhall, B., 1968. Interim report on the Camp 30 Merton Area, Tasmania. Aberfoyle Tin Development Partnership. Annual report to the Tasmanian Mines Dept.

- Owen, S. 2005. 2005 Annual exploration report EL18/2002, Serpentine Ridge, NW Tasmanian. Adamus Resources Ltd
- Reid, A. M., 1932. Osmiridium in Tasmania. Tasmania Department of Mines, Geological Survey Bulletin 32.
- Roger Poltock Geological Pty Ltd (1987). Annual report for period Oct. 1986 – Sept. 1987, EL18/86, John Lynch Creek, Tasmania. Company Report to the Tasmanian Mines Dept 87\_2713.
- Venture Minerals Ltd. (2011): Venture discovers second DSO hematite prospect at Mt Lindsay, Tasmania. VMS ASX Announcement on 22<sup>nd</sup> Nov. 2011.
- Venture Minerals Ltd. (2012): Major high grade tin discovery, Tasmania. VMS ASX Announcement on 8th Aug. 2012.
- Xie, J. (2012). First Annual Progress Report, EL46/2010, for the period 26 May 2011 to 25 May 2012, Huskisson River, Tasmania. Unpublished report, Yunnan Tin Australia TDK Resources P/L.
- Xie, J. (2013). Second Annual Progress Report, EL46/2010, for the period 26 May 2012 to 25 May 2013, Huskisson River, Tasmania. Unpublished report, Yunnan Tin Australia TDK Resources P/L.
- Xie, J. & Veska, L.T. (2014). Third Annual Progress Report, EL46/2010, for the period 26 May 2013 to 25 May 2014, Huskisson River, Tasmania. Unpublished report, Yunnan Tin Australia TDK Resources P/L.
- Xie, J. & Veska, L.T. (2015). Fourth Annual Progress Report, EL46/2010, for the period 26 May 2014 to 25 May 2015, Huskisson River, Tasmania. Unpublished report, Yunnan Tin Australia TDK Resources P/L.
- Xie, J. (2017). Sixth Annual Progress Report, EL46/2010, for the period 26 May 2016 to 25 May 2017, Huskisson River, Tasmania. Unpublished report, Yunnan Tin Australia TDK Resources P/L.
- Xie, J. (2018). Seventh Annual Progress Report, EL46/2010, for the period 26 May 2017 to 25 May 2018, Huskisson River, Tasmania. Unpublished report, Yunnan Tin Australia TDK Resources P/L.
- Xie, J. (2019). Ninth Annual Progress Report, EL46/2010, for the period 26 May 2018 to 25 May 2019, Huskisson River, Tasmania. Unpublished report, Yunnan Tin Australia TDK Resources P/L.
- Yu, J. 2016. Fifth annual progress report, EL46/2010, Huskisson River, Tasmania, for the period 26 May 2015 – 25 May 2016. Yunnan Tin Australia TDK Resources Pty Ltd.