

200'							F/4/205
205'	<u>SANDY ARKOSIC SILT</u> white streaks of f'd pathic sandy silt in a predom brown silt		np	fine black rods of C1 ~5%	~8%		F/4/210
210'	7' f'd par ↓ 2 depth to a predom brown sandy silt at 215'						F/4/215
215'	<u>CARBONACEOUS SANDY SILT</u> rnd to sub rnd, fine (≤ 1/4 mm) predom qtz (~5%) in a brown silt matrix (80%) ↓ in silt matrix & T in C. content.		np	black to brown peat -10% + rare wood frag.	np.		F/4/220
220'				1 in % peat to 20-30%			F/4/225
230'							F/4/230
240'				peat -30-40% + ~5% soft black to dark brown wood frag.			F/4/235
250'	-some grey green sandy silt due to contamination			1 in % wood frag. to ~20%			F/4/240
260'				~20% peat < 3% small wood frag.			F/4/245
270'							F/4/250
280'							F/4/255
290'							F/4/260
300'							F/4/265
310'							F/4/270
320'							F/4/275
330'							F/4/280
340'							F/4/285
350'							F/4/290
360'							F/4/295
370'							F/4/300
380'							F/4/305
390'	<u>WEATHERED DOLERITE</u> Light grey clay after dolerite	+ +					F/4/310
400'		+ +					F/4/315
410'		+ +					F/4/320
420'		+ +					F/4/325
430'		+ +					F/4/330
440'	END						

