

DRILL ADVANCE				LITHOLOGY							
LOST CORE	DEPTH	DRILL ADVANCE INTERVAL	CORE RECOVERY	PERCENT RECOVERY	INTERVAL	DESCRIPTION	ALTERATION	GRAPHIC LOG	STRUCTURE	MINERALISATION	VISUAL PERCENTAGE MINERALISATION
	30.7	0.7	0.7	100%	31						
	31.4	3.1	3.0	96%	31.6 31.9				31.2 fracture / joint: 40° 31.7 quartz vein contact: 70°	31.8 Thin pyritic layer in quartz.	10%
	34.5	1.9	1.9	100%	35				35.6 Bedding to core axis: 45° 36.0 Bedding to core axis: 90°		
	36.4	1.1	1.1	100%	37						
	37.5	3.0	2.9	96%	37.0	Brecciated and deformed laminated chloritic, in part carbonaceous mudstone.	chlorite, ?epidote and minor quartz.		Breccia clasts up to 30mm. Laminar bedding folded and deformed.		
	39.0				39.0	Laminated chloritic in part carbonaceous siltstone (? tuffaceous). Thin 20 mm thick whitish carbonate bearing beds at 39.2m.	chlorite alteration.				
	40.8	2.1	2.1	100%	40.2	Laminated carbonaceous to chloritic siltstones to dark argillites. Some of the thin lighter coloured laminae have a small carbonate component (~5%).			41.6 Bedding to core axis: 55°		
	42.6	2.0	2.0	100%	42				42.4 quartz vein: 5° 43.4 Bedding to core axis: 60°	42.4 quartz vein with pyrite	5%
	44.6	3.0	3.0	100%	44.6				44.4 quartz vein: 45°		

SCALE 1:100 (1cm = 1 m)

COMSTAFF PROPRIETARY LIMITED

DRILLHOLE LOG FOR DDH RBE 7

LOGGED BY \_\_\_\_\_ FROM 30 TO 45

DATE 24 / 5 / 80

PAGE 3 OF 13