


LITHOLOGY

DRILL ADVANCE				LITHOLOGY							
LOST CORE	DEPTH	DRILL ADVANCE INTERVAL	CORE RECOVERY	PERCENT RECOVERY	INTERVAL	DESCRIPTION	ALTERATION	GRAPHIC LOG	STRUCTURE	MINERALISATION	VISUAL PERCENTAGE MINERALISATION
	136.5	3.0	2.8	93%	136.5 - 137.5	zone of intense carbonate veining in core					
	139.5	2.6	2.7		137.5 - 140.1	Massive dull green-grey medium to fine tuffaceous sandstones with thin interbedded siltstone and black argillite horizons.			138.5 Bedding to core axis: 30° facing up the hole from graded bedding.		
	142.1	3.1	3.1	100%	140.1 - 143.2				41.1 5mm thick carbonate vein : 70°		
	145.2	3.1	3.1	100%	143.2 - 146.3						
	148.3	3.1	3.1	100%	146.3 - 149.4	100mm zone of thin carbonate veins. Carbonate tends to develop in the core both as cement and small veinlets.			147.1 Bedding to core axis: 25° 147.5 Vein : 50°		147.3 Pyrite disseminations and concentrations occur parallel to bedding. Locally concentrations may achieve 15%.

SCALE 1:100 (1cm = 1m)



COMSTAFF PROPRIETARY LIMITED

DRILLHOLE LOG FOR DDH RBE 7

LOGGED BY FROM 135 TO 150

DATE 24/5/80

PAGE 10 OF 13