

DRILL ADVANCE				LITHOLOGY							
LOST CORE	DEPTH	DRILL ADVANCE INTERVAL	CORE RECOVERY	PERCENT RECOVERY	INTERVAL	DESCRIPTION	ALTERATION	GRAPHIC LOG	STRUCTURE	MINERALISATION	VISUAL PERCENTAGE MINERALISATION
	45.1	2.1	0.85	40.4%			of mild hydrothermal alteration.		45.1 Fracture/fault zone.		
	47.2	1.2	0.96	80%		Broken and fractured siltstone to fine grained sandstone.	Limonite staining along fracture planes. Minor quartz veins at 47.9m.		48.0 Core badly fractured and broken.	47.9 Pitted and oxidised zone, possible leached sulphides?	
	48.4	1.4	1.26	90%							
	49.8	0.9	0.9	100%	50						
	50.7	1.2	1.2	100%	50.7-51.7	black carbonaceous siltstone to fine tuff sandstone. Presence of thin laminate bedding in part of the core.	Core shows evidence of mild hydrothermal alteration. Infrequent thin leached and partially limonite-stained quartz veins.		50.7 Bedding to core axis: 45°	Very minor pyrite scattered through the core or concentrated on fracture zones.	40%
	51.9	0.5	0.5	100%	52						
	52.4	0.9	0.9	100%	53	Dull greenish-grey, fine-medium grained buffaceous sandstones and minor, in part, carbonaceous siltstones. Core is broken and fractured in places.					
	53.2	0.6	0.6	100%							
	53.9	0.2	0.2	100%	54						
	54.1	0.5	0.5	100%							
	54.6	1.2	1.15	96%	55						
	55.8	0.5	0.5	100%	56						
	56.3	1.7	1.3	76%	57						
	58.0	1.7	1.7	100%	58						
					59						
					59.5						
	59.7	0.6	0.6	100%	59.7-60	Prominent quartz-sulphide vein.	Part oxidised and leached		quartz vein: 530	Pyrite altered sulphides	10%

SCALE 1:100 (1cm = 1 m)

