


DRILL ADVANCE				LITHOLOGY					VISUAL PERCENTAGE MINERALISATION		
LOST CORE	DEPTH	DRILL ADVANCE INTERVAL	CORE RECOVERY	PERCENT RECOVERY	INTERVAL	DESCRIPTION	ALTERATION	GRAPHIC LOG	STRUCTURE	MINERALISATION	VISUAL PERCENTAGE MINERALISATION
	135.0	1.0	1.0	100%							
	136.0	3.0	3.0	100%	135.2 - 136.2	Massive tuffaceous fine-grained sandstone with very minor associated mudstone. Very minor quartz-carbonate veins cut the core.			135.9 B.C.A: 40°		
	139.0	3.0	3.0	100%	139.0 - 139.0				139.7 calcite vein, 10 mm; 40°		
	142.0	0.8	0.8	100%	142.0 - 142.0						
	142.8	2.2	2.2	100%	143.1 - 143.1	Finely laminated - in part brecciated, pale-medium grey mudstone.			143.7 B.C.A: 55°		
	145.0	3.0	3.0	100%	145.0 - 145.0						
	146.0	2.2	2.2	100%	145.9 - 146.1	Deformed, in part brecciated, black carbonaceous mudstone. Numerous thin pygmatic quartz-carbonate veins cut the core.			147.3 Intense quartz-carbonate veining: 55°		
	148.0	3.0	3.0	100%	148.0 - 148.0						
	149.0	3.0	3.0	100%	149.0 - 149.0						
	150.0	3.0	3.0	100%	150.0 - 150.0						

SCALE 1:100 (1cm = 1m)



138.1 Thin 10mm brecciated zone with minor chalcopyrite. <1%

Pyrite concentrations associated with deformed veins in black argillite. Some of the pyrite is streaked out along deformed bedding planes. Locally pyrite may achieve concentrations of ~10%. 2%

149.8 100 mm quartz, pyrite, chalcopyrite, arsenopyrite vein. 10%

COMSTAFF PROPRIETARY LIMITED

DRILLHOLE LOG FOR DDH RBE 8

LOGGED BY FROM 135 TO 150

DATE 6-7/6/80

PAGE 10 OF 21