


DRILL ADVANCE				LITHOLOGY							
LOST CORE	DEPTH	DRILL ADVANCE INTERVAL	CORE RECOVERY	PERCENT RECOVERY	INTERVAL	DESCRIPTION	ALTERATION	GRAPHIC LOG	STRUCTURE	MINERALISATION	VISUAL PERCENTAGE MINERALISATION
	15.4	1.0	0.3	30%				15.3 60°B			
	16.4	1.5	0.5	33.3%							
	17.9	1.5	0.4	26.6%							
	19.4	0.3	0.3	100%	19.4	Feldspathic sandstone, tuffaceous, medium grained with interlayered black carbonaceous shale. Black shale component increases down hole. Sheared lithic fragments of shale occur in sandstone.	Soft, possibly minor clay alteration.	Massively bedded.	Ry in shale intercalations. Limonite staining in sandstone and xaln Ry in fractures.	1%	
	19.7	0.2	0.2	100%							
	19.9	0.8	0.5	62.5%							
	20.7	0.8	0.4	50%							
	21.5	0.9	0.56	62.2%							
	22.4	1.2	0.9	75%							
	23.6	0.5	0.5	100%							
	24.1	1.1	0.5	45.4%				24.0 20°S			
	25.2	2.2	0.65	29.5%	25.3	Pyritic black shale, with very minor green tuffaceous sandstone interbeds.	End of weathering effects. Graphite developed on cleavage planes.	Well laminated.	Xaln Ry concentrated along bedding planes or remobilized into vuggy patches.	5%	
	27.4	1.0	0.4	40%				27.5 60°B			
	28.4	1.5	0.5	33.3%							
	29.9	0.4	0.4	100%				30.0 100°B			



SCALE 1:100 (1cm = 1 m)