

## DIAMOND DRILL RECORD

HOLE NUMBER : BT 132

LOGGED BY : AFR

NWPS

INTERVAL (m)		RECOVERY		DESCRIPTION	FORM.	% Sn.										
FROM	TO	m	%			FROM	TO	TOTAL	ACIDSOL.	% Cu.	% As.	% Mn	% Pb.	% Zn.	% Bi.	g/t Ag
				SUMMARISED LOG												
0	2.6			NON-CORING.												
2.6	18.8			ALKALI GRANITE WITH ZONES OF INTENSE HEMATITE ALTERATION, AND VARIABLE ZONES OF GRANITE GREISEN AND MINOR GRANULAR GREISEN (ANCHOR GRANITE).												
18.8	44.0			ALKALI GRANITE VARYING TO GRANITE-GREISEN, MINOR GRANULAR GREISEN.												
				DETAILED LOG												
						5	6	0.01	0.0025	0.025	0.0050	<1				
							7	40.01	0.0030	0.025	0.0035	1				
0	2.6	0.0	0.0	Coring but no recovery.			8	0.10	0.0025	0.03	0.0050	<1				
							9	0.04	0.0025	0.045	0.0055	1				
2.6	5.6	1.2	40.0	Weathered pinkish-cream, sericitised medium grained alkali granite (although may be close to P.A. contact, as there is weathered P.A. in the bank adjacent to the drill site).			10	0.24	0.0020	0.06	0.0105	<1				
							11	0.15	0.0015	0.135	0.0135	1				
							12	0.36	0.0015	0.085	0.0125	<1				
							13	0.18	0.0015	0.09	0.0135	<1				
5.6	5.9	0.3	100	Broken pegmatite/(quartz).			14	0.03	0.0015	0.045	0.0060	1				
							15	0.12	0.0015	0.06	0.0095	1				
5.9	6.1	0.2	100	Zone of layered pink aplite to coarser pegmatite. Definite layering at 40-45° CA. Slight lime green sericite.			16	0.06	0.0020	0.165	0.0255	2				
							17	0.04	0.0015	0.23	0.0315	2				
							18	0.25	0.0025	0.27	0.0430	1				
6.1	8.7	2.6	100	Grades into pink-white medium grained alkali granite. Common weathering as limonitic joints, patches etc. Weak sericitisation. Disseminated dark biotite.			19	0.05	0.0015	0.09	0.0150	1				
							20	0.17	0.0030	0.12	0.0210	1				
							21	0.06	0.0015	0.065	0.0090	<1				
							22	0.63	0.0015	0.065	0.0080	<1				
8.7	10.9	2.2	100	Grades into very limonitic and sericitised (for 60cms) of greisenised alkali granite. Then fresher granular greisen-(granite) with pervasive hematite staining.			23	0.03	0.0010	0.035	0.0065	<1				
							24	0.03	0.0010	0.04	0.0060	<1				
							25	0.04	0.0010	0.035	0.0055	<1				
							26	0.08	0.0010	0.035	0.0055	<1				
10.9	12.3	1.4	100	Grades into complex zone of altered alkali granite. Clayey orange material for 10cms, then cream and hematite spotted, stained granular alkali greisen (granite). The hematite replaces coarse micas.			27	0.01	0.0010	0.035	0.0055	<1				
							28	<0.01	0.0010	0.035	0.0050	1				
							29	0.01	0.0010	0.030	0.0055	<1				
							30	40.01	0.0015	0.030	0.0055	<1				
12.3	13.1	0.8	100	Grades into grey-green granular greisen with sparse disseminated hematite.			31	"								
							32	0.01								
							33	0.02								
13.1	16.7	3.6	100	Complex, coloured and variable. Gradational cream-grey granite-(greisen) to a more pink pervasive siliceous granular greisen. Then into a pronounced scarlet granular rock with bright green coarse mica. Locally there are patches of coarse micas in the last 60cms.			34	0.01								
							35	"								
							36	"								
							37	"								

892130