

## APPENDIX C (Continued)

4. Costean 4 L22N (see figure 12)

Sampled from E end of costean	Cu ppm	Pb ppm	Zn ppm	Ag ppm	Mn ppm	Ba (sol) ppm
0- 5 m	435	30	140	< 2	650	< 100
5- 10	200	50	150	-	650	-
10- 15	185	70	135	-	320	-
15- 20	315	30	145	-	340	-
20- 25	285	20	50	-	90	-
25- 30	115	30	35	-	100	-
30- 35	95	20	45	-	130	-
35- 40	120	30	100	-	250	-
40- 45	170	70	55	-	110	-
45- 50	85	90	50	-	170	-
50- 55	85	70	55	-	240	-
55- 60	105	50	75	-	300	-
100-105	110	160	225	-	130	-
105-110	170	70	220	-	120	-
110-115	105	50	70	-	140	-
115-120	165	70	150	-	200	-
120-125	95	50	105	-	220	-
125-130	110	30	90	-	160	-
10 m (30983) (Resampled)	70	40	80	-	180	-
0- 8 m	55	30	130	-	840	-
8- 10	75	100	130	-	380	-
10- 12	95	190	75	-	70	-
12- 15	130	80	200	-	590	-
30- 32	40	30	40	-	130	-
32- 37	65	20	100	-	320	-

5. Costean 5 100' north of L23.4N 1700W  
Previously sampled 1979 (see results Meares et al, 1980)

Sampled from E end of costean	Cu ppm	Pb ppm	Zn ppm	Ag ppm	Mn ppm	Ba (sol) ppm
0 -10'	390	<10	10	< 2	20	<100
10'-20'	190	<10	150	< 2	1600	<100
20'-30'	360	10	70	2	710	-
30'-40'	420	70	85	2	11000	-
40'-50'	200	60	65	<2	7100	-
50'-60'	140	130	180	4	49000	200
60'-70'	120	40	60	<2	5200	-
70'-80'	40	50	40	<2	1200	-