

HSK.504	C2/4 353,025.E 5,358,260.N	fine W, micro peg' with tourm
HSK.505	C2/4 353,125.E 5,358,075.N	porph m-c W, g'mass minor, felds phenos to 20mm dom.
HSK.506	C2/4 352,870.E 5,358,700.N	porph pale pink/brn, felds > (red 10mm) qtz.
HSK.512	C2/2 351,390.E 5,360,850.N	med W. med R/W see append (ii)
HSK.517	A3/4 344,100.E 5,365,200.N	see append (ii) med W with tourm
HSK.518	A3/4 343,725.E 5,365,575.N	porph fine W g'mas, biot prom, F > Q with nodules
HSK.519	A3/4 342,525.E 5,365,650.N	coarse W see append (ii)
HSK.520	A3/4 342,160.E 5365,675.N	see append (ii)
HSK.529	A4/3 344,875.E 5,364,750.N	porph fine W g'mas, F > Q
HSK.535	A4/4 349,650.E 5,365,100.N	see append (ii)
HSK.539	C2/2 349,995.E 5,362,160.N	see append (ii) m-c R/W
HSK.546	C2/1 349,400.E 5,361,500.N	med R/W, minor qtz. felds phenos
HSK.547	C2/1 349,475.E 5,361,325.N	med-C R/W fine W scattered qtz felds phenos - 11mm
HSK.554	C2/3 347,950.E 5,358,670	med-c W ~ chlorit' biot' see append (ii)