RGC EXPLORATION PTY. LIMITED

083001

MICROFILMED

GEN

E.L. 12/90 AND E.L. 15/90

WARATAH PARTIAL RELINQUISHMENT REPORT

FOR THE PERIOD 1990 TO 1992

92-3357.

RECEIVED

RECEIVED

Covering letter
on EL 15/90
Folio 77
EL 12/90 Folio 4

Compiled by: S.W. Halley
Geologist

Report No. T/92/9

Endorsed by: M.J. Fleming
Senior Exploration Geologist

May, 1992

Distribution:
- Tasmania Mines Department (1)
- RGC Exploration (1)

SUMMARY

E.L.s 12/90 and 15/90 were granted to Renison Ltd., as a result of a successful tender application during 1990. The leases cover a combined area of 332 km^2. The area was acquired because of its potential for carbonate replacement tin deposits of the type mined at Mount Bischoff and Luina.

The E.L.s cover the northern and eastern margins of the Meredith Granite and the surrounding rocks, which includes Oonah Formation sediments, Crimson Creek volcaniclastics, Gordon Limestone and Eldon Group sediments.

Work completed by RGCE to date included a literature review of previous exploration work, reassessment of the existing stream sediment geochemistry database, reassessment of the 1981 Mines Department aeromagnetic data, regional mapping and rock chip sampling and a regional gravity survey. In 1991, as a result of a cut in RGCE's tin exploration budget, further work in these EL's concentrated on specific prospects rather than work of a more regional nature. Budgetary constraints also necessitate this partial relinquishment of the EL's. Because of the intensely magnetic rocks around the south eastern margin of the granite and the basalt cover along the eastern side of the EL's, these areas are perceived to be more difficult exploration areas, and have subsequently been relinquished.

CONTENTS

SUMMARY

1. INTRODUCTION
2. LAND TENURE
3. PREVIOUS WORK
   - 3.1 Comstaff EL 5/63
   - 3.2 Goldfields Exploration EL 17/77
   - 3.3 Aberfoyle EL 16/78
   - 3.4 Billiton EL 46/88
   - 3.5 Soloriens EL 59/88
   - 3.6 Department of Mines
4. STRATIGRAPHY
   - 4.1 Oonah Formation
   - 4.2 Crimson Creek Formation
   - 4.3 Dundas Group
   - 4.4 Ordovician-Devonian Sediments
   - 4.5 Tertiary Basalt
   - 4.6 Meredith Granite
5. STRUCTURE
6. WORK COMPLETED
   - 6.1 Geology
   - 6.2 Geochemistry
   - 6.3 Geophysics
7. RESULTS
8. CONCLUSIONS
9. REFERENCES

APPENDICES

1. Interpretation of regional aeromagnetic data from the Waratah-Mt. Ramsay area, Tasmania.
2. An interpretation form of Waratah Granite - D. Leaman.
3. Rock Chip Geochemistry

PLANS

PLAN	DRAWING NO.	TITLE	SCALE
1	5530/033	Meredith Project EL 12/90 & EL 15/90 Area to be relinquished	1:50000
2	5530/029	Factual Geology Sheet 8	1:10000
3	5530/031	Factual Geology Sheet 11	1:10000
4	5530/017	Waratah Area bouger anomaly D. Leaman, Map 1	1:50000
5	5530/018	Waratah Area residual anomaly D. Leaman, Map 2	1:50000
6	5530/019	Waratah Area Form of Meredith Granite D. Leaman, Map 3	1:50000

1. INTRODUCTION

EL 12/90 was granted in 1990 as a result of a successful tender application. An adjoining area also applied for was granted as EL 15/90. The combined area of the leases is 332 km^2. The EL's are regarded as being geologically continuous and have been explored and reported as one single block. The licences are held by Renison Limited and explored by RGC Exploration, both wholly owned subsiduaries of RGC Limited.

Much of the area is vacant crown land. It is covered by wet schlerophyll forest with patches of rain forest, ti-tree scrub and button grass plains. Access is provided by the Corinna Road, numerous logging tracks and old exploration tracks and by newly-cut walking tracks. Much of the area is accessible only by foot.

Work on the EL's has concentrated on locating tin mineralisation of the style present at Mount Bischoff and Cleveland. Leaman and Richardson (1989) highlighted the potential of this area by demonstrating the relationship between the shape of buried granite bodies and the associated mineralization, in particular the north-east trending ridge of granite beneath Mount Bischoff. Carbonate units are known to occur within the Oonah Formation and in the Crimson Creek Formation. Gordon Limestone also occurs in proximity to the Meredith Granite.

Initially a literature study was conducted to assess the quality and results of previous exploration. Regional mapping and rock chip sampling was a major part of this years work, providing a framework for assessment of geophysical and geochemical data and identifying areas requiring more detailed follow-up work.

2. LAND TENURE

In March 1990, Renison Limited tendered for ETA 160 "Mt. Ramsay" and ETA 161 "Waratah". The tender application was successful and EL 12/90 with an area of 149 km^2 was granted on 6/7/90. A further 183 km^2 was also applied for and this area was granted as EL 15/90 on 6/7/90. EL 12/90 includes the township of Waratah. Excluded from EL 12/90 is RL 8807 which covers 4 km^2 around Mount Bischoff and several small MLs over alluvial deposits on the Waratah River and near the margins of the Meredith Granite. These ML's are:

19M/72	Campelane Nominees,	Waratah River alluvials
44M/90	A. Sporer,	Wombat Flat
11M/77	A. Sporer,	Wombat Flat
14M/77	A. Sporer,	Wombat Flat
5M/75	Seaborn Pty. Ltd.	Wombat Flat
82M/77	M.G. Glozier	below Waratah Falls

There are also a number of water easements associated with 19M/72.

Following relinquishment, EL 12/90 will be reduced to approximately 105 km^2, excluding the above mentioned areas and EL 15/90 will be reduced to 36 km^2 (plan 1).

3. PREVIOUS WORK

3.1 Comstaff EL 5/63

Comstaff held EL 5/63 covering much of the present EL's 12/90 and 15/90, for a period of twenty five years from 1963 to 1988. During this period, a vast multitude of reports were produced. A summary of work completed in the relevant areas is contained in the report by BHP (1988), including a complete list of references, and in Shaw and Everett, (1985a, b).

Work in the Magnet-Bishoff area started in the early 1970's with extensive stream sediment sampling and geological mapping. This was followed by a regional TURAM EM survey. A prominent tin anomaly was reported from Deep Gully Creek. Three grids were cut over the TURAM anomalies, but CRONE EM and magnetic surveys revealed nothing of interest. An enlarged grid was cut between Magnet and Bischoff. One EM anomaly was drill tested with disappointing results. Two drill holes at Magnet intersected old workings hence missed the vein.

A belated follow-up of the Deep Gully Creek anomaly in 1981 led to the discovery of mineralised Devonian quartz-porphyry float and unmineralised outcropping dykes, identical to the intrusives at Mount Bischoff. The dykes intrude Oonah Formation, near the Precambrian-Cambrian boundary about 3km northeast of Mount Bischoff. Comstaff followed up this discovery with gridding, soil geochemistry, geological mapping, ground magnetics and aeromagnetics.

Two sources of anomalous tin geochemistry were identified:
(i) Tin occurs in Tertiary sediments which lie between basalt flows; this tin was most likely shed from Mount Bischoff, and
(ii) Blocks of porphyry, particularly in Ethol Creek, are altered to mineralized topaz greisen typical of the Bischoff mineralization. However the outcropping porphyries are strongly sericitized but unmineralized. It is possible that the topaz

greisen float originated from Mount Bischoff, but since outcropping porphyries occur in the area, a local source seems more likely.

In 1984 five holes were drilled through the Tertiary cover. Four of the holes intersected dolomite within the Precambrian basement. The dolomite lies along the faulted contact between the Oonah and Crimson Creek Formations. The source of the tin anomalous porphyries was never identified.

There was a regional DIGHEM survey in 1983 which identified significant anomalies in the Deep Gully Creek area, but none of these were ever followed up. One of these anomalies occurred along the faulted contact between the Oonah dolomite and the Crimson Creek Formation, but outside Comstaff's EL. Although further drilling was recommended, no more work was done on this prospect after the initial five holes were completed.

The Butler's Road anomaly was selected from the 1983 DIGHEM data. It occurs at the north eastern tip of the granite where tourmaline-veined, greisenized granite is surrounded by hornfelsed sediments. BHP joint-ventured into EL 5/63 in 1985 and subsequently drilled one short hole in this area, which returned anomalous tin and base metal values.

Comstaff started exploring the eastern margin of the Meredith Granite in 1969 with a regional stream sediment and reconnaissance mapping program. Some of the anomalies were gridded and covered by soil sampling. This led to the Will O'Wisp target area where subsequent drilling intersected Precambrian dolomite and pyritic black shales. An INPUT survey in 1975 identified 8 conductors. These were investigated by gridding, soil sampling, ground EM and magnetic surveys and geological mapping. Five of these prospects were subsequently drilled. The most interesting of these was a skarn on the foothills of Mount Ramsay which however had very disappointing assays. Comstaff also investigated the old South Bischoff mine.

Five holes were drilled, intersecting very narrow greisen zones within the granite.

3.2 Goldfields Exploration EL 17/77

EL 17/77 covered the southern margin of the Meredith Granite, including the northern end of the Husskisson Syncline, now included within EL 15/90. Goldfields Exploration followed up stream sediment geochemical anomalies around the northern edge of the Huskisson syncline with soil and bedrock geochemical surveys, ground magnetics, IP, and drilling. This work defined a zone of magnetite skarns over 4 km in length in the Gordon Limestone along the granite contact. Four drill holes successfully tested the skarn, with a best result of 7.0m at 0.65% Sn. No further work was done here due in part to the complexity of the tin mineralogy.

3.3 Aberfoyle EL 16/78

EL 16/78 covered the northern margin of the Meredith Granite, south of the Cleveland Mine. Stream sediment geochemistry led to the discovery of a tungsten-bearing skarn and a tin-bearing skarn, both outside the area of the current EL 12/90. Interest was drawn to a third area by an anomaly from the Department of Mines 1981 aeromagnetic survey. In the Whyte River area, the Gordon Limestone - Eldon Group sequence has a faulted contact with Cambrian andesites to the east and is intruded by the Meredith Granite to the south. The magnetic anomaly apparently occurs on the contact between the granite and the limestone. Tremolitic calc-silicate rocks were reported to be possible skarn material, but more likely were hornfelsed Cambrian andesites from the vicinity of the granite contact. Aberfoyle located a stream sediment tin anomaly coincident with the magnetic anomaly. A small grid was cut however no mention of this grid, or of any work carried out on it was recorded in any Aberfoyle reports. It is possible that following the cutting of the grid, a decision to drop the EL was made before any work was completed. The

source of the magnetic and geochemical anomalies was never identified.

3.4 Billiton EL 46/88

EL 46/88 covered much of the northern part of EL 5/63, relinquished in 1988 by Comstaff-BHP. Billiton concentrated on a combined aeromagnetic, EM anomaly identified by the 1983 Comstaff DIGHEM survey. This occurs at Stonedam Creek, a few km E of Waratah, on the edge of the basalt plateau. The area was gridded, geologically mapped and surveyed with ground magnetics and UTEM. A drill hole to test the anomaly had background levels of tin and base metals.

3.5 Soloriens Mining P.L. EL 59/88

EL 59/88 covered the northern and eastern margins of the Meredith Granite. David Leaman had begun an interpretation of the 1981 Mines Department aeromagnetic data when financial constraints resulted in the EL being surrendered. No field work was done.

3.6 Mines Department

The area from Zeehan to Waratah was the subject of a major regional mapping project undertaken by the Department of Mines. This work was reported in the Geological Survey Bulletin 62 (Brown, 1986). This is a detailed account of the geology, stratigraphy and petrology of the region.

4. STRATIGRAPHY

4.1 Oonah Formation

Within the exploration area, there are two blocks of Oonah Formation, one surrounding Mt. Bischoff and the other in the Ramsay-Coldstream area. Drilling through Tertiary Basalt suggests that these two blocks are continuous beneath the basalt cover. The Proterozoic Oonah Formation contains pale grey quartz sandstones, generally finer-bedded pale grey siltstones, dark grey shales, dolomites and minor mafic lavas and volcanoclastics.

The Oonah has been divided into upper and lower successions on the basis of lithology. The Lower Oonah is dominated by micaceous quartz sandstones and siltstone with minor interbedded phyllitic mudstone. The Upper Oonah has a greater abundance of mudstone and shales, with dolomite, mafic volcanics and relatively minor sandstone.

The Oonah Formation in the Ramsay block can be divided into three zones. The central zone is dominated by quartz sandstone in beds 10 to 50 cm thick. On lithological criteria, this is correlated with the Lower Oonah. In the western zone there are thinnly bedded calcareous siltstones and shales. In the eastern zone there are also siltstones, shales and dolomite. Although these two zones are correlated with the Upper Oonah, no mafic rocks have been noted in this area. The Upper Oonah rocks tend to be finer grained and thinner bedded than those of the Lower Oonah. They are less competent and more ductile and are often strongly deformed, with abundant parasitic and short wavelength folds. A curious outcrop of very pure quartz occurs on the Coldstream Road. It is composed of saccharoidal quartz with patches of clear coarse quartz crystals. This may be a silicified dolomite. Silicification is quite common in the Precambrian dolomites.

Around Mt. Bischoff, a similar association is observed. Dolomite occurs with shale and siltstone rather than in sandstone-

dominated successions. As well as at Mount Bischoff, dolomite has been mapped at Deep Gully Creek and on the northern side of the Waratah River.

4.2 Crimson Creek Formation

The Crimson Creek Formation occurs along the eastern side of the Meredith Granite, around the Cleveland mine area and north of Mount Bischoff. The sequence is composed largely of basaltic volcanoclastic turbidites and finely bedded siltstone and mudstone. Basalts and chert are quite common in the northern areas. Thin carbonate horizons occur within the Crimson Creek Formation but rarely outcrop. Chemically, the basalts are tholeiitic in character (Brown, 1986). Due to the basaltic component within the sediments, the Crimson Creek rocks are usually deeply weathered and outcrop is generally poor.

4.3 Dundas Group

A correlate of the Dundas Group occurs in EL 15/90 along Huskisson Drive. This correlation is made on the basis of lithology and the presence of ultramafic and mafic detritus. The sequence contains conglomerate, interbedded with lithic wacke, siltstone and mudstone. The conglomerate is calcareous, with clasts of chert, basalt, mudstone and pyroxenite. It has a calcareous, sandy matrix. These rocks are faulted against the Oonah Formation to the west.

4.4 Ordovician - Devonian Sediments

A sequence of Ordovician to Devonian sediments unconformably overlies Cambrian mafic to ultramafic rocks within the Huskisson Syncline and in a smaller syncline north of the Meredith Granite between Mt. Stewart and Heazlewood. These sediments belong to the Gordon Limestone - Eldon Group sequence and are part of the same sequence that occurs near Zeehan, Queenstown and in the King River Valley. The Gordon Limestone occurs at the base of the

sequence and may have a thickness of up to 500m. It generally does not outcrop. Alluvium filled valleys tend to develop above the limestone. The Gordon Limestone is overlain by the Crotty Sandstone, a white, friable quartz rich sandstone, up to 400m thick. This sandstone unit tends to outcrop as resistant ridges. The Crotty Sandstone is overlain by calcareous laminated siltstone and mudstone, the Amber Slate. This also tends to be a poorly outcropping unit. Successive units include the Keel Sandstone, the Austral Creek Siltstone, the Florence Sandstone and the Bell Shale. In the Whyte River area, only that part of the sequence from the Gordon Limestone to the Amber Slate has been mapped. The Crotty Sandstone forms a prominent ridge around the edge of the syncline, but the rest of the sequence is poorly exposed, most of it covered by alluvium.

IGNEOUS ROCKS

4.5 Tertiary Basalt

An extensive basalt plateau covers the area east and south of Waratah with erosional remnants to the west. Individual flows range from less than 1m to greater than 10m thick. Fluvial and lacustrine sediments occur between the basalt flows. The sediments range from mud to gravel and are often poorly consolidated. The Tertiary cover is up to 300m thick.

4.6 Meredith Granite

The Meredith Granite has been radiometrically dated at 356 Ma (with revised decay constants). It has a wide range of textural variety. Around the north-eastern tip and eastern margin, the granite is porphyritic close to the contact, with feldspar phenocrysts up to 25mm long and quartz phenocrysts up to 8mm. Moving away from the contacts towards the centre of the granite, the rock becomes less porphyritic and more equigranular. It is biotite-bearing throughout. Along the northern margin, south of the Whyte River, coarse-grained equigranular biotite granite

occurs right up to the contact. Zones of greisenization and concentrations of tourmaline veining are conspicuous close to the margins of the granite but are relatively scarce towards the interior. Quartz feldspar porphyry dykes related to the Meredith Granite occur at Mount Bischoff and at Deep Gully Creek.

5. STRUCTURE

The same stratigraphic and structural elements occur both S and N of the Meredith Granite. South of the granite the structural trends are N-S but they swing to the NE further north. The most obvious structure in the area is the Husskisson Syncline. Ultramafic-mafic rocks were thrust over the Crimson Creek Formation along low angle structures. These rocks are unconformably overlain by Dundas Group sediments, Ordovician limestone and Silurian to Devonian siliciclastics. This sequence is folded about a N-S axis. The same units occur north of the granite in the Whyte River-Bald Hill area.

The blocks of Oonah formation at Mt. Bischoff and in the Coldstream valley have possibly also been thrust over the top of Crimson Creek Formation. Similar situations have recently been recognised on the Sorell Peninsula and near Zeehan, where Precambrian rocks have been thrust over Crimson Creek Formation. The Magnet Dyke is a ductile ultramafic/mafic slice which may have "lubricated" the fault movement. To the east of the granite, there are probably a series of fault slices running parallel to the granite contact. The block of Oonah formation may in fact contain several faulted slices of Precambrian rocks.

The structural control on the granite emplacement is apparent from the shape of the granite, particularly with the steep eastern and western sides, and the shallow plunging NE ridge under Mount Bischoff, following the regional structural trends.

6. WORK COMPLETED

6.1 Geology

Geology base maps were prepared at 1:10,000 scale from enlargements of the Tasmania 1:25,000 topographic series. A literature review was completed to determine what exploration has already been undertaken on the area covered by the E.L.s. Twenty-four kilometres of tracks were cut to provide access for a gravity survey. All of these tracks were cut on the portion of the EL's to be retained.

Reconnaissance mapping was undertaken along roads, tramways, walking tracks and some creeks. The aim of the mapping was to gain some information on the structural styles in the area, to check on the mapping of Brown (1986), to provide a geological basis for the interpretation of the geochemistry and geophysics and to determine what areas, if any, warranted more detailed follow-up work.

6.2 Geochemistry

A total of 99 rock chip samples were collected during reconnaissance mapping, however only 3 of these were from the area being relinquished. These were analysed at Analabs, Cooee for Sn, Cu, Pb, Zn, Ag, As and some for W. The samples were crushed and disc-ground to 80% < 150 microns. A riffle split of 500g was taken and pulverized in a Labtech Mill and a split was then taken for analysis.

The analytical methods employed were:

Method	Element	
401	Sn, W:	XRF
101	Cu, Pb, Zn, Ag:	AAS
114	As:	Hydride Generation/AAS

During the 25 year period that Comstaff held EL 5/63, over 3000 stream sediment samples were collected and analysed in the area covered by the present E.L.'s. Rather than duplicate this effort, it was decided to have a closer look at the existing data and be more selective for areas warranting check or follow-up sampling. A complete compilation of all stream sediment data is presented by Comstaff (1977). This data was entered on the RGC computer system and geochemical maps were plotted using the "Geochem" program. This highlighted several areas of interest where more detailed work could be undertaken. Most of these were in the portions of the EL's to be retained. However the area south east of Mount Ramsay, within 3 to 4 km of the granite contact had high background levels of Pb, Zn, and Cu. Although Comstaff concentrated much of their work in this area, it probably still warrants a detailed follow-up.

Detailed stream sediment sampling to check anomalies from the Comstaff database was carried out but all the anomalies tested were in the areas to be retained.

6.3 Geophysics

The EL's 12/90 and 15/90 lie within an area covered by the Tasmanian Department of Mines' regional aeromagnetic survey of the West Coast. This survey was flown in May 1981 by Geoex Pty. Ltd. with east-west flight lines 500m apart and a mean sensor height of 135m. The Tasmanian Department of Mines presented the data as a 5 nt contour plan at 1:50,000 scale which was obtained by RGC in 1990 and supplied to Geoimage who used image processing techniques to present the data as coloured and grey scale images of residual magnetics and vertical derivative, with various directional filters. This information was sent to Wyatt and Associates to be interpreted.

The existing gravity data base for the Waratah-Ramsay area was acquired as part of the Mount Read Volcanics Project (Leaman and Richardson, 1989). Although this survey defined the gross form

of the Meredith Granite, the station spacing was too coarse to allow identification of exploration targets. Inaccessible parts of the region had no gravity readings at all.

An infill survey was proposed to better define the form of the granite, particularly in those areas where buried ridges of granite occurred at relatively shallow depths. All available vehicular and foot tracks were used and an additional 24 km of walking tracks were cut. Readings were taken at approximately 500m spacings along these lines of access. The survey was undertaken by Mackie Martin Pty. Ltd. In total, 337 gravity station readings were added to the data base. This information was processed and interpreted by David Leaman of Leaman Geophysics.

7. RESULTS

The Comstaff stream sediment sampling highlighted areas of known mineralization. Drainages from Mount Bischoff and Magnet were particularly conspicuous. Creeks from the South Bischoff greisen and from the Wombat Flat and Upper Ramsay alluvial areas were not surprisingly also anomalous in tin.

The rock chip geochemistry is presented in appendix 3. These three samples were collected from the Coldstream Road. Sample T20090 was a limonitic breccia in Oonah quartzite from the crest of the ridge between the Coldstream and Ramsay valleys. It was anomalous in base metals and arsenic. This area was gridded and drill tested by Comstaff and was one of their more interesting prospects. The drilling encountered extensive sequences of dolomite with weak, patchy alteration.

Several areas of known mineralization have distinct magnetic signatures and are evident on the 1981 aeromagnetic survey. These include Mt. Bischoff and particularly the magnetite skarns at the northern end of the Husskisson syncline. A number of the geological units in the area can be distinguished from the magnetics. These include the Meredith Granite, the Tertiary basalt, the inliers of Oonah Formation, the ultramafic complex and particularly the metamorphic aureole in the Crimson Creek Formation surrounding the Meredith Granite. Following an inspection of the data, both Wyatt and Mudge recommended flying a low-level high resolution aeromagnetic survey (Wyatt, appendix 1). However, it was decided not to proceed with the recommendations for the following reasons.

a) The magnetic nature of the basalt makes it difficult to detect anomalies beneath the basalt cover.

b) The Meredith Granite has produced a highly magnetic contact metamorphic aureole in the adjacent Crimson Creek rocks. This effect covers some highly prospective parts of the EL's. It is unlikely that a magnetic anomaly of the magnitude of that produced by the Mount Bischoff mineralization could be distinguished within this highly magnetic contact zone.

c) Given the topography of the area, it would be difficult to maintain a constant survey height with a fixed-wing aircraft. Using a helicopter would add substantially to the cost of a high resolution survey.

It should however be pointed out that some areas of mineralization have magnetic anomalies that can be distinguished from the available data, even given its obvious limitations. A large number of anomalies, although probably related to the country rock, have never been investigated. A more detailed survey as proposed by Wyatt would certainly generate numerous exploration targets.

The gravity survey indicates that the Meredith Granite is largely unroofed (Leaman, appendix 2). The eastern face of the granite is very steep and seems to have little potential for large deposits.

However gravity interpretation has limitations due to the lack of a good structural model and a lack of rock density data. For example it failed to detect the cupola associated with the Cleveland deposit. Similarly the mass of dykes at Mount Bischoff suggests closer proximity to a granite mass than indicated by the gravity interpretation.

If the gravity interpretation was to be constrained by a better structural model and by appropriate rock density measurements, infill gravity data and reinterpretation could greatly assist in defining prospective areas.

8. CONCLUSIONS

Regionally, the margins of the granite and its surrounds are difficult to explore. Magnetic targets are obscured by the very high magnetic background in the contact metamorphic aureole. Although magnetic anomalies abound, most will be due to the magnetic properties of the country rock rather than to mineralized bodies.

The granite stands out as a topographic high and sheds abundant stream sediment anomalies from the small but widespread greisen zones occurring near its margins. Using stream sediment geochemistry to explore for carbonate replacement style tin mineralization is difficult because of the widespread tin in the streams. Since the western Tasmanian tin deposits have broad base metal halos, an association with elevated stream sediment base metal geochemistry may distinguish Bischoff-type mineralization from granite-hosted greisens.

A detailed gravity survey is one way of selecting potentially mineralized areas since most of the granite related mineralization sits above shallowly buried ridges of granite. The gravity survey completed suggests that the southern and eastern margins of the Meredith Granite are steeply dipping and that the main body of the Meredith Granite has been unroofed. It is also apparent that a better knowledge of the regional structures and rock properties is needed to refine the gravity interpretation.

The basalt cover over the eastern parts of the EL's remains a problem. Although it undoubtedly covers some very interesting rocks, drilling to test the sub-basaltic geology is considered to be a very expensive way to explore at this time.

9. REFERENCES

Aberfoyle, 1985: Final report, Exploration Licence 16/78, Meredith, Tasmania. Unpub. rep. Aberfoyle.

BHP, 1988: Exploration Licence 5/63. Final report on the relinquished area: Results of exploration for 1987-1988 and a summary of activities for the period 1963-1988. Unpub. rep. BHP.

Brown, A.V., 1986: Geology of the Dundas - Mt. Lindsay - Mt. Youngbuck region. Bull. geol. Surv. Tasm. 62.

Cartwright, A.J., Komyshan, P., and Roberts, P.A., 1984: E.L. 17/77 Wilson River area: Annual report for 1983/84. Unpub. rep. GFEL.

Hofto, V. and Morrison, K.C., 1989: Exploration Licence 59/88 - Mount Ramsay. Relinquishment report. Unpub. rep. Soloriens Mining Pty. Ltd.

Leaman, D.E., and Richardson, R.G., 1989: The granites of west and north west Tasmania - A geophysical interpretation. Bull. Geol. Surv. Tasm. 66.

Levings, J.A., 1984: Deep Gully Creek interim report, EL 5/63 Area 1. Unpub. rep. Comstaff

Randell, J.P., and Hungerford, N., 1989: E.L. 46/88 - Waratah. Progress report on exploration to 23rd. December 1989. Unpub. rep. Billiton.

Shaw, R.W.L. and Everett, M.P., 1985a: Final report on areas relinquished to the Department of Mines Tasmania (June 1985), Exploration Licence 5/63, Area 1, Arthur River. Unpub. rep. Comstaff.

Shaw, R.W.L. and Everett, M.P., 1985b: Final report on areas relinquished to the Department of Mines Tasmania (June 1985), Exploration Licence 5/63, Area 2, Ramsay. Unpub. rep. Comstaff.

083024

APPENDIX 1

Interpretation of Regional Aeromagnetic data from the Waratah-Mt. Ramsay area, Tasmania

083025

Interpretation of Regional Aeromagnetic data from the Waratah - Mt Ramsay area, Tasmania

E.L.'s 12/90, 15/90
BURNIE SK 55-3

for

RGC Exploration Pty. Limited

Bruce Wyatt

December 1990

Wyatt & Associates
21 Custance St, FARRER
PO Box 705 MAWSON ACT 2607
Phone 06 2863519

083026

Distribution

RGC Exploration, Hobart (2 copies)

RGC Exploration, Canberra

Wyatt & Associates

Summary

The area is part of the Tasmanian Department of Mines' 500 metre line spaced aeromagnetic survey of the west coast of Tasmania. The data were acquired using a fixed wing aircraft with a proton precession magnetometer. Navigation was achieved using aerial photography. The mean sensor height was 135 metres above ground level, but varied between 80 and 500 metres.

The regional aeromagnetic data clearly demonstrate the differing magnetic characteristics of the major rock types in the area. The Meredith Granite is non-magnetic but gives rise to a metamorphic halo with characteristics determined by the nature of the country rocks. Small isolated anomalies within the granite may indicate skarns. The Tertiary basalts form complex anomalous zones which mask the weaker anomalies from underlying Palaeozoic rocks. However, some of the stronger anomalies due to Palaeozoic sources can be delineated within the complex zones.

Strong anomalies are associated with volcanic units within the Crimson Creek Formation and with ultramafic units. The potential for recognising structures controlling mineralisation and smaller anomalies due to possible mineralisation is dependent on acquiring more detailed data which is more amenable to filtering and detailed modelling techniques.

A review of the previous use of other geophysical techniques in the area is considered necessary: to determine the response of the known ore bodies to various methods; to delineate areas which have been adequately tested; and then to define the areas with the most potential for exploration and the most promising techniques to use.

The discrete magnetic anomalies within and adjacent to the margins of the Meredith Granite should be investigated as primary targets for skarn and pyrrhotite bodies. High resolution ground magnetics (and possibly other techniques) with detailed computer modelling is necessary to define targets for testing by drilling. Previous detailed geophysics, geology, and drilling should be taken into account before further work is undertaken on any magnetic anomalies. A review on the application of various geophysical techniques to the area is recommended.

The survey data have limitations which can be attributed to the data acquisition phase. The area should be reflown with high resolution equipment and 100 to 150 metre line spacing. The steep topography of about half of the area warrants the consideration of the use of helicopter rather than fixed wing aircraft. Consideration should also be given to acquiring EM as well as magnetic data.

A detailed survey using state-of-the-art technology could be expected to provide more detailed information regarding: the definition of smaller amplitude targets within generally non-magnetic rock units; the resolution of any small anomalies adjacent to large amplitude anomalies over ultramafics; the resolution of units within the Crimson Creek Formation; the definition of faults and other structures; the thickness of Tertiary basalt; the location of any windows of relatively thin basalt cover; and the structure and composition of the basement beneath the basalts.

Table of Contents

Summary

Table of Contents

List of Tables

List of Figures

List of Plates

1 Introduction 1
1.1 Topography 1

2 Geology 2

3 Magnetic interpretation 4
3.1 Data processing and presentation 4
3.2 Magnetic characteristics of mapped units 4
3.3 Lineaments and structure 6
3.4 Modelling 7

4 Conclusions 21

5 Recommendations 23

6 References 24

List of Tables

1 Stratigraphic table (after Brown 1984) 2
2 Modelled body characteristics - tabular prisms 9
3 Modelled body characteristics - ellipsoids 9
4 Modelled body characteristics - 3-dimensional prisms 10

List of Figures

1 Magnetic profile, line 2320, 5415000 mN 11
2 Magnetic profile, line 2270, 5412500 mN 12
3 Magnetic profile, line 2220, 5409800 mN 13
4 Magnetic profile, line 2170, 5407500 mN 14
5 Magnetic profile, line 2140, 5406000 mN 15
6 Magnetic profile, line 2071, 5402500 mN 16
7 Magnetic profile, line 1990, 5398500 mN 17
8 Magnetic profile, line 1970, 5397500 mN 18
9 Magnetic profile, line 1880, 5393000 mN 19
10 Magnetic profile, line 1837, 5390500 mN 20

List of Plates

Note: All images (Plates 1 to 12) are at 1:100,000 scale

1 Residual magnetics image - greyscale

2 Residual magnetics image - coloured

3 Magnetics image - vertical derivative greyscale

4 Magnetics image - vertical derivative coloured

5 Magnetics image - vertical derivative with AGC greyscale

6 Magnetics image - minimum gradient greyscale

7 Magnetics image - HSI transform using VD as intensity

8 Magnetics image - HSI transform using VD with AGC as intensity

9 Magnetics image - vertical derivative with AGC & 0° azimuth shade

10 Magnetics image - vertical derivative with AGC & 45° azimuth shade

11 Magnetics image - vertical derivative with AGC & 90° azimuth shade

12 Magnetics image - vertical derivative with AGC & 135° azimuth shade

13 Stacked profiles - second horizontal difference bipole 1:100,000 scale

14 Location of modelled bodies and flight lines 1:100,000 scale

15 Residual magnetic contours, 5 nT interval 1:50,000 scale

16 Magnetic interpretation 1:50,000 scale

Note: The following maps have also been generated but are *not* included with this report.

Residual magnetic intensity contours (5 nT interval) 1:100,000 scale

Laminated magnetics image - vertical derivative with AGC greyscale 1:100,000 scale

Flight path 1:100,000 scale

Laminated stacked profile plots at 1:100,000 scale, second horizontal difference bipole

1 Introduction

The area is part of the Tasmanian Department of Mines' regional aeromagnetic survey of the west coast of Tasmania. This survey was flown in May of 1981 by Geoex Pty. Limited, with east west flight lines nominally 500 metres apart and a mean sensor height of 135 metres above ground level. The data were acquired using a Cessna A185E fixed wing aircraft with a Sonotek magnetometer. The resolution was 0.1 nanoTesla (nT), accuracy 0.5 nT, and sample interval 0.8 second (40 metres). Navigation was achieved using aerial photography and flight path recovery using 16 mm ground tracking camera.

This interpretation is part of an exploration effort which aims to discover Renison-style pyrrhotite-rich replacement tin mineralisation. The aim is to locate major structures which act as conduits for the mineralising solutions that are known to have emanated from the Meredith Granite. Of interest is where these structures intersect stratigraphic carbonate horizons that act as hosts for the tin mineralisation. Deposits of this type in the area are the Cleveland Tin deposit at Luina and the Mt Bischoff deposit at Waratah. Carbonate horizons would be expected in the Precambrian Oonah Formation, the Cambrian Crimson Creek Formation, and the Ordovician Gordon Limestone. Some Cambrian carbonate alteration may occur adjacent to the ultramafics. Magnetite-bearing skarns are also known to occur in the area.

1.1 Topography

The EL's comprise about equal areas of plateau (between about 560 and 660 metres above height datum) and fairly rugged dissected terrain down to 150 metres metres above height datum.

Mt Ramsay (855 metres), Wombat Hill (850 metres) and Mt Pearse (1001 metres, just off the eastern edge of EL 15/90) rise above the plateau. The dissected areas include parts of Deep Gully Creek, Waratah River, Magnet Creek, Ritchie Creek, and Arthur River around Mt Bischoff (784 metres) in the north of EL 12/90; and the Ramsay, Coldstream, Hatfield, Wilson, Que, and Huskisson Rivers and their tributaries in the south of EL 15/90.

The plateau area comprises mainly outcrop of Meredith Granite (in the west, and an extensive area of Tertiary basalt.

2 Geology

Table 1 Stratigraphic Table - (after Brown, 1984)

Permian-Carboniferous		
	Pt	tillite
-------- landscape unconformity ----------		
Silurian-Devonian		
Eldon Group	S-Du	
---------- disconformity ----------		
Ordovician		
Gordon Limestone	Ogl	limestone
Owen Formation	Os	conglomerate, sandstone, siltstone
---------- disconformity ----------		
Cambrian		
Dundas Group	Ed	
Huskisson Group	Eh	
Rosebery Group	Er	
---------- erosional break ----------		
eoCambrian		
Crimson Creek Form.	Ecc	with lava flows (Eccb)
Success Creek Group	Esc	
--------- landscape unconformity ----------		
Precambrian		
Oonah Formation	Po	

<u>Igneous Rocks</u>

Tertiary	Tb	basalt
Jurassic	Jdl	dolerite (intrusive sills)
Devonian	Dg	granitoids
Cambrian		
Mt Read Volcanics	Emrv	pyroclastic and epiclastic acid to intermediate volcanics
	Ebm	low-titanium tholeiite
	Eba	high-magnesian andesite and associated coarse-grained pyroxenite
EoCambrian		
	Es	serpentinised ultramafic-mafic rocks

The outcrop limits of both Tertiary basalt and Meredith Granite, as mapped by Brown (1984) are shown on Plate 16.

Burrett and Martin (1989) should be consulted for general description of the geology and mineral deposits of the area, as well as a comprehensive list of references.

Leaman and Richardson (1989) have analysed regional gravity data over the Meredith Granite using 2-dimensional modelling along five profiles which cross the pluton. Their provisional interpretation indicates that the granite pluton has steep edges on the west and much of the northwest and southeast, but shelves to the northeast (beneath Cleveland, Magnet, and Bischoff mines). A northeast trending zone in the southeast of EL 15/90 has also been indicated as a zone with "complex alternative structures possible".

3 Magnetic Interpretation

3.1 Data Processing and Presentation

The original data were gridded (125 * 125 metre cells) and 5 nT contours were produced at 1:50,000 scale. Unfortunately, the use of a fixed wing aircraft over the rugged terrain produced large variations in ground clearance, particularly between adjacent lines flown on reciprocal headings. This resulted in marked "herring-bone" patterns on the quieter parts of the contour maps and ruled out the possibility of using a finer contour interval to display more detail.

More recently, RGC Exploration has used several other techniques to enhance the processing and presentation of the data.

In 1990, Geoimage Pty Limited used image processing techniques and presented the data as greyscale and coloured images of: residual magnetics; vertical derivative (VD); and VD with automatic gain control (AGC). Directional filters were also applied to the VD with AGC data sets (Walker, 1990). The locations of Mt Bischoff and Cleveland tin mines are shown on the images as 'MB' and 'CL' respectively.

In 1990, Surtec Pty Limited reprocessed the data and produced flight path map and second horizontal difference bipole stacked profiles at 1:50,000 scale.

The vertical derivative and horizontal difference operators are high pass filters and both effectively reduce the effect of the levelling problems in the data and allow the delineation of lower amplitude, short wavelength anomalies.

The magnetic data are shown as Plates 1 to 14. Additional displays of images exist as 35 mm photographic slides. The contour map is most useful for accurate positioning of all anomalies and other interpreted features. Profiles generated for modelling from the digital data provide the best display of the relative amplitudes of anomalies and anomaly shape for matching computer generated models. The other images have been filtered to enhance various trends and spectral characteristics, particularly short wavelength features which may be related to faulting.

3.2 Magnetic Characteristics of Mapped Units

The Eldon Group coincides with regions of extensive magnetic gradients in the far west and the southwest of the area. In the far west, the gradient probably reflects the geometry of underlying Cambrian rocks which outcrop further to the southwest. In the southwest, the gradient is due to Cambrian outcrop to the east and west. There are several smaller anomalies within the outcrop in the southwest. Most of these are adjacent to the margin of the Meredith Granite and are recommended for further investigation.

Gordon Limestone crops out in the west of the area, between Eldon Group and Cambrian rocks. The limestone itself is completely non-magnetic and is coincident with gradients due to the Cambrian rocks. A couple of anomalies are adjacent to the margin of the Meredith Granite and are recommended for further investigation.

The Owen Formation crops out at Mt Pearse in the far east of the area. It appears on the magnetic contours as a relative low, due to the window in the surrounding

(magnetic) Tertiary basalt.

Rocks of the Dundas Group and its correlates crop out in a north-northeast trending belt in the southeast of the area. The more northwesterly, undifferentiated part is magnetically quiet with only a couple of fairly small anomalies, up to a few tens of nT in amplitude, which occur on a line subparallel to strike and topography.

The Huskisson Group (a correlate of the Dundas Group) contains a zone with northeast trending linear anomalies up to 150 nT in amplitude. This zone presumably reflects the presence of volcanic units within the Huskisson Group, or possibly within underlying Mt Read Volcanics. An anomaly due to a reversely magnetised source occurs at 5397000 mN, 383000 mE (just outside EL 15/90). This is similar to some anomalies over the Tertiary basalts and possibly indicates an unmapped basalt outlier. If this is not the case, it should be further investigated to determine the source.

The Crimson Creek Formation crops out in a north-south zone through the centre of the area and in an extensive area including the northwest part of the area. The Formation is generally quite magnetic, especially those areas mapped as dominantly basalt flows. The magnetic data indicate that the basalts are more extensive than indicated by mapping. The higher amplitude anomalies also indicate where the Formation extends beneath the westernmost area of Tertiary basalt cover. The acquisition of more detailed aeromagnetic data over the Crimson Creek Formation would allow much better resolution of magnetic units within the Formation and the direct detection of any anomalies due to pyrrhotite associated with mineralisation.

The Oonah Formation and its correlates crop out in the south of the area, between outcrop of Dundas Group and Crimson Creek Formation. The only magnetic anomalies (in the northeast) have amplitudes of a couple of hundred nT and indicate that the Tertiary basalts probably extend further than mapped. Ground checking is required to verify this assumption.

Oonah Formation also occurs around Mt Bischoff. The only detected magnetic anomaly is the 300 nT anomaly over Mt Bischoff.

Tertiary basalt covers an extensive area in the northeast, and also occurs as other smaller outliers. The magnetic field over these outcrops is characterised by complex, short wavelength anomalies with amplitudes of a few hundred nT. Some of the anomalies are negative, indicating reverse magnetisation in some flows.

While it is possible to delineate the anomaly due to a major magnetic unit within the Crimson Creek Formation where it lies beneath the basalt because of its high amplitude (about 1000 nT), the direct detection of magnetic anomalies associated with mineralisation beneath the basalt cover is not very likely.

The basalt surface is relatively flat, so the main factor determining the anomaly shape will be the depth of weathering and the configuration of the lower surface of the flows, the pre-basalt topography. Better quality magnetic data could be used to determine the thickness of basalt and in particular, any windows of relatively thin basalt cover.

The Meredith Granite has an extensive outcrop area in the west. The granite is not magnetic. There are quite a few anomalies adjacent to the margins of the pluton which are recommended for further investigation. A couple of isolated anomalies

within the granite are probably due to skarns or unmapped outliers of Tertiary basalt. These are also recommended for further investigation.

Porphyry dykes at Mt Bischoff are also associated with the Meredith Granite. An unusual shaped anomaly over Mt Bischoff is probably partly due to equipment and buildings associated with the mine, but modelling studies indicate that the bulk of the anomaly is due to a quite extensive source with significant depth extent (see Figure 2).

Cambrian gabbros crop out northwest of the EL's and coincide with part of some significant magnetic anomalies. The magnetic data suggest that the gabbros may be much more extensive in the subsurface than mapped on the surface.

The Mt Read Volcanics crop out extensively to the southeast of the EL's and contain some magnetic anomalies with amplitudes up to several tens of nT.

Cambrian massive ultramafic cumulates occur to the north of the EL's (north of Mt Bischoff). These outcrops appear to be magnetic but their effects are not easily distinguished from the adjacent Tertiary basalt.

Other Cambrian and eoCambrian rocks occur near the northwest boundary of EL 12/90. The magnetic response of these is variable. The low-titanium tholeiite is generally non magnetic to the west of Mt Bischoff, but strongly magnetic west of Luina (around Heazlewood). High-magnesian andesite and associated coarse-grained pyroxenite shows a complex pattern in the west of EL 12/90, with anomalies of a few tens of nT.

The derivative maps (Plates 5 and 13) show different character over some of the rock units, generally due to the effect of the automatic gain control (AGC) filter amplifying noise over essentially non-magnetic rocks. The most marked zones occur over the Meredith Granite, Eldon Group, Dundas Group, and part of the Oonah Formation (outside the EL's, 12 km north-northwest from Luina).

Plate 16 shows the position of magnetic sources and contacts as interpreted from the magnetic contour map and the second horizontal difference profiles. The delineated areas correspond to the surface projections of the sources of the magnetic anomalies. The second horizontal difference profiles (Plate 13) have an inherent bias against east-west features because of the use of a high pass filter along the lines only, and the stacked profile display. However, given that very few of the geological features trend east-west and that the contours provide additional information, the bias does not detract significantly from the interpretation. The areas of highest second horizontal difference correspond well with the interpretation of source positions from the contours. Some of the areas delineated in Plate 16 show the extra detail that can be extracted by the use of appropriate filtering and presentation techniques.

3.3 Lineaments and Structure

Several magnetic lineaments have been interpreted from a study of all available data displays, particularly the images of vertical derivative with automatic gain and with shading applied from various directions (Plates 9 to 12). These lineaments have generally been defined by steeper gradients or by colinear anomalies or truncations of anomalies. The lineaments are shown on Plate 16 and are generally considered to be indications of faults or, in some cases, of units within the

stratigraphy. In the northeast of the area, where the magnetic field is dominated by the effect of Tertiary basalts, only the strongest lineaments have been delineated.

A few strong anomalies trend northeast in a line from Mt Ramsay to just north of Mt Pearse. These are parallel to linear anomalies over the Huskisson Group and may also reflect units within the Palaeozoic basement. An alternative explanation is that they reflect greater thickness of basalt within older river valleys.

The majority of the interpreted lineaments trend northwest to north-northwest, with most of the remainder trending north-northeast. In the western part of the Meredith Granite (outside the EL's) two lineaments extend through the granite at 350° and 355°. The most dominant lineaments are one which passes through Mt Bischoff and another which truncates most of the Meredith Granite.

The large north-south anomalies over Crimson Creek Formation are interpreted to indicate dyke-like bodies which dip away from the Meredith Granite at its margin (Figures 4 to 7). More detailed analysis of these anomalies, along with gravity anomalies, may allow definition of the subsurface shape of the granite.

The delineation of structure would be much easier and more reliable with a more detailed data set.

3.4 Modelling

Most of the more prominent anomalies have been computer modelled to provide a better appreciation of the shape and nature of their sources. It is stressed that the models are not unique, and that various simplifying assumptions have been made in terms of the geometries of the bodies employed in the models. More detailed airborne or ground magnetic readings and modelling studies would be required before drill targets could be defined.

Remanent magnetisation has not been taken into account in the interpretations, except in the case of some Tertiary basalt flows which are reversely magnetised (see Figures 7 and 8).

The characteristics of all bodies used in the modelled profiles are listed in Tables 2, 3, and 4. Units on all profile plots are kilometres and nanoTeslas, with arbitrary origins. Remanence has not been included in any of the models because of lack of adequate control. The earth's main field has been taken as 62535 nT amplitude, inclination -71.8°, and declination 12.8°.

Data from ten flight lines have been quantitatively interpreted using a 3-dimensional modelling package (Almond, 1990). These lines were chosen to demonstrate the effects of most interpreted magnetic sources in the survey area.

Data on these lines were reprocessed by digitising ground elevation and adding this to the recorded aircraft ground clearance. The modelled anomalies can then be computed at their true locations in space. The ground elevation lies between 150 and 1000 metres above sea level. The nominal sensor height was 135 metres above ground level, but actually varied from less than 75 metres to more than 500 metres above ground level. Differences in aircraft elevation on reciprocal headings are responsible for some "herring-boning" on the contour presentations.

The results of the profile modelling studies are presented as Plate 14 and Figures 1 to 10. Each of the figures comprises a profile showing the observed and computed total magnetic fields and a cross section showing all bodies used for the model on that profile. The modelled profiles are oriented 'looking North'. The observations are shown as '+' symbols, the height of the observations as the lower solid line, and the computed magnetic anomaly field as the upper solid line. The scales of the ten figures are all identical to give a better appreciation of anomaly amplitudes and body dimensions.

The initial positions of the bodies and their strike and strike length were determined by examination of the contour map (Plate 15) and the second horizontal difference map (Plate 13). Plate 14 is a plan view showing the surface projections of the limits of each body, and the location of each ninth data point along the ten modelled lines. The observation points are shown as '+' symbols.

The calculated field for each profile generally only includes the effects of those bodies which are intersected by the profile, with the following exceptions. Body 1 was not used for line 2320. Body 15 and 18 were not used for lines 2170 or 2140. Body 23 was not used in lines 2170 or 2220. Body 26 was not used for line 2220. Body 32 was not used for line 1990. Bodies 47 and 48 were used for line 1880. Body 42 was used for line 1837.

In each profile, a (separate and arbitrary) linear regional has been removed from the observed field. The magnetic field in the area shows a strong east-west gradient, with the measured field being about 200 nT higher in the east than in the west. This may reflect the presence and thickening of Tertiary basalt to the east or may be due to another deeper source.

Tables 2, 3, and 4 list the parameters used to define each of the bodies used for all models. Xr, Yr, and Depth refer to the "reference point" of each body. For the ellipsoid bodies, the reference point is the top of the body. For the tabular prisms, the reference point is at the centre of the upper surface of the body. For the 3-D bodies, the cross-section is defined by several pairs of coordinates whose location is given relative to the reference point (corner 1) which is taken to be the local origin. All distances are kilometres. Susceptibility is in SI units. The susceptibilities assigned to the bodies is generally about 0.01 for the Tertiary basalt sheets and between 0.03 and 0.05 for the ultramafic bodies.

It should be noted that the interpretation of any particular body is not unique. In particular, an approximately inverse relationship exists between the susceptibility and size of the body.

Little reliance should be placed on those parts of the modelled bodies at depths of greater than a few hundred metres below the aircraft.

It should be noted that body 35 is only half the width on line 1970 that it is on line 1990. Body 9 corresponds to Mt Bischoff.

Body 38 has been assigned a negative susceptibility to denote reversed direction of magnetisation.

Attempts have been made in a few cases to model the general effects of the basalt, but these should be considered as demonstrations of some of the possible effects of edges and windows of sheets rather than a rigorous attempt to match the observed field. Bodies 4, 5, 10, 11, 12, 27, 28, 29, and 38 on Figures 1, 2, 5, 6, 7, and 8 were

designed to show the shapes and amplitudes of anomalies over basalt sheets of varying shape, attitude and thickness.

Table 2 Modelled Body Characteristics - Tabular Prisms

No	Xr (east)	Yr (north)	depth	strike (deg)	dip (deg)	susc (SI)	width	breadth	height
1	374.011	5414.500	-0.250	30	115	0.017	0.60	5.0	1.0
2	377.188	5415.537	-0.543	315	90	0.010	2.30	3.0	0.8
3	377.829	5414.765	-0.543	340	90	0.030	0.60	1.3	0.6
4	380.490	5415.000	-0.633	20	90	0.010	1.10	3.5	0.25
5	385.500	5417.972	-0.678	0	90	0.015	8.00	9.0	0.6
6	370.306	5411.265	-0.498	30	110	0.010	0.30	2.8	0.7
7	371.432	5412.313	-0.543	70	90	0.010	0.50	3.0	0.7
8	372.328	5413.009	-0.543	320	90	0.010	0.50	1.4	0.6
13	367.800	5409.832	-0.500	30	120	0.015	0.30	1.5	2.0
14	368.847	5409.988	-0.707	10	110	0.015	0.30	1.0	2.0
15	369.481	5407.972	-0.500	5	130	0.035	0.50	3.7	2.0
16	370.589	5409.988	-0.550	95	120	0.035	0.70	1.3	2.0
17	371.640	5409.212	-0.424	95	120	0.035	0.70	3.7	2.0
18	373.381	5408.050	0.200	355	35	0.040	0.50	4.0	4.0
19	374.193	5410.220	-0.509	340	90	0.020	0.50	2.0	2.0
20	374.915	5408.980	-0.537	0	90	0.020	0.30	3.0	2.0
21	376.187	5409.987	-0.594	5	90	0.015	1.00	2.0	2.0
38	379.867	5397.830	-0.500	35	90	-0.020	1.20	0.9	0.2
43	375.480	5390.790	-0.100	35	60	0.010	0.60	5.0	1.5
44	380.001	5393.073	-0.400	37	90	0.010	0.15	10.0	2.0
45	380.001	5392.373	-0.400	37	90	0.010	0.15	10.0	2.0
46	380.001	5391.673	-0.400	37	90	0.010	0.15	10.0	2.0

All distances are in kilometres.
Depths are positive down, relative to mean sea level (negative depths are above sea level).

Table 3 Modelled Body Characteristics - Ellipsoids

No	Xr (east)	Yr (north)	depth	strike (deg)	dip (deg)	susc (SI)	a	b	c	plunge (deg)
33	368.293	5397.094	-0.464	15	105	0.017	0.75	0.40	0.10	0
34	369.446	5397.113	-0.520	0	80	0.060	0.15	0.10	0.50	0
39	366.151	5392.768	-0.170	-90	90	0.027	0.30	0.45	0.40	-20
40	367.175	5392.769	-0.368	-90	90	0.020	0.15	0.30	0.20	-20
41	369.243	5392.535	-0.204	-90	90	0.030	0.60	0.80	0.70	0

a, b, c are the ellipsoid axis lengths.
All distances are in kilometres.
Depths are positive down, relative to mean sea level (negative depths are above sea level).

Table 4 Modelled Body Characteristics - 3-Dimensional prisms

No	Xr (east)	Yr (north)	depth	strike (deg)	susc (SI)	length	coordinates of corners (X on first line, Y on second line)							
9	376.484	5412.201	-0.633	0	0.010	1.8	0.0	1.928	3.085	-2.769	-0.280			
							0.0	0.724	4.433	4.478	0.045			
10	379.850	5413.001	-0.678	0	0.010	3.0	0.0	1.648	1.332	-1.613	-1.683			
							0.0	0.090	0.407	0.452	0.090			
11	382.233	5412.989	-0.633	0	0.010	3.0	0.0	0.280	0.070	-0.245	-0.561			
							0.0	0.000	0.317	0.317	0.000			
12	385.187	5412.975	-0.701	0	0.010	3.0	0.0	2.830	2.865	0.516	-2.007	-2.148		
							0.0	0.023	0.384	0.701	0.430	0.023		
22	362.852	5406.211	-0.396	0	0.010	5.0	0.0	0.351	0.701	0.942	-0.329			
							0.0	-0.042	0.141	1.371	1.173			
23	365.556	5406.935	-0.400	30	0.030	7.0	0.0	0.439	1.318	1.784	1.942	-0.237		
							0.0	-0.194	-0.307	-0.240	1.927	1.927		
24	368.285	5406.140	-0.650	10	0.010	5.0	0.0	0.351	0.500	0.419	-0.100			
							0.0	-0.042	0.141	1.173	1.173			
25	369.738	5406.077	-0.650	30	0.035	5.5	0.0	0.803	0.992	-0.274	-1.230			
							0.0	-0.113	0.000	5.456	5.400			
26	373.024	5406.778	-0.537	345	0.035	5.0	0.0	0.306	0.786	1.335	3.906	2.557		
							0.0	-0.254	-0.170	0.057	6.587	6.530		
27	382.224	5405.616	-0.622	30	0.010	3.8	0.0	1.059	0.819	-0.251				
							0.0	0.000	0.452	0.396				
28	384.984	5405.923	-0.622	10	0.020	2.5	0.0	0.694	1.196	0.158				
							0.0	0.000	0.452	0.452				
29	386.210	5406.235	-0.707	0	0.020	5.0	0.0	1.000	0.904	0.028				
							0.0	0.000	0.396	0.396				
30	368.329	5402.507	-0.664	355	0.050	2.5	0.0	0.219	0.382	-0.317				
							0.0	-0.014	0.452	0.481				
31	371.865	5402.497	-0.664	0	0.010	4.0	0.0	21.141	21.229	2.180	1.818	1.150	0.712	-0.055
							0.0	-0.071	0.269	0.325	0.212	0.184	0.325	0.325
32	372.564	5402.111	-0.339	20	0.040	5.5	0.0	0.409	1.033	1.371	4.000	2.650		
							0.0	-0.141	-0.113	0.000	6.757	6.700		
35	373.001	5397.466	-0.548	0	0.045	4.5	0.0	0.300	-0.100	-0.400	4.500			
							0.0	3.000	3.000	0.000				
36	373.730	5397.499	-0.448	0	0.015	5.0	0.0	20.001	20.002	-0.301				
							0.0	0.003	0.601	0.598				
37	375.650	5397.849	-0.400	340	0.030	5.0	0.0	0.300	-0.150	-0.450				
							0.0	1.500	1.500	0.000				
42	372.201	5394.001	-0.430	355	0.042	4.0	0.0	0.500	0.600	-0.500	-0.500	-0.230		
							0.0	0.270	6.000	6.000	0.210	0.000		
47	371.850	5389.998	0.250	355	0.040	5.0	0.0	0.400	-0.100	-1.050	-0.600	-0.200		
							0.0	0.283	5.000	5.000	0.300	0.000		
48	372.850	5390.100	0.226	355	0.040	4.0	0.0	0.150	-0.350	-0.950	-0.450	-0.262		
							0.0	0.283	5.000	5.000	0.226	0.007		

All distances are in kilometres.
Depths are positive down, relative to mean sea level (negative depths are above sea level).




Figure 1 Magnetic profile, line 2320, 5415000 mN




Figure 2 Magnetic profile, line 2270, 5412500 mN




Figure 3 Magnetic profile, line 2220, 5409800 mN




Figure 4 Magnetic profile, line 2170, 5407500 mN




Figure 5 Magnetic profile, line 2140, 5406000 mN




Figure 6 Magnetic profile, line 2071, 5402500 mN




Figure 7 Magnetic profile, line 1990, 5398500 mN




Figure 8 Magnetic profile, line 1970, 5397500 mN




Figure 9 Magnetic profile, line 1880, 5393000 mN




Figure 10 Magnetic profile, line 1837, 5390500 mN

4 Conclusions

The regional aeromagnetic data clearly demonstrates the differing magnetic characteristics of the major rock types in the area. The granite is non-magnetic but gives rise to a metamorphic halo with characteristics determined by the nature of the country rocks. Small isolated anomalies within the granite may indicate skarns. The Tertiary basalts form complex anomalous zones which mask the weaker anomalies from underlying Palaeozoic rocks. However, some of the stronger anomalies due to Palaeozoic sources can be delineated within the complex zones.

Strong anomalies are associated with volcanic units within the Crimson Creek Formation and with ultramafic units. The potential for recognising structures controlling mineralisation and smaller anomalies due to possible mineralisation is dependent on acquiring more detailed data which is more amenable to filtering and detailed modelling techniques.

The acquisition of high resolution ground magnetic data (and possibly data using other techniques) with detailed computer modelling is necessary to define targets for testing by drilling. Any ground magnetic surveys should be designed with lines perpendicular to strike (if practicable) and extending beyond the anomaly far enough to allow accurate definition of the background field, and with sufficient lines to define the strike and anomaly maximum. It is recommended that ground magnetic surveys be conducted along lines spaced 50 metres apart with readings spaced no more than 5 metres apart.

The survey data have limitations which can be attributed to the data acquisition phase. In steep terrain, it is impossible to maintain comparable elevation while flying adjacent flight lines on reciprocal headings with a fixed wing aircraft. The sensor height actually varied from less than 75 metres to greater than 500 metres above ground level. The data were acquired on lines 500 metres apart, with sample spacing about 40 metres and accuracy 0.5 nT.

This interpretation has not included a review of the previous use of other geophysical techniques in the area. Such a review is considered necessary: to determine the response of the known ore bodies to various methods; to delineate areas which have been adequately tested; and then to define the areas with the most potential for exploration and the most promising techniques to use.

In particular, it is suggested that detailed airborne magnetic surveying would assist greatly in definition of pyrrhotite, skarns, or mafic units which may be associated with tin mineralisation. A much more detailed interpretation of the structure would be possible with higher resolution data.

The use of a helicopter as survey platform allows the variations in ground clearance to be very significantly less than with a fixed wing aircraft. In this area, a flight line spacing of 100 to 150 metres is considered to be optimum, with recording of both radar and barometric altitude mandatory. Current alkali vapour magnetometers are capable of providing data with sample interval 5 metres, and resolution 0.01 nT.

The usefulness of detailed airborne electromagnetic surveying over prospective rock units should be evaluated in terms of discovering additional ore bodies or any of the other styles of mineralisation which are present within the area.

The upper surface of the Tertiary basalt is relatively flat, so the main factor determining the anomaly shape will be the depth of weathering and the

configuration of the lower surface of the flows, the pre-basalt topography. Better quality magnetic data could be used to estimate the thickness of basalt and in particular, to determine the location of any windows of relatively thin basalt cover. It is also possible that additional information regarding the structure of the pre-basalt basement rocks could be obtained using appropriate filtering techniques on high quality data.

5 Recommendations

1 The discrete magnetic anomalies within and adjacent to the margins of the Meredith Granite should be investigated as primary targets for skarn and pyrrhotite bodies.

2 Previous detailed geophysics, geology, and drilling should be taken into account before further work is undertaken on any particular magnetic anomalies. A review of the application of various geophysical techniques to the area is recommended.

3 The area should be reflown with high resolution equipment and 100 to 150 metre line spacing. The steep topography of about half of the area may warrant the use of helicopter rather than fixed wing aircraft. Consideration should also be given to acquiring EM as well as magnetic data.

4 High resolution ground magnetics (and possibly other techniques) with detailed computer modelling is necessary to define targets for testing by drilling.

6 References

Brown, A.V., 1984, Geological compilation of the Zeehan-Waratah area, Dundas Trough. 1:100000 scale. Geological Survey of Tasmania.

Burrett, C.F. and Martin, E.L. (editors), 1989, Geology and Mineral Resources of Tasmania. Special Publication Geological Society of Australia 15.

Leaman, D.E., and Richardson, R.G., 1989, The granites of north-west Tasmania - a geophysical interpretation. Bulletin geological Survey of Tasmania 66.

Walker, R.N., 1990, Processing of Magnetics for the West-Coast Survey, Tasmanian Department of Mines. Unpublished report by GeoImage Pty Limited for RGC Exploration Pty Limited.

360 000 mE

370 000 mE

380 000 mE

5 420 000 mN

5 410 000 mN

5 400 000 mN

5 390 000 mN

Plate 1 Greyscale magnetics image, 1:100,000 scale
Waratah - Mt Ramsay area EL 12/90 & EL 15/90

5 cm

083055




Plate 2 Coloured magnetics image, 1:100,000 scale
Waratah - Mt Ramsay area EL 12/90 & EL 15/90

083056

360 000 mE

370 000 mE

380 000 mE

5 420 000 mN

5 410 000 mN

5 400 000 mN

5 390 000 mN

Plate 3 Magnetics image, 1:100,000 scale
vertical derivative greyscale
Waratah - Mt Ramsay area EL 12/90 & EL 15/90

5 cm

083057

360 000 mE

370 000 mE

380 000 mE

5 420 000 mN

5 410 000 mN

5 400 000 mN

5 390 000 mN

Plate 4 Magnetics image, 1:100,000 scale
vertical derivative coloured
Waratah - Mt Ramsay area EL 12/90 & EL 15/90

5 cm

083058

360 000 mE

370 000 mE

380 000 mE

5 420 000 mN

5 410 000 mN

5 400 000 mN

5 390 000 mN

Plate 5 Magnetics image, 1:100,000 scale
vertical derivative with AGC greyscale
Waratah - Mt Ramsay area EL 12/90 & EL 15/90

5 cm

083059




Plate 6 Magnetics image, 1:100,000 scale
minimum gradient greyscale
Waratah - Mt Ramsay area EL 12/90 & EL 15/90

083060.

360 000 mE

370 000 mE

380 000 mE

5 420 000 mN

5 410 000 mN

5 400 000 mN

5 390 000 mN

Plate 7 Magnetics image, 1:100,000 scale
HSI transform using VD as intensity
Waratah - Mt Ramsay area EL 12/90 & EL 15/90

5 cm

083061

360 000 mE

370 000 mE

380 000 mE

5 420 000 mN

5 410 000 mN

5 400 000 mN

5 390 000 mN

Plate 8 Magnetics image, 1:100,000 scale
HSI transform using VD with AGC as intensity
Waratah - Mt Ramsay area EL 12/90 & EL 15/90

5 cm

083062




Plate 9 Magnetics image, 1:100,000 scale
vertical derivative with AGC & 0° azimuth shade
Waratah - Mt Ramsay area EL 12/90 & EL 15/90

083063




Plate 10 Magnetics image, 1:100,000 scale
vertical derivative with AGC & 45° azimuth shade
Waratah - Mt Ramsay area EL 12/90 & EL 15/90

083064

360 000 mE

370 000 mE

380 000 mE

5 420 000 mN

5 410 000 mN

5 400 000 mN

5 390 000 mN

Plate 11 Magnetics image, 1:100,000 scale
vertical derivative with AGC & 90° azimuth shade
Waratah - Mt Ramsay area EL 12/90 & EL 15/90

5 cm

083065

360 000 mE

370 000 mE

380 000 mE

5 420 000 mN

5 410 000 mN

5 400 000 mN

5 390 000 mN

Plate 12 Magnetics image, 1:100,000 scale
vertical derivative with AGC & 135° azimuth shade
Waratah - Mt Ramsay area EL 12/90 & EL 15/90

5 cm

083066

360 000 mE

370 000 mE

380 000 mE

5 420 000 mN

5 410 000 mN

5 400000 mN

5 390 000 mN

Plate 13 Stacked profiles, 1:100,000 scale
second horizontal difference bipole
Waratah - Mt Ramsay area EL 12/90 & EL 15/90


5 cm

083067

92-3357.

083068

Plate 14 Location of modelled bodies and flight lines 1:100,000 scale



360000mE 370000mE 380000mE 390000mE

5420000mN 5410000mN 5400000mN 5390000mN 5380000mN

E.L. 15/90

E.L. 12/90

E.L. 12/90

E.L. 15/90

E.L. 12/90

E.L. 15/90

MOUNT BISHOFF

R.L. 8807

WARATAH

MT CLEVELAND

LUINA

MT PEARCE

MT RAMSAY

MT CHARLOTTE

MT BLOOMFIELD

MT KERSHAW

BURNS PEAK

PARSONS HOOD

LINDSAY

Webstertite Hill

Merton Hill

BRONZITE HILL

Fingerpost

HIGHWAY

MURCHISON

RIVER

Wilson River

Little Wilson River

Boco

Sophia

MACKINTOSH

AIRBORNE SURVEY SPECIFICATIONS — WESTERN TASMANIA — SURVEY, MAY, 1981

MAGNETOMETER: SONOTEK IGSS 1 SYSTEM, Cessna 185 E aircraft
Sensitivity 0.1 nT

RECORDING INTERVAL 40m sampling (approx.)

MEAN SENSOR HEIGHT 135m

NOMINAL LINE SPACING 500m

FLIGHT PATH RECORD 16mm ground tracking camera

RESIDUAL MAGNETIC CONTOURS

MAGNETICS Tie Line levelled

GRID MESH SIZE 125 x 125 m

GRID FILTER Simple 3 pt filter

CONTOUR INTERVAL 5nT

Flown by Geox Pty Ltd

ARTHUR RIVER | HELLYER | PIEMAN | SOPHIA

083069

RGC EXPLORATION PTY. LIMITED

INCORPORATED IN NEW SOUTH WALES

COMPILED	B.W.W.	
DRAWN		
DATE	OCT., 1990	
CHECKED	B.W.W.	
1:250,000 Reference		

WARATAH - MT. RAMSAY AREA

RESIDUAL MAGNETIC CONTOURS

BASE PLAN No.

OVERLAY PLAN No.

SCALE 1:50,000 0 0.5 1.0 1.5 2.0 3.0 Km.

PLATE 15

360000mE
370000mE
380000mE
390000mE
5420000mN
5410000mN
5400000mN
5390000mN
5380000mN

E.L. 15/90
E.L. 12/90
MT. BISHOF
R.L. 8807
WARATAH
△MT. CLEVELAND
LUINA
CLEVELAND
E.L. 12/90
△MT. PEARSE
E.L. 15/90
MT. RAMSAY△
E.L. 12/90
E.L. 15/90

5 cm

MAGNETIC SOURCE
MAGNETIC SOURCE (REVERSELY MAGNETISED)
MAGNETIC LINEAMENT
MEREDITH GRANITE
TERTIARY BASALT

ARTHUR RIVER | HELLYER
PIEMAN | SOPHIA

083070

RGC EXPLORATION PTY. LIMITED
INCORPORATED IN NEW SOUTH WALES

B.W.W.
L.D.K.
DEC., 1990
B.W.W.

WARATAH - MT. RAMSAY AREA

MAGNETIC INTERPRETATION

1:50,000 0 0.5 1.0 1.5 2.0 3.0Km.

PLATE 16

083071

APPENDIX 2

An interpretation form of Waratah Granite

085072

LEAMAN GEOPHYSICS
Survey Review, Specification, Reduction, Interpretation
Gravity, Magnetic and Seismic Methods
Structure and Prospect Evaluation

Registered office:
3 MALUKA STREET, BELLERIVE, TAS. 7018
All correspondence to:
GPO BOX 320 D, HOBART, TAS. 7001
Telephone: (002) 44 1233
Fax: (002) 44 6674

AN INTERPRETATION FORM OF MEREDITH GRANITE

WARATAH AREA
EL 12/90 AND 15/90

for
RGC Exploration Pty Limited

by
Dr. D.E. Leaman

January 1991

MEREDITH

T/91/25

CONTENTS

page

SUMMARY

INTRODUCTION 1

GRAVITY SURVEY 2

INTERPRETATION
Rocks and rock properties 3
Features of the gravity field ... 4
Methods 5
Discussion 6

CONCLUSIONS AND RECOMMENDATIONS 12

REFERENCES 13

APPENDIX: DETAILS OF GRAVITY DATA BASE AUGMENT

FIGURES

1 Location map
2 Previous interpretation of Meredith Granite, Waratah area
3A Map of residual Bouguer anomaly
3B Map of residual magnetic field
3C Overlay of gravity field on magnetic field
4 Trends
5 Location of profiles discussed in text
6 2D model line 5
7 8
8 9
9 10
10A 2D gravity model regional line 7
10B 2D magnetic model reigonal line 7
10C Geological integration of line 7 models
11 3D model line 1
12 1 source components
13 2
14 3
15 4
16 5
17 6
18 8
19 9
20 10
21 11
22 12
23 13
24 14
25 15
26 Current granite model, Meredith Granite
27 Granite model and regional trends
28 Granite model and target zones
(All figures follow the text)

MAPS

1 Bouguer anomalies showing licence boundaries
2 Residual Bouguer anomaly
3 Current granite model interpretation
(Maps are included in separate pockets)

SUMMARY

Review of extant gravity data, supplemented by a new infill survey, in the Waratah area has enabled considerable refinement of previous models and concepts of the form of the Meredith Granite in the region north and east of its principal exposure south west of Waratah.

This interpretation, although limited by assumptions concerning the bulk properties of the intruded rocks and the scale and nature of the disruption of the sequences about the intrusion, has defined several definite elements of the granite form.

There is a clear correlation between granite form and many observed fault, fold and trend patterns across the licence areas. It may be inferred that the granite has been affected by, and modified high level manifestations of, pre-existing fractures.

There is also a positive association with known mineralisation, particular forms of the granite and those structures which presented primary controls on intrusion. Few of these are obvious in surface mapping which may not distinguish them due to existence of other faults, or due to their modification upon intrusion or deformation.

The association between granite form and mineralisation is of two types.
Small prospects or mines may be related indirectly to the granite by significant structures. The Magnet mineralisation is of this type; the structural link being the intrusion-fault which has plumbed a fluid source beneath.
Other sites, including Cleveland and Mt Bischoff, are more directly associated with the primary granite fracture system developed during intrusion. Prospects of this type are associated with projections of the wall forms.
Both associations can only be appreciated or recognised after basic interpretation of the form of the granite.
Due to the form of the Meredith Granite the potential for other sites is generally limited to an area about the northern extemity of the pluton.

The interpretation also indicates some depth limitation of potential hosts and deposits. Modelling and profile analysis has shown that a variable density contrast profile is required between the granite and the intruded rocks. This may mean local siliceous cappings but the general implication is for termination of Dundas Group, Oonah, Crimson Creek and Success Creek Formations at an average depth of about 1500 m - where such a volume is available above the granite - and that the root of the granite is emplaced in predominantly siliceous basement.

The present work provides a guide for further exploration.
It has also identified anomalous structures and sequences within the Cambrian rocks to the east of the granite and further work, using both gravity and magnetic data, is recommended in order to satisfactorily define principal relationships.

083075

INTRODUCTION

ELs 12/90 and 15/90, titled Waratah, extend south from Waratah and Mt Bischoff in north western Tasmania (see Figure 1).

This area contains a number of small prospects and a few significant deposits; base metals at Magnet, tin at Cleveland and Mt Bischoff.

In order to understand the relationships between these deposits and the Devonian Meredith Granite, exposed several kilometres southwest of Waratah, gravity data have been reviewed and interpreted.

Previous interpretations of more regional and indicative nature have been reported which demonstrate the efficacy of this approach. These were summarised in Chapter 15 of Leaman & Richardson (1989a). Figure 2 presents the net conclusion of this previous work which shows that the granite was inferred to shelve shallowly north toward Waratah. Many contacts were thought to dip very steeply.
This diagram suggests the general relationship between mineralisation and granite shelf but does not provide any specific guidance for prospect or EL evaluation.
It was based on simplified analysis of the regional data base current in 1987.

This report examines the gravity data within the Waratah - Mt Ramsay - Guildford area in order to provide a more refined and specific description.

The gravity data used were obtained from the state gravity data base (TASGRAV) maintained by the Department of Mineral Resources and Energy. This data was supplemented by infill surveys undertaken by RGC Exploration. Map 1 presents a compilation of this data for the area centred on the licences. All Bouguer reductions include a 22 km terrain correction and use of crustal density 2.67 gm/cc. The nominal station spacing is about 700-900 m although some large gaps are present. All station locations are shown in Maps 1 and 2.

The present report describes preliminary analysis of this data using three dimensional methods applied to residual profiles generated after removal of the deep crustal and oceanic components as defined by Leaman & Richardson (1989b). This presentation provides a normal relativity of response patterns free of regional gradient distortions.

Interpretation has been halted at the point at which the general form of the granite has been defined (as requested) and any secondary issues, such as anomalous elements of the Cambrian succession, have only been treated in detail sufficient to ensure that the granite interpretation is sound. There is scope for much further analysis of these issues.

GRAVITY SURVEY

The extant Tasmanian gravity data base for the Waratah - Guildford - Ramsay area consisted of stations approximately 1 km apart. These had been acquired during a late phase of the Mt Read Volcanics Project by the Department of Mines and given a first interpretation by Leaman and Richardson (1989a).

The coverage was adequate to define the gross extent of the Meredith Granite and suggest elements of its form. It was not sufficient for any specific exploration or evaluation of moderate areas. An infill survey with a nominal spacing of 500 m was proposed in the area surveyed by the Department for those areas in which the gravity field appeared depressed and for which shallow granite might be an explanation. Such sites occur in the basalt-covered area south of Waratah.
Much of the Mt Ramsay - Cleveland area, and especially the region north of the Luina road west of Waratah, had never been surveyed previously. Terrain, budget and objectives led to survey design compromises such that reasonable areal definition was obtained. All availabe access, foot and vehicle, was used to obtain on-line spacings of about 500 m even though large gaps in coverage remain. Some lines were cut in areas where no access was available. The ultimate coverage was determined by available access and gridding. Map 1 indicates the final status of coverage. A large gap in coverage north of Luina has not been significantly reduced due to limited access and its location beyond the present licence areas.

The survey was undertaken by Mackie Martin Pty Ltd (Bettina Townrow observer) during November and December 1990 to specifications compatible with the Mt Read coverage. It was linked to state datum at Waratah and Que River. All stations were corrected for instrumental drift (and tide) and referred to an internal tie network. Reproducibility of observed gravity is of the order of 0.03 mGal.
All elevations were determined barometrically using survey spot points for control. Elevations are considered reproducible and reliable within 2 to 3 m generally although several spot ties were made within the practical limit of 1 m.
All stations were terrain corrected to a radius of 22 km. Corrections range up to 3 mGal and may have an error of 5% due to uncertainties about precise location within the terrain and limited local descriptions about the station. Most corrections were, however, less than 0.5 mGal.

The data have been contoured using a 1 mGal interval due to these limitations.

Data was reduced using the 1930 ellipsoid compatibly with the Tasmanian data base with a Bouguer density of 2.67 gm/cc.

A tabulation of the infill survey is given in the Appendix.

INTERPRETATION

ROCKS AND ROCK PROPERTIES

No specific study of rock properties has been undertaken in order to provide the present interpretation. All work has been referenced against the geological maps of Groves et al (1972) and Corbett and McNeill (1988) pending preparation of original maps by RGC exploration (in progress at time of writing).

Late Precambrian rocks of the Oonah Formation are exposed within the area. These include carbonates and various other sediments but are intensely deformed and faulted. Leaman (1986, 1990) has argued that the fault-bounded blocks of Oonah Formation represent pop-up blocks dislocated from thrust slices. Dundas Group and various lower Cambrian rock suites are exposed across the bulk of the area studied. These units are intensely faulted and contain numberous slices of mafic and ultramafic rocks. Several deformed lower Cambrian thrust sheets have been inferred (e.g. Figure 10).
Leaman (1986), and Berry and Crawford (1988), have suggested major thrusts within this association of structures. Eastward translation of the Oonah Formation across a large part of the Lower Palaeozoic sections was implied by Leaman (1986). Similar displacements are implied within the Cambrian rocks. East facing structures are impressed upon these and the most notable is the Rosebery Fault.
The resultant pattern is structurally complex and not yet adequately mapped or defined.

The Meredith Granite has intruded this complex structural regime.

Nominal rock properties based on general experience of the exposed materials have been employed but tested against the controls afforded by the magnitude of anomalies across the granite itself.
Many figures (e.g., 11, 20 and 21) provide examples of these tests to suggest that the appropriate bulk contrast between granite and country rock - as a package - does not exceed 0.12 gm/cc.
Since the bulk density of the granite is of the order of 2.61 to 2.63 gm/cc then the implied bulk density of the intruded rocks is about 2.75 gm/cc. This is consistent with sample determinations obtained during the Mt Read Volcanics Project for both Oonah Formation and most Cambrian rocks (excluding mafics).
Ordovician, Silurian and Devonian rocks are also variable but generally much less dense and have little bulk occurrence within the Waratah area. Siliceous rocks may have densities of 2.5-2.55 gm/cc while carbonates may be 2.74-2.84 gm/cc. Siliceous Tyennan-style Precambrian rocks have bulk densities of the order of 2.67 gm/cc.
Modelling of the marked regional negative Bouguer anomaly due to the granite has shown, however, that the maximum contrast cannot be applied to the entire intrusion implying that there are significant density variations within either granite or country rocks at high levels or that the contrast between granite and intruded rocks decreases with depth. High level variations in intruded rocks are to be expected in this area but the second observation suggests a predominantly siliceous section at depths greater than about 2000 m.

083078

FEATURES OF GRAVITY FIELD

Maps 1 and 2 present the gravity field; Map 1 shows the observed Bouguer anomalies and Map 2 residual anomalies after removal of crustal effects using the method of Leaman & Richardson (1989b). Figure 3 presents the residual anomalies and compares them with the magnetic field. All discussion is based on the residuals.

The gravity field is variable in intensity across the entire area but the dominant anomaly is associated with the exposure of the Meredith Granite. A strong negative gradient is associated with the outcrop limits.

Other large anomalies, usually positive, are not completely defined being beyond the objectives of this project.. They include the effect of the Heazlewood ultramafic complex, the southern extension of the Burnie Formation north of Waratah, and a belt of positive anomalies extending north east of Pinnacles. Other variations in the field are much subtler.

Review of the available geological compilations and correlations with the gravity field shows that many secondary gradients observed within the principal anomalies rarely reflect mapped patterns - in terms of orientation and nominal position. There is also no distinguishable interference due to the blanket of Tertiary sediment or basalt. The gravity field, as displayed in Figure 3, is applicable to the rocks beneath this cover. In particular to many of the rocks beneath the exposed sequences as well. This is an essential corollary to the limited correlation between mapped features and anomaly patterns. Relatively thin coverings, either as units or structures, are implied throughout the area, much as suggested by Leaman (1986). The additional definition afforded by the upgraded coverage leaves little doubt on this point.

Figure 4 presents a basic trend compilation using available geophysical data; geological data is too restricted to be reliable. Dominant trends are NE-SW, NW-SE although elements of near east-west and north-south trends are also visible. The irregular distribution of data suggests that all these orientations are real and not artifacts of coverage.

Figure 3C shows that the positive anomaly belt north of Pinnacles correlates directly with a magnetic feature. Surface mapping of this area indicates Cambrian sediments (correlates of the Stitt Quartzite?) and the geophysical response is not consistent with either the implied mapping or the correlation. Many such couplings can be identified but the generally low detail of extant geological mapping limits evaluation. The association of mineralisation with the feature described above suggests that is may well be prospective and worthy of inspection. Such anomalies, generated within the sedimentary succession (?), are not treated in detail in this report due to the prescribed emphasis on the form of the Meredith Granite.

METHODS

An array of observed profiles was established based on Map 2. These profiles were designed to provide a representative sampling of semi-regional anomaly forms. The nominal line spacing was about 5 km and may be contrasted with the nominal station spacing of 0.5 km. The particular location of the test profiles reflects the actual station coverage in order to minimise interpolations - especially north of Luina.
This report represents a preliminary interpretation which is also an order of magnitude upgrade of previous interpretations. It is not an overinterpretation of the data available.

The "observed" profile used as reference in all model calculations is the residual derivation and thus excludes long wavelength effects such as the response due to the continental margin and continental thickening. The effects of other regional granites remain and have been modelled along with the particular object of study. In this study the crustal formulation of Leaman and Richardson (1989b) was used to create the residual responses from the observed data and the primary model of the other granites of NW Tasmania given by Leaman and Richardson (1989a) was used to estimate the contributions of other granites. The other granites are sufficiently distant that any inadequacies in this component are not significant. The observed profile may be used for direct comparison with model calculations.

The analysis has been directed at primary review of granite form only.

Some N-S and E-W profiles were then modelled using simple two dimensional methods in order to scale the general relief of the granite and estimate the order of the density contrast applicable. This was achieved by extending some lines eastward but many lines close to the boundaries of the main anomaly (due to granite) cannot be modelled using 2D methods. These simple methods were also used to test whether the density contrast applied over considerable depths, say greater than 3 km, and whether it was possible to model viably using the granite density alone as a bulk reference.

Those profiles referred to in this text are located in Figure 5.

Figures 6, 7, 8 and 9 exemplify typical 2D models. These suggest that a variable contrast must be applied against the granite, with depth, since the Cambrian sequence appears less than 2 km thick.

Initial 3D tests confirmed that a bulk contrast of about +0.12 gm/cc cannot be used across the entire depth range of the granite. The particular requirement of contrast varies little for each line, the nature of surface rocks and structures or the orientation of the profile. These tests showed that no

single contrast can be employed but that the minimum contrast required is about +0.05 gm/cc. These indications are consistent with intrusion into a predominantly siliceous sequence.
It was found that the effective contrast pattern for the intrusion was about 0.12 gm/cc from 1000 to 2000 m reducing rapidly to 0.05 gm/cc at 3000 to 3500 m. There is no suggestion in this area that any spiky crests of the granite contain granite which is more siliceous and of lower density than the bulk of the body. This would suggest that a substantial part of the body has been unroofed. The density contrast pattern suggests that the Cambrian-Oonah Precambrian sequence is not much thicker than 2000 m and that the bulk of the intruded rocks is composed of Tyennan type Precambrian rocks.

Since the form of the granite is irregular and generally offers angular projections to the north three dimensional methods must be employed to generate reliable proportions for the form of the granitic mass. Two dimensional methods are quite unreliable for many profiles. The information derived from 2D tests and previous work was used to produce an initial 3D model which was passed through six iterations to produce the solution offered in this report.

Each profile was examined after each iteration and tested for consistency in both body geometry and curve fit parameters. The latter provide a crucial test of model and assumptions and must be consistent for all lines of the array. The difference between the zero shift and the observed and calculated difference is a measure of this and has been found to be about 5.0 mGal for this data set in the form used and using the variable density contrast profile defined above. The deviation for best fits is about 0.5 mGal which approximates the noise level in the data.
Note that a residual data set should yield a near zero difference, not 5.0 mGal, and would if the density contrasts were referred to the Bouguer density; viz -.06 gm/cc for granite and +0.07 mGal for other formations and the granite and all other units are properly depth scaled. The fit depends on a balance of negative effects due to granite - the root depth assumed becomes critical in this - and positive effects due to other formations. A consistent root of 9 km has been used since this element is the least sensitive and important to the study..

DISCUSSION

Fifteen profiles of the profile array established to evaluate the gravity field and define the form of the Meredith Granite in this area are shown in Figures 6 to 25. All granite profiles shown represent the current status of the 3D model. Further refinement of this model (Map 3) is not justified until some control is established within the study area, or the contrast assumptions can be significantly revised.

Each profile has been discussed separately in order to indicate the nature of the conflicts, issues and targets implied in different parts of the structure.

No attempt has been made to include geological details in each section due to the uncertainties in extant mapping. Figure 10C, however, suggests the general relationships which can be inferred from application of both gravity and magnetic data sets. Extension of these concepts requires concurrent magnetic modelling to the entire area. This has been beyond the scope of the present study which has been directed toward definition of the form of the Meredith Granite and which has only considered other elements of the geology, in gross terms, in order to assess implications for the granite model. There is clearly scope for comprehensive analysis of the structural setting.

The 2D models of the granite presented in Figures 6, 7 and 9 indicate that the Meredith Granite is largely unroofed and that its walls dip steeply. These models also show that the Cambrian sequences, or Oonah Formation, are relatively thin. Local anomalous sequences are also evident. These are clearly evident in Figure 9. The source appears to lie either in the thin sequence above the Rosebery Fault, or within the thickened overthrust section beneath it.
Figure 8 examines the possible contribution due to the Heazlewood mafic complex. It assumes that granite does not contribute significantly to the profile (contrast Figure 19) and shows that such sources may be significant and may counterweight negative effects due to granite.
Only 3D modelling and comprehensive interrelationships between profiles can resolve many aspects of such problems. Some ambiguity must remain until all geological units are treated in the detail applied to the granite.

The section line (line 7) used for research in NW Tasmania by the Geology Department of the University of Tasmania has been modelled and evaluated in order to gain a better idea of unit relationships and their influence on granite response.
Figures 10A and B provide concordant gravity and magnetic evaluations. These complex models are consistent with current geological knowledge. The natural scale, labelled section shown in Figure 10C transforms these models into a true section. This solution should not be accepted outright; it is simply a solution which explains the two fields and is consistent with current control information. It includes assumptions about the continuity of Crimson Creek Formation, for example, which may not be valid. All such presumptions are secondary and the style and implications of the section are probably valid.

The 2D analysis demonstrates that it is not possible, in this area, to simply treat the granite in isolation and refer it against relatively uniform or thick materials. The 3D models have, therefore, included gross assessment of both Cambrian rocks to the east of the granite, and mafics and Burnie Formation to the north and west.

Refer to Figure 5 for the location of all lines.

Line 1 (Figures 11, 12).
This section samples the eastern part of the intrusion. The model accounts for the general character of the profile but the imperfections evident at the northern end of the profile are partly due to limited data control and partly to the gross mass balance between Burnie Formation and the granite. This is illustrated in Figure 12 where the primary components of the calculated resultant are displayed separately. It is evident that it would be possible to counter some variation in the granite form by some variation in the Burnie Formation - although there are limitations. The outer limit of the granite wall cannot be shifted any great distance, however.
The steep-sided nature of the granite is clearly evident. The Magnet mineralisation can be linked to possible wall fractures.

Line 2 (Figure 13)
This profile samples the Cambrian rocks east of the granite, which due to its very steep eastern face, barely appears in this section. The principal elements in the fit are the thickness of Burnie Formation in the north and the extent of denser Dundas Group locally. The spiky zone with an amplitude of about 4 mGal represents the belt of anomalous units. But all are far in the roof of the granite.

Line 3 (Figure 14)
This section passes through the Waratah area and Mt Bischoff is barely off line. It shows, more clearly than line 2, the relative responses of the granite and Burnie Formation (to the north) and their mass balance which yields a smooth integration, and the influence of the anomalous units in the Dundas(?) Group to the south. The deviation is due to these units and not any variation in form of the granite.
The Bischoff mineralisation is clearly associated with the fracture system and pinnacles on the margin of the intrusion. The general asymmetry of the granite roof may also prove of interest within the basalt covered areas.

Line 4 (Figure 15)
This profile illustrates the very steep margin of the granite. The fit is imperfect due to non inclusion of any concept for the anomalous belt within the Dundas(?) Group and the detailed nature of the exposed granite. The presence of dense units off section nearby also affect the end of the profile.
Corbett & McNeill (1988) indicate two granite outcrops far removed from the main exposure at the northing of this line. These are clearly no part of the Meredith Granite and may not be in situ - if properly located.

Line 5 (Figure 16)
This line is similar to line 4 in all characteristics except the clear accounting of the form of the profile at its western end due to sampling and presence of more granite. The anomalous zone within the Cambrian rocks to the east is much more limited.

Line 6 (Figure 17)
The profile also samples the tip of the granite in the Waratah area. The Magnet Mine is virtually on profile and Mt Bischoff is a little distance from it. There are obvious associations between mineralised sites and the margins of the intrusion. The Magnet dyke, and the associated fracture system, may also be related.
The western end of the profile is not well fitted indicating the presence of more mafic material than included.

Line 8 (Figure 18)
This profiles examines the structure near Cleveland. The observed field is not well defined in this region. Nor have the regional assumptions related to the Heazlewood Complex been adequate. No further refinements are justified until more data is available. The shape of the granite is not absolutely determined in the Cleveland area due to the possible, uncontrolled, mass trade offs. It does suggest that granite is not far away and that the controlling fracture systems related to the granite margin have been important.

Line 9 (Figure 19)
The shape of this profile disguises the effect of the granite pillar due to the balance between mafics to the west and Burnie Formation to the north. The overall relationships, extracted only by 3D methods, are evident. The association between the margins of the intrusion and the mineralisation north of Magnet and at Mt Bischoff is unambiguously anticipated.

Line 10 (Figure 20)
The profile and its implications are similar to those described for line 5.

Line 11 (Figure 21)
This line conceals some of the ambiguities present in the present interpretation. It is not evident by inspection how far the mass balance between granite and Burnie Formation in the the north allows a shift of the interpreted granite margin. The dip in observed profile at 29 km could be argued as being due to a further granite spine. Such a spine would then require the main granite wall to be at least 5 km further north than shown in the figure. This is not sustainable three dimensionally and the anomaly dip is probably due to Tertiary elements.
The positive change at 27 km and the steepened gradient to the south can be related directly to the granite margin even though concealed by Tertiary cover. It is possible that the model shown places the granite margin 2 km too far north. The model is a compromise given present control. The position of the margin can be further evaluated only after consideration of anomalies sourced in the country or roof rocks and these details have been beyond the scope of the present project.

Line 12 (Figure 22)
This line samples the eastern wall of the Meredith Granite and the glancing orientation has produced the high relief nature of the model shown. The distribution of the Burnie Formation, in

the north, has been inadequately treated but the model does account for the eastern edge of the pluton.

Line 13 (Figure 23)
Granite is not significant along the profile which essentially samples the anomalous belt within the Dundas Group. This profile clearly suggests the scale of responses due to these materials. The mineralisation at Silver Falls is off line and appears marginal to one of the dense sources. The pattern is not dissimilar to that at Hellyer and is worthy of verification. The match for Burnie Formation at the northern end of the profile is adequate.

Line 14 (Figure 24)
This section also samples the Cleveland area. Although the problems of data coverage (see line 9) remain the model and control is better at the orientation of this line and it is possible to establish a clear structural correlation between the mineralisation and the granite.

Line 15 (Figure 25)
This profile re-inforces the suggestion of a relationship between the location of the Magnet dyke and mineralisation and the granite margin. The mismatch near the crest of the granite may imply that the model in bulk terms is too shallow in this area.

It should be recognised that the current model solution and interpretation of the Meredith Granite as presented in Map 3 and Figure 26 has incorporated some mass balance assumptions, has ignored some contributing details (as within the Cambrian succession) and is only defined where model profiles have been sampled and examined. Most of those profiles are included in this report and it is clear that some large gaps do exist between them. The shape of the gravity field and the effects and requirements of 3D forms and models, however, do minimise the possibilities and the deficiencies are not considered large.
The model is generally untested and uncontrolled at this date.

The model establishes the very steep contact zones along the eastern face of the Meredith Granite. Any ribbing present is minimal and restricted to fracture corners. The shape of the granite is clearly defined in regional stress terms by the extensive lineaments in both gravity and magnetic data sets (Figure 4) which are simply extensions of the granite corner orientations. Intrusion of the granite has almost certainly been controlled by pre-existing features, few of which are unambiguously exposed, but which dominate the upper crust.

The granite shelves northward with some crestal pinnacles. Mineralisation between Luina and Waratah is associated with these pinnacles, or extensions of the fracture systems which define the margin of the pluton. The pluton is more extensive to the north west and may extend more than 10 km NW of Waratah. It is, however, largely unroofed to the south of Cleveland.

There is some scope for further exploration in this region although much of the granite-bearing zone is beyond the present ELs, good data coverage and firm geological knowledge.

The basalt-covered area SE of Waratah may also disguise granite related mineralisation (near 374 000 mE, 5404 and 5407 000 mN). More detailed review of these areas is necessary.

The association between trends and granite form is illustrated in Figure 27 and the inferred zones of prospective interest for granite-related mineralisation are shown in Figure 28.

The surveys have also drawn attention to the positive anomalies within the Cambrian rocks. These may have exploration significance given the lack of current knowledge and the presence of base metal mineralisation within the zone. Evaluation of these materials is suggested grossly in Figure 10C but has been generally beyond the scope of the present interpretation. Possible links and correlation with Que- Hellyer mineralisation may be inferred but there is need for more analysis and ground checking of rock units.
Figure 28 also outlines the anomalous zones within the intruded Cambrian rocks which may carry other mineralisation styles.

CONCLUSIONS AND RECOMMENDATIONS

The present interpretation of gravity data in the Waratah - Guildford - Mt Ramsay area has established that the Meredith Granite is mostly unroofed with steeply dipping margins to east and south. The roof of the granite plunges shallowly northward beneath Waratah and it is in the area between Luina, north Bischoff and south Waratah that some potential exists for granite-related or controlled mineralisation at shallow depths. The known mineralisation at Cleveland, Magnet and Mt Bischoff has been explained by the interpreted granite forms. Some other sites exist within the region.

There seems limited potential for large deposits along the eastern face of the granite due to its sheer form. The fractures which impinge on this may be important, however, in transferring fluids. Many of these lateral and transverse structures appear to be large and are not generally to be inferred from current mapping. Many clearly pre date the granite and have controlled its emplacement. Many may also have influenced other types of mineralisation.

Attention has been drawn to the anomalous zones within the Cambrian(?) rocks adjacent to the granite although the present evaluation has not reviewed them in detail. At least one of these zones is mineralised. The data available (gravity and magnetic) are able to contribute to appraisal and definition of these zones.

The present work has been restricted to primary appraisal of the Meredith Granite and has considered incidental issues only in so far as this allows some estimation of the reliability of the granite model. Many assumptions are implicit although it is believed that the general form reported is sound. Depth estimates must be treated as estimates pending control and revision. Any secondary revision must incorporate refined judgments about other rock sources (ultramafics, Burnie Formation, anomalous Cambrian) and cannot be justified until this is possible. Improved base mapping may well provide such control.

The principal recommendations relate to features east of the granite. The gross trends identified by the survey and their implication for non granite mineral styles must be evaluated. The extant gravity and magnetic data are able to support any structural evaluation of these features.
The general form of the interpreted granite around Waratah is sufficient to guide exploration. In the basalt-covered area south of Waratah, near the granite rib at 375 000 mE, 5404 000 mN, further gravity analysis should not be undertaken without control afforded by use of the magnetic data. The latter can provide estimates of basalt thickness.

The interpretation also suggests that the Late Precambrian and Early Cambrian rocks of the area are stratigraphic or structural veneers little more than 2 km thick.

083087

REFERENCES

Berry, R.F., & Crawford, A.J., 1988. The tectonic significance of the Cambrian allochthonous mafic-ultramafic complexes in Tasmania. Aust. J. Earth Sci., 35, 161-171.

Corbett, K.D., & McNeill, A.W., 1988. Map 6. Geological compilation map of the Mount Read Volcanics and associated rocks. Hellyer to South Darwin Peak. Mt Read Volcanics Project Dep. Mines Tasm.

Groves, D.I., Martin, E.L., Murchie, H., & Wellington, H.K., 1972. A Century of tin mining at Mount Bischoff 1871-1971. Bull. Geol. Surv. Tasm., 54.

Leaman, D.E., 1986. Gravity interpretation west and north west Tasmania. Mt. Read Volcanics Project Report Dep. Mines Tasm.

Leaman, D.E., & Richardson, R.G., 1989a. The granites of west and north west Tasmania - a geophysical interpretation. Bull. geol. Surv. Tasm., 66.

Leaman, D.E., & Richardson, R.G., 1989b. Production of a Residual Gravity Field Map for Tasmania and some implications. Exploration Geophysics, 20, 181-184.

Leaman, D.E., 1990. The shifting basement of Tasmania. Paper given to the Max Banks Symposium, Hobart, Dec. 1990. Geol. Soc. Australia publication.

083088

Report submitted on behalf of
Leaman Geophysics
by

D. Leaman

Dr. D.E. Leaman, B.Sc., Ph.D
F. Aus.I.M.M., M.M.I.C.A

APPENDIX

TABLE OF REDUCED GRAVITY DATA
Survey code 9049

The table lists all data acquired by RGC Exploration during 1990 in the ~~Wa~~ area.
The reduction density used was 2.67 gm/cc.

The table lists the station number (with year and survey code)
position (easting, northing, elevation)
gravity (observed, theoretical)
terrain correction
Bouguer anomaly

Observed gravity is based on state datum tie points and the theoretical gravity is based on the 1930 ellipsoid.

RGC Exploration - Waratah Area Gravity Data

Station No.	East (m)	North (m)	Elev (m)	Gobs	Gtheo	Terr Corr	B.A. mgal
9049.9900	[redacted]			980.178698	980.309596	0.53	-9.97
9049.9901	[redacted]			980.187758	980.305434	0.13	2.46
9049.9902	[redacted]			980.172837	980.313296	0.22	-10.99
9049.9903	379560.0	403580.0	640.00	980.176180	980.315214	0.16	-12.97
9049.9904	[redacted]			980.177862	980.311662	0.58	-9.08
9049.1001	[redacted]			980.138480	980.307978	4.44	-10.62
9049.1002	[redacted]			980.180991	980.313632	0.12	-11.53
9049.1003	[redacted]			980.181215	980.313960	0.10	-12.84
9049.1004	[redacted]			980.180920	980.314012	0.12	-12.77
9049.1005	[redacted]			980.177791	980.314207	0.16	-14.48
9049.1006	[redacted]			980.182061	980.314550	0.20	-14.45
9049.1007	378230.0	404300.0	634.00	980.175447	980.314616	0.15	-14.29
9049.1008	379070.0	403820.0	657.00	980.171594	980.315014	0.16	-14.01
9049.1009	[redacted]			980.161168	980.313580	0.47	-17.97
9049.1010	[redacted]			980.165448	980.313910	0.30	-19.11
9049.1011	[redacted]			980.162095	980.314094	0.36	-19.63
9049.1012	[redacted]			980.163756	980.314723	0.18	-20.75
9049.1013	[redacted]			980.167293	980.314983	0.15	-21.24
9049.1014	[redacted]			980.170819	980.313353	0.21	-12.09
9049.1015	[redacted]			980.177465	980.312460	0.19	-9.88
9049.1016	[redacted]			980.171349	980.312340	0.25	-9.13
9049.1017	[redacted]			980.169821	980.312275	0.31	-11.91
9049.1018	[redacted]			980.163970	980.312648	0.53	-10.83
9049.1019	[redacted]			980.161198	980.312220	0.68	-10.27
9049.1020	[redacted]			980.162421	980.312090	0.73	-10.83
9049.1021	[redacted]			980.178443	980.311989	0.33	-6.92
9049.1022	[redacted]			980.179085	980.311606	0.48	-6.53
9049.1023	[redacted]			980.165999	980.312861	0.23	-12.26
9049.1024	[redacted]			980.187065	980.304893	0.17	2.15
9049.1025	[redacted]			980.189532	980.304366	0.32	3.72
9049.1026	[redacted]			980.187534	980.304478	0.41	2.88
9049.1027	[redacted]			980.190368	980.304688	0.71	2.06
9049.1028	[redacted]			980.199194	980.304604	0.92	2.72
9049.1029	[redacted]			980.206991	980.304447	0.83	-1.22
9049.1030	[redacted]			980.218549	980.304192	0.86	3.94
9049.1031	[redacted]			980.231319	980.303569	1.07	5.54
9049.1032	[redacted]			980.245150	980.303207	0.97	3.89
9049.1033	[redacted]			980.247769	980.303192	0.86	3.47
9049.1034	[redacted]			980.245537	980.303557	0.70	1.69
9049.1035	[redacted]			980.243600	980.304067	0.77	1.28
9049.1036	[redacted]			980.241542	980.304595	1.10	0.60
9049.1037	[redacted]			980.226549	980.303962	1.61	2.88
9049.1038	[redacted]			980.223430	980.304695	1.69	2.06
9049.1039	[redacted]			980.229658	980.304639	1.57	1.54
9049.1040	[redacted]			980.242021	980.304275	1.42	2.51
9049.1041	[redacted]			980.247881	980.303623	0.86	3.15
9049.1042	382570.0	414900.0	623.00	980.184599	980.306127	0.11	1.14
9049.1043	[redacted]			980.183804	980.305928	0.31	-2.01
9049.1044	[redacted]			980.186964	980.305565	0.34	-0.03
9049.1045	[redacted]			980.184538	980.306316	0.48	-3.46
9049.1046	[redacted]			980.184956	980.306456	0.50	-1.19
9049.1047	[redacted]			980.185628	980.305815	0.24	0.45
9049.1048	[redacted]			980.182275	980.310304	0.32	-9.08
9049.1049	[redacted]			980.180043	980.310683	0.23	-10.41
9049.1050	[redacted]			980.181541	980.311271	0.25	-9.67
9049.1051	[redacted]			980.191570	980.310822	0.72	-7.58
9049.1052	[redacted]			980.195688	980.310269	1.69	-8.04

RGC Exploration - Waratah Area Gravity Data

Station No.	East (m)	North (m)	Elev (m)	Gobs	Gtheo	Terr Corr	B.A. mgal
9049.1053	[redacted]			980.200366	980.309824	0.92	-8.60
9049.1054	[redacted]			980.203169	980.309467	0.81	-10.08
9049.1055	[redacted]			980.207348	980.309154	0.66	-9.28
9049.1056	[redacted]			980.179228	980.312142	0.30	-10.45
9049.1057	[redacted]			980.176150	980.311942	0.56	-10.11
9049.1058	[redacted]			980.179493	980.311718	1.07	-10.17
9049.1059	[redacted]			980.186179	980.308728	0.98	-5.89
9049.1060	[redacted]			980.201069	980.308484	1.31	-6.56
9049.1061	[redacted]			980.208010	980.308591	1.11	-6.42
9049.1062	[redacted]			980.180777	980.314494	0.16	-12.18
9049.1063	[redacted]			980.182092	980.314902	0.18	-13.02
9049.1064	[redacted]			980.181185	980.315317	0.20	-13.73
9049.1065	[redacted]			980.181256	980.314441	0.10	-11.31
9049.1066	[redacted]			980.182459	980.314851	0.12	-10.10
9049.1067	[redacted]			980.183050	980.315228	0.13	-12.83
9049.1068	[redacted]			980.190582	980.305534	0.61	4.28
9049.1069	[redacted]			980.189491	980.305873	0.97	3.61
9049.1070	[redacted]			980.184324	980.306213	0.29	1.36
9049.1071	[redacted]			980.160556	980.313564	0.17	-18.47
9049.1072	[redacted]			980.164225	980.313996	0.22	-19.32
9049.1073	[redacted]			980.172491	980.314342	0.19	-20.48
9049.1074	[redacted]			980.175844	980.314754	0.18	-20.50
9049.1075	[redacted]			980.182989	980.315118	0.17	-18.84
9049.1076	[redacted]			980.186169	980.315565	0.18	-18.07
9049.1077	[redacted]			980.175905	980.315908	0.26	-17.18
9049.1078	[redacted]			980.174448	980.316338	0.29	-19.04
9049.1079	[redacted]			980.180359	980.316709	0.28	-19.02
9049.1080	[redacted]			980.175396	980.316964	0.26	-20.32
9049.1081	[redacted]			980.174692	980.316693	0.21	-20.21
9049.1082	[redacted]			980.174274	980.317654	0.26	-20.16
9049.1083	[redacted]			980.173663	980.317674	0.28	-19.79
9049.1084	372290.0	399920.0	600.00	980.182744	980.318065	0.26	-17.03
9049.1085	371730.0	399230.0	580.00	980.181613	980.318613	0.42	-22.48
9049.1086	371720.0	398820.0	576.00	980.182683	980.318944	0.50	-22.45
9049.1087	371810.0	398390.0	581.00	980.182357	980.319292	0.61	-22.03
9049.1088	372070.0	398060.0	585.00	980.183998	980.319562	0.74	-19.74
9049.1089	372260.0	397490.0	599.00	980.183559	980.320024	0.79	-17.84
9049.1090	372360.0	396730.0	624.00	980.178647	980.320639	0.97	-18.26
9049.1091	372270.0	395720.0	688.00	980.162513	980.321452	2.81	-20.78
9049.1092	371950.0	394800.0	565.00	980.194842	980.322190	1.97	-14.23
9049.1093	371780.0	395050.0	651.00	980.173235	980.321985	2.13	-18.55
9049.1094	372330.0	394850.0	620.00	980.181236	980.322155	2.72	-16.23
9049.1095	372320.0	394340.0	618.00	980.182438	980.322566	2.67	-15.88
9049.1096	372380.0	393850.0	604.00	980.187758	980.322962	2.99	-13.39
9049.1097	373510.0	390170.0	160.00	980.286896	980.325947	1.28	-6.30
9049.1098	374140.0	390200.0	252.00	980.269070	980.325932	1.65	-5.64
9049.1099	374580.0	390920.0	245.00	980.268367	980.325357	0.95	-7.85
9049.1100	374520.0	390360.0	283.00	980.259286	980.325808	1.36	-9.50
9049.1101	375060.0	390660.0	294.00	980.261773	980.325574	1.51	-4.46
9049.1102	375890.0	391250.0	381.00	980.241582	980.325109	0.99	-7.59
9049.1103	376580.0	391780.0	433.00	980.230300	980.324691	1.01	-8.20
9049.1104	375430.0	391560.0	245.00	980.269733	980.324853	1.27	-5.66
9049.1105	375760.0	392470.0	225.00	980.271975	980.324123	1.39	-6.50
9049.1106	376600.0	393040.0	402.00	980.234346	980.323675	2.66	-7.59
9049.1107	[redacted]			980.180206	980.310738	0.62	-7.15
9049.1108	[redacted]			980.183029	980.310655	0.59	-6.25
9049.1109	[redacted]			980.183101	980.310648	0.52	-5.84

RGC Exploration - Waratah Area Gravity Data

Station No.	East (m)	North (m)	Elev (m)	Gobs	Gtheo	Terr Corr	B.A. mgal
9049.1110	[redacted]			980.189807	980.310582	0.95	-4.74
9049.1111	[redacted]			980.189562	980.310722	1.10	-4.19
9049.1112	[redacted]			980.188421	980.310547	0.98	-5.67
9049.1113	[redacted]			980.174815	980.310425	3.17	-8.30
9049.1114	[redacted]			980.189858	980.310270	1.73	-5.57
9049.1115	[redacted]			980.181817	980.310090	1.68	-8.56
9049.1116	[redacted]			980.174978	980.310235	2.45	-11.82
9049.1117	[redacted]			980.187004	980.310379	1.79	-11.42
9049.1118	379370.0	403290.0	644.00	980.175477	980.315445	0.17	-13.10
9049.1119	378660.0	402880.0	652.00	980.173673	980.315766	0.21	-13.62
9049.1120	[redacted]			980.177006	980.315656	0.24	-15.26
9049.1121	[redacted]			980.182989	980.315279	0.23	-14.81
9049.1122	[redacted]			980.181439	980.315676	0.22	-14.21
9049.1123	[redacted]			980.174590	980.315304	0.22	-15.96
9049.1124	378410.0	403730.0	652.00	980.171777	980.315078	0.19	-14.84
9049.1125	380240.0	404290.0	641.00	980.175630	980.314651	0.17	-12.75
9049.1126	380700.0	405240.0	647.00	980.173959	980.313891	0.14	-12.51
9049.1127	380970.0	405580.0	652.00	980.173102	980.313620	0.13	-12.12
9049.1128	381360.0	405170.0	660.00	980.172644	980.313956	0.16	-11.31
9049.1129	381340.0	404650.0	654.00	980.174020	980.314375	0.15	-11.54
9049.1130	381560.0	405920.0	663.00	980.170677	980.313354	0.13	-12.11
9049.1131	381690.0	405360.0	663.00	980.172318	980.313807	0.13	-10.93
9049.1132	382030.0	404830.0	660.00	980.173959	980.314239	0.14	-10.30
9049.1133	381920.0	406060.0	661.00	980.171839	980.313246	0.12	-11.25
9049.1134	382300.0	406520.0	649.00	980.173836	980.312880	0.10	-11.27
9049.1135	381740.0	407790.0	642.00	980.175895	980.311848	0.09	-9.56
9049.1136	383770.0	408780.0	636.00	980.175620	980.311077	0.08	-10.26
9049.1137	383520.0	408310.0	628.00	980.177587	980.311452	0.08	-10.24
9049.1138	378790.0	404170.0	657.00	980.171584	980.314728	0.15	-13.74
9049.1139	378980.0	404600.0	644.00	980.172185	980.314384	0.16	-15.35
9049.1140	379000.0	405250.0	634.00	980.175518	980.313860	0.15	-13.47
9049.1141	379390.0	406330.0	622.00	980.178096	980.312994	0.13	-12.40
9049.1142	379500.0	407520.0	623.00	980.179370	980.312036	0.09	-10.01
9049.1143	380710.0	408080.0	638.00	980.176150	980.311601	0.09	-9.85
9049.1144	380070.0	408350.0	638.00	980.176710	980.311375	0.10	-9.05
9049.1145	379370.0	408670.0	621.00	980.179666	980.311107	0.10	-9.17
9049.1146	378480.0	410570.0	618.00	980.179136	980.309563	0.32	-8.53
9049.1147	379700.0	410440.0	631.00	980.176547	980.309685	0.19	-8.81
9049.1148	380480.0	410500.0	629.00	980.178025	980.309647	0.11	-7.77
9049.1149	381580.0	410640.0	639.00	980.176843	980.309549	0.11	-6.89
9049.1150	382330.0	409720.0	632.00	980.177944	980.310300	0.10	-7.92
9049.1151	381850.0	409250.0	633.00	980.177628	980.310673	0.08	-8.44
9049.1152	382980.0	409630.0	625.00	980.178484	980.310381	0.09	-8.85
9049.1153	378060.0	409650.0	605.00	980.181205	980.310299	0.15	-9.92
9049.1154	378190.0	409340.0	606.00	980.181592	980.310551	0.12	-9.62
9049.1155	378240.0	408690.0	605.00	980.181868	980.311076	0.11	-10.08
9049.1156	[redacted]			980.181358	980.311572	0.13	-10.67
9049.1157	[redacted]			980.181144	980.311878	0.11	-11.21
9049.1158	[redacted]			980.181144	980.311530	0.10	-10.68
9049.1159	378800.0	408210.0	609.00	980.180950	980.311471	0.10	-10.61
9049.1160	378770.0	408800.0	608.00	980.181164	980.310994	0.11	-10.11
9049.1161	[redacted]			980.178891	980.308449	1.31	-6.28
9049.1162	[redacted]			980.182255	980.315739	0.16	-13.52
9049.1163	[redacted]			980.183784	980.316204	0.18	-13.22
9049.1164	[redacted]			980.181674	980.316772	0.19	-13.72
9049.1165	[redacted]			980.183366	980.316960	0.22	-13.76
9049.1166	[redacted]			980.181113	980.317161	0.28	-13.80


Station No.	East (m)	North (m)	Elev (m)	Gobs	Gtheo	Terr Corr	B.A. mgal
9049.1167	[redacted]			980.184966	980.317398	0.30	-13.51
9049.1168	[redacted]			980.183896	980.317936	0.41	-15.00
9049.1169	375190.0	399610.0	604.00	980.185037	980.318356	0.78	-13.72
9049.1170	375700.0	399000.0	577.00	980.193252	980.318855	1.15	-10.94
9049.1171	375800.0	398450.0	609.00	980.188115	980.319300	1.43	-9.95
9049.1172	375760.0	397880.0	607.00	980.189297	980.319759	2.10	-8.95
9049.1173	375730.0	397200.0	598.00	980.190653	980.320307	1.75	-10.26
9049.1174	375460.0	396650.0	596.00	980.190622	980.320747	1.99	-10.89
9049.1175	375840.0	393200.0	200.00	980.271516	980.323535	1.41	-11.27
9049.1176	376040.0	393600.0	215.00	980.272046	980.323215	1.72	-7.16
9049.1177	375660.0	393400.0	210.00	980.272739	980.323371	1.92	-7.41
9049.1178	375270.0	394400.0	388.00	980.234795	980.322559	1.88	-9.56
9049.1179	374800.0	394500.0	398.00	980.231900	980.322472	2.23	-10.05
9049.1180	374260.0	394280.0	240.00	980.260937	980.322642	2.73	-11.77
9049.1181	375220.0	395360.0	510.00	980.209957	980.321784	2.63	-8.87
9049.1182	376560.0	393680.0	230.00	980.265207	980.323158	1.33	-11.38
9049.1183	377380.0	394140.0	311.00	980.248309	980.322798	2.15	-11.16
9049.1184	377610.0	394520.0	342.00	980.242082	980.322495	2.40	-10.74
9049.1185	377800.0	395320.0	407.00	980.230076	980.321852	2.18	-9.54
9049.1186	377930.0	395700.0	428.00	980.224959	980.321547	1.87	-10.53
9049.1187	377900.0	396050.0	463.00	980.218365	980.321265	2.03	-9.79
9049.1188	377760.0	396450.0	499.00	980.212821	980.320940	1.59	-8.37
9049.1189	377900.0	397200.0	567.00	980.197043	980.320337	1.32	-10.43
9049.1190	378290.0	397750.0	601.00	980.187544	980.319899	1.21	-12.91
9049.1191	378510.0	398450.0	611.00	980.185190	980.319337	0.77	-13.18
9049.1192	378710.0	398850.0	605.00	980.186637	980.319017	0.56	-12.80
9049.1193	379400.0	399430.0	603.00	980.188798	980.318559	0.42	-10.72
9049.1194	380290.0	403040.0	648.00	980.177200	980.315659	0.18	-10.80
9049.1195	380990.0	402440.0	651.00	980.177872	980.316153	0.20	-10.01
9049.1196	381240.0	401420.0	652.00	980.178504	980.316979	0.31	-9.90
9049.1197	380660.0	400410.0	641.00	980.181990	980.317785	0.33	-9.36
9049.1198	379870.0	399920.0	607.00	980.189522	980.318170	0.35	-8.89
9049.1199	379450.0	406970.0	632.00	980.175854	980.312479	0.09	-12.20
9049.1200	378600.0	406690.0	622.00	980.177220	980.312693	0.09	-13.02
9049.1201	378240.0	406400.0	621.00	980.176710	980.312922	0.12	-13.92
9049.1202	378310.0	406660.0	640.00	980.173062	980.312714	0.13	-13.62
9049.1203	378100.0	406130.0	627.00	980.176170	980.313138	0.13	-13.49
9049.1204	377620.0	406140.0	621.00	980.178219	980.313123	0.15	-12.59
9049.1205	377130.0	406130.0	616.00	980.179564	980.313125	0.14	-12.24
9049.1206	376420.0	406170.0	622.00	980.179075	980.313083	0.13	-11.51
9049.1207	381110.0	406590.0	651.00	980.173541	980.312808	0.11	-11.09
9049.1208	381060.0	407330.0	649.00	980.175049	980.312210	0.10	-9.38
9049.1209	[redacted]			980.181164	980.308794	1.41	-7.79
9049.1210	[redacted]			980.195188	980.308748	1.53	-9.74
9049.1211	[redacted]			980.210497	980.308525	1.39	-10.08
9049.1212	[redacted]			980.222218	980.308308	1.15	-8.03
9049.1213	[redacted]			980.226794	980.308125	1.75	-6.60
9049.1214	[redacted]			980.226233	980.308243	2.02	-5.83
9049.1215	[redacted]			980.227018	980.308375	1.51	-6.08
9049.1216	[redacted]			980.179982	980.308053	1.69	-7.95
9049.1217	[redacted]			980.196045	980.307597	1.29	-6.59
9049.1218	[redacted]			980.214115	980.307188	1.83	-6.46
9049.1219	[redacted]			980.235814	980.306819	1.57	-3.34
9049.1220	[redacted]			980.238148	980.306199	1.26	-2.47
9049.1221	[redacted]			980.242500	980.305640	1.22	0.24
9049.1222	[redacted]			980.228343	980.307171	1.46	-6.95
9049.1223	[redacted]			980.181643	980.311123	0.61	-6.70

RGC Exploration - Waratah Area Gravity Data

Station No.	East (m)	North (m)	Elev (m)	Gobs	Gtheo	Terr Corr	B.A. mgal
9049.1224	[redacted]			980.175885	980.310269	0.73	-5.98
9049.1225	[redacted]			980.189552	980.309662	1.05	-3.98
9049.1226	[redacted]			980.217754	980.309065	2.37	-3.96
9049.1227	[redacted]			980.219578	980.308860	2.45	-4.21
9049.1228	[redacted]			980.223971	980.308795	1.75	-6.75
9049.1229	[redacted]			980.214095	980.308370	1.47	-7.23
9049.1230	[redacted]			980.192916	980.307991	1.42	-7.43
9049.1231	[redacted]			980.176690	980.312178	0.20	-7.81
9049.1232	[redacted]			980.183763	980.311462	0.77	-8.70
9049.1233	[redacted]			980.202353	980.311332	0.92	-6.35
9049.1234	[redacted]			980.222503	980.311157	1.12	-3.93
9049.1235	[redacted]			980.238229	980.311226	0.83	-2.53
9049.1236	[redacted]			980.248309	980.311462	0.88	-0.31
9049.1237	[redacted]			980.248228	980.311495	1.02	1.29
9049.1238	[redacted]			980.242490	980.311461	0.71	5.11
9049.1239	[redacted]			980.252896	980.311502	0.66	8.74
9049.1240	[redacted]			980.272658	980.311944	0.57	18.72
9049.1241	[redacted]			980.234540	980.311393	1.72	16.93
9049.1242	[redacted]			980.221779	980.310803	0.97	15.81
9049.1243	[redacted]			980.251530	980.310252	1.08	20.26
9049.1244	[redacted]			980.250918	980.311899	0.95	0.36
9049.1245	[redacted]			980.236517	980.312358	1.58	-1.87
9049.1246	[redacted]			980.243580	980.312414	1.18	-0.18
9049.1247	[redacted]			980.253161	980.312794	1.15	1.31
9049.1248	[redacted]			980.248982	980.312296	0.86	2.26
9049.1249	[redacted]			980.238209	980.312771	0.94	0.93
9049.1250	[redacted]			980.248411	980.312858	1.83	-0.65
9049.1251	[redacted]			980.248136	980.313566	1.12	-1.36
9049.1252	[redacted]			980.232134	980.314251	1.16	-8.57
9049.1253	[redacted]			980.221321	980.314009	1.42	-5.70
9049.1254	[redacted]			980.218915	980.313014	1.27	-0.57
9049.1255	[redacted]			980.214278	980.313359	1.39	-2.28
9049.1256	[redacted]			980.207531	980.313696	1.85	-2.22
9049.1257	[redacted]			980.251509	980.313329	2.02	-2.56
9049.1258	[redacted]			980.197696	980.311843	1.36	-4.59
9049.1259	[redacted]			980.202486	980.311679	1.67	-5.03
9049.1260	[redacted]			980.213718	980.312017	1.78	-3.08
9049.1261	[redacted]			980.192895	980.307776	1.91	-7.33
9049.1262	[redacted]			980.202466	980.307690	2.19	-5.66
9049.1263	378210.0	411250.0	609.00	980.181878	980.309012	0.70	-6.63
9049.1264	[redacted]			980.186118	980.306038	0.19	-1.10
9049.1265	[redacted]			980.186230	980.305805	1.22	-0.71
9049.1266	[redacted]			980.185068	980.305533	1.41	-0.82
9049.1267	[redacted]			980.192814	980.305780	1.37	-1.04
9049.1268	372960.0	390030.0	160.00	980.290341	980.326052	0.75	-3.49
9049.1269	373180.0	390720.0	175.00	980.286815	980.325499	0.63	-3.63
9049.1270	372690.0	390480.0	225.00	980.273290	980.325685	0.74	-7.40
9049.1271	372080.0	390430.0	220.00	980.274350	980.325717	0.79	-7.30
9049.1272	371610.0	390820.0	311.00	980.256279	980.325395	1.10	-6.84
9049.1273	371580.0	391410.0	373.00	980.236160	980.324919	1.36	-14.03
9049.1274	371670.0	391930.0	387.00	980.236823	980.324501	1.75	-9.80
9049.1275	371620.0	392510.0	402.00	980.230300	980.324032	1.88	-12.77
9049.1276	370870.0	392650.0	401.00	980.225938	980.323908	1.94	-17.15
9049.1277	370640.0	392160.0	320.00	980.245608	980.324300	1.76	-13.99
9049.1278	371700.0	392960.0	351.00	980.238566	980.323670	2.66	-13.40
9049.1279	372050.0	392660.0	393.00	980.234091	980.323917	1.94	-10.58
9049.1280	372230.0	392990.0	527.00	980.204891	980.323654	2.33	-12.76

RGC Exploration - Waratah Area Gravity Data

Station No.	East (m)	North (m)	Elev (m)	Gobs	Gtheo	Terr Corr	B.A. mgal
9049.1281	[redacted]			980.168261	980.312723	0.22	-9.09
9049.1282	[redacted]			980.167476	980.313173	0.26	-10.87
9049.1283	[redacted]			980.166416	980.313867	0.29	-11.81
9049.1284	[redacted]			980.163726	980.314216	0.39	-12.58
9049.1285	[redacted]			980.165815	980.314669	0.43	-13.47
9049.1286	[redacted]			980.171339	980.314939	0.41	-13.94
9049.1287	[redacted]			980.170717	980.315196	0.53	-12.14
9049.1288	[redacted]			980.169678	980.315436	0.66	-14.27
9049.1289	[redacted]			980.152515	980.314419	1.30	-12.26
9049.1290	[redacted]			980.146838	980.314024	1.22	-12.31
9049.1291	[redacted]			980.152749	980.313645	0.91	-10.86
9049.1292	[redacted]			980.159873	980.313553	0.48	-9.98
9049.1293	[redacted]			980.168180	980.313488	0.31	-10.43
9049.1294	388180.0	401790.0	661.00	980.169861	980.316770	0.17	-16.70
9049.1295	387820.0	402030.0	669.00	980.169454	980.316572	0.18	-15.33
9049.1296	387380.0	402280.0	663.00	980.170361	980.316364	0.19	-15.38
9049.1297	386960.0	402410.0	669.00	980.169036	980.316254	0.17	-15.44
9049.1298	386520.0	402220.0	667.00	980.170055	980.316402	0.17	-14.96
9049.1299	386040.0	402020.0	651.00	980.173245	980.316557	0.16	-15.08
9049.1300	385540.0	401670.0	683.00	980.167466	980.316833	0.20	-14.80
9049.1301	385180.0	401560.0	676.00	980.168618	980.316917	0.23	-15.08
9049.1302	384550.0	401360.0	662.00	980.171645	980.317070	0.27	-14.92
9049.1303	383840.0	401350.0	649.00	980.176568	980.317069	0.26	-12.56
9049.1304	383340.0	401490.0	643.00	980.178494	980.316950	0.27	-11.69
9049.1305	383000.0	401810.0	646.00	980.177821	980.316687	0.29	-11.49
9049.1306	382690.0	402170.0	637.00	980.181857	980.316393	0.31	-8.91
9049.1307	382350.0	402280.0	652.00	980.176364	980.316300	0.32	-11.35
9049.1308	381910.0	402030.0	646.00	980.179483	980.316495	0.34	-9.59
9049.1309	380070.0	411470.0	642.00	980.177261	980.308859	0.19	-5.11
9049.1310	379610.0	411860.0	636.00	980.175966	980.308539	0.23	-7.22
9049.1311	[redacted]			980.178198	980.308334	0.27	-6.52
9049.1312	[redacted]			980.180634	980.307948	0.41	-3.95
9049.1313	[redacted]			980.178861	980.307647	1.00	-5.42
9049.1314	[redacted]			980.183733	980.307440	1.08	-4.39
9049.1315	[redacted]			980.196452	980.307225	1.13	-3.02
9049.1316	[redacted]			980.186036	980.306985	1.19	-4.68
9049.1317	[redacted]			980.187147	980.306699	1.61	-2.27
9049.1318	[redacted]			980.182683	980.306379	1.74	-2.35
9049.1319	[redacted]			980.186270	980.315581	0.17	-17.60
9049.1320	[redacted]			980.189685	980.316874	0.28	-14.38
9049.1321	[redacted]			980.192212	980.317025	0.39	-12.29
9049.1322	[redacted]			980.196717	980.317177	0.45	-10.83
9049.1323	[redacted]			980.182214	980.316665	0.20	-11.89
9049.1324	[redacted]			980.181358	980.316537	0.17	-12.05
9049.1325	[redacted]			980.182479	980.316393	0.14	-13.38
9049.1326	[redacted]			980.182204	980.316242	0.13	-12.13
9049.1327	[redacted]			980.182214	980.316114	0.14	-11.59
9049.1328	[redacted]			980.185078	980.315962	0.16	-11.31
9049.1329	[redacted]			980.183254	980.315804	0.21	-11.55
9049.1330	[redacted]			980.187932	980.315693	0.23	-13.63
9049.1331	[redacted]			980.162207	980.313381	0.18	-15.25
9049.1332	[redacted]			980.165723	980.313045	0.22	-15.69



5 cm

083096

LOCATION MAP WARATAH AREA EL 12/90 AND 15/90 FIGURE 1

083097





Figure 84. *Provisional interpretation of form of the Meredith granite. Mineralised sites after Bamford and Green (1986).*


PREVIOUS INTERPRETATION OF MEREDITH GRANITE AND ASSOCIATION WITH MINERALISED SITES (from Leaman & Richardson, 1989a)

FIGURE 2



RESIDUAL BOUGUER ANOMALY WARATAH AREA

FIGURE 3A



Airborne Geophysical Survey and Compilation by GEOEX PTY LTD


RESIDUAL MAGNETIC ANOMALY
1981 Mines Dept Survey

WARATAH AREA
FIGURE 3B

CONTOURS OF RESIDUAL TOTAL MAGNETIC INTENSITY

SCALE 1:50000

083099



Airborne Geophysical Survey and Compilation by GEOEX PTY LTD

for

DEPARTMENT OF MINES TASMANIA


WEST COAST AREA TASMANIA

CONTOURS OF RESIDUAL TOTAL MAGNETIC INTENS

SCALE 1:50000


RESIDUAL GRAVITY + MAGNETIC FIELD QUE RIVER-LUINA AREA

FIGURE 3C




TRENDS WARATAH REGION

FIGURE 4



LOCATION OF PROFILES REFERRED TO IN TEXT

FIGURE 5



POSSIBLE ALTERNATE SECTION : HUSKISSON – QUE/HELLYER MACKINTOSH REGION


Note: Solution consistent with gravity/magnetic data other structural configurations (incl drusted sequences) not tested.

D. E. Leaman
Jan 91


INTEGRATED GEOLOGICAL SECTION FOR LINE 7
FIGURE 10C

WARATAH AREA


083102



WARATAH AREA CURRENT MODEL OF FORM OF MEREDITH GRANITE
Mineralised sites indicated FIGURE 26



WARATAH AREA GRANITE MODEL AND TREND SUMMARY FIGURE 27



WARATAH AREA GRANITE MODEL AND TARGET ZONES FIGURE 28

APPENDIX 3

Rock Chip Geochemistry

RGC EXPLORATION PTY.LTD. DATA SHEET

27/ 5/92

PROJECT: Waratah rock chip geochemistry.Meredith Granite surrounds.

SAMPLE NUMBER	TNORTH metres	TEAST metres	AS PPM	CU PPM	PB PPM	SN PPM	W PPM	ZN PPM	AG PPM
t 20090	394640	375100	50	95	715	-3		1300	-0.5
t 20091	394110	375070	-2	5	5	-3		5	-0.5
t 20092	394080	375120	-2	5	5	-3		10	-0.5

Laboratory:	ANALAB	ANALAB	ANALAB	ANALAB	ANALAB	ANALAB	ANALAB
Method :	GA114	GA101	GA101	GX401	GX401	GA101	GA101
Det. Limit:	1.000	5.000	5.000	3.000	1.000	5.000	0.500

20140

083108

5 cm

92-3357.1

RGC EXPLORATION PTY. LIMITED
Incorporated in New South Wales

COMPILED:		MEREDITH PROJECT
DRAWN:	M. WALTER	E.L. 12/90 & E.L. 15/90
DATE:	JUNE 1990	AREA TO BE RELINQUISHED
CHECKED:		
1:250 000 REFERENCE:		
BASE PLAN NO.	5530/039	
OVERLAY PLAN NO.	-	
SCALE 1:50 000	0 1 2 3 4km	PLAN 1

5400000 m N

5390000 m N

370000 m E

378000 m E

Ramsay River

Creek

Celistream River

Hay

CROWN LAND

COLDSTREAM RIVER

MOUNT RAMSAY

RIVER

RAMSAY

HØSKISSON

Webbs Creek

Ross

medium forest with low dense vegetation

forestry operations

DRIVE

1950

T 2009

T 2009a

scrub

talus

STRATIGRAPHY

QUATERNARY		Qa	alluvium
TERTIARY		Tb	basalt
		Ts	sediment
DEVONIAN		Dg	Meredith granite
SILURIAN	ELDON Gp.	Sa	Amber shale
		Sc	Crotty sandstone
ORDOVICIAN		Og	Gordon limestone
CAMBRIAN	DUNDAS Gp	Edc	conglomerate
		Edg	greywacke
		Eds	siltstone
	CRIMSON CREEK Fm.	Ecg	greywacke
		Ecs	siltstone
		Ecb	basalt
PRECAMBRIAN	OONAH Fm.	Pog	quartzite
		Posh	shale
		Pos	siltstone
		Posd	sandstone
		Pod	dolomite

strike and dip of bedding

strike and dip of foliation


strike and dip of jointing

anticline

syncline

fault

F vein



LOCALITY PLAN
SCALE 1:500 000

siltstone (s)
conglomerate (c)
sandstone (sd)
greywacke (g)
shale (sh)
basalt (b)
granite (g)

Q Quaternary
Ed Dundas Group
Ec Crimson Creek Formation
Po Oonah Formation
T Tertiary

5 cm

92-3357.

RGC EXPLORATION PTY. LIMITED
Incorporated in New South Wales

083109

COMPILED:
DRAWN:
DATE:
CHECKED:
1:250 000 REFERENCE:

WARATAH AREA. E.L. 12/90 & 15/90
FACTUAL GEOLOGY

Sheet 8 of 11

BASE PLAN NO. 5530/029
OVERLAY PLAN NO.

SCALE 1:10 000

PLAN 2

5 390 000 m N

WARATAH 84 EAST

CROWN LAND

CROWN LAND

MOUNT KERSHAW

5380 000 m N


370 000 m E

379 000 m E

STRATIGRAPHY

DEVONIAN	Dg	Meredith Granite
	Qp	Quartz-feldspar porphyry
SILURIAN ELDON Gp.	Sa	Amber shale
	Sc	Crotty quartzite
Ordovician	Og	Gordon limestone
CAMBRIAN CRIMSON CREEK Fm.	Ecg	greywacke
	Ecs	siltstone
	Ecb	basalt
PRECAMBRIAN Oonah Fm.	Poq	quartzite
	Posh	shale
	Pos	siltstone
	Pod	dolomite
	Posd	sandstone

- strike and dip of bedding
- strike and dip of foliation
- strike and dip of jointing
- fault
- anticline
- syncline



LOCALITY PLAN
SCALE 1:500 000

5 cm

92-3357.

RGC EXPLORATION PTY. LIMITED 083110
Incorporated in New South Wales

COMPILED:	
DRAWN:	
DATE:	
CHECKED:	
1:250 000 REFERENCE:	


WARATAH AREA. E.L. 12/90 & 15/90
FACTUAL GEOLOGY

Sheet II of II

BASE PLAN NO. 5530/032
OVERLAY PLAN NO.

SCALE 1:10 000

PLAN 3



RGC EXPLORATION

Compilation by D.E. Leaman, Leaman Geophysics, Hobart Jan 1991
Data sources : Division of Mines, RGC
Reduction density : 2.67 t/m³
1930 ellipsoid
22km terrain correction


083111

WARATAH AREA
BOUGUER ANOMALY

PLAN 4 MAP 1

92-3357.

5530/017



083112

5530/018

RGC EXPLORATION

Regional - residual separation using MANTLE 2 process
Bouguer anomalies at 2.67 t/m³ reduction density
Compilation by D.E. Leaman, Leaman Geophysics, Hobart
Jan 1991

92-3357. WARATAH AREA
RESIDUAL ANOMALY

PLAN 5

MAP 2

5420 000 mN

380 000 mE

390 000 mE

Relationship of Burnie Fm and granite not resolved in detail at depth.

5410 000mN

9000

Granite limit not resolved due to complex anomaly patterns in Cambrian rocks near surface

5400 000 mN

5400 000 mN

360 000 mE

Meredith Granite

400

6500

200

750

1500

2500

4500

6500

9000

Sites of "granite exposures" after Corbett & McNeill (1988) Conclusion: Not in place/or part of Meredith pluton

Approximate position of granite exposure boundary based on large scale regional mapping (see text)

5390 000 mN

370000 mE

380 000mE

390 000mE

92-3357.

083113

Interpretation incorporates data sampling, density assumptions as explained in the associated text.

5530/019

RGC EXPLORATION

Interpretation by D.E. Leaman
Leaman Geophysics
Jan. 1991

Contours in metres of estimated depth to granite – below surface

WARATAH AREA

FORM OF MEREDITH GRANITE

PLAN 6

MAP 3