

TCR 98-4116

# Aberfoyle Resources Limited

655001

EXPLORATION DIVISION

ACN 004 664 108

MINERAL RESOURCES		
FILE REF: EL 106/87		
29 JAN 1998		
DOC. REF.		
OFFICER	FOR ACTION	FOR INFO
See Folio 65		
RESUBMIT TO	DATE	

EXPLORATION LICENCE 106/87

LAKE MACKINTOSH

**TASMANIA**  
**MICROFILMED**  
 FICHE No. 014439-41

Relinquishment Report

February 1998

VOLUME 1 OF 1

Prepared by:

Andrew McNeill  
Senior Geologist

Richard de Bomford  
Services Officer

Steven Richardson  
Senior Geologist

Issued by:

A M Hespe  
MANAGER - BASE METALS

DISTRIBUTION

- Aberfoyle - Burnie (1/4)
- Aberfoyle - Townsville (2/4)
- Aberfoyle - Melbourne (3/4)
- Mineral Resources Tas (4/4)

98\_4116



655002

## LIST OF APPENDICES

I	EL 2/70 and 15/73 - Report references.
II	EL 106/87 - Details of untested structural / stratigraphic targets
III	MAC 43 - Drill log and logging codes
IV	MAC 43 - Assays
V	MAC 43 - Pb isotope results
VI	MAC 43 - Downhole EM survey results
VII	MAC 43 - Magnetic susceptibility results

## LIST OF FIGURES

		Follows page
1	Original Licence Boundary	2
2	Area of First Relinquishment	2
3	Area of Second Relinquishment	2
4	Current Tenure	2
5	Untested Structural / Stratigraphic Targets on EL 106/87	3
6	Prospect Location Guide	4
7	Stratigraphic Diagram	7
8	Drill Hole Location Guide	18

## LIST OF PLATES

<u>No.</u>	<u>Title</u>	<u>Scale</u>
MAC 478	South Mackintosh / Mt. Charter Area - Rehabilitation Plan.	1:10000
MAC 481A	Cross Section 14500N - DDH MAC 43	1:1000
MAC 481B	Cross Section 14500N - DDH MAC 43	1:1000

**TABLE OF CONTENTS**

	Page No.
1.0 SUMMARY	1
2.0 INTRODUCTION	2
3.0 EXPLORATION PHILOSOPHY	3
4.0 PREVIOUS EXPLORATION	4
4.1 Introduction	4
4.2 Geology	4
4.2.1 Mineral Resources Tasmania mapping	4
4.2.2 Aberfoyle Resources mapping	4
4.2.3 Geological interpretation	7
4.2.4 Mineralisation and alteration	8
4.3 Geochemistry	9
4.3.1 Stream sediments	9
4.3.2 Soils	10
4.3.3 Drill core	12
4.3.4 Rock-chip / channel sampling	13
4.3.5 Pb-isotopes	13
4.4 Geophysics	13
4.4.1 Ground-EM	13
4.4.2 Seismic	15
4.4.3 Magnetics	16
4.4.4 Radiometrics	17
4.4.5 Gravity	17
4.5 Diamond Drilling	18
4.6 Research	22
4.6.1 Theses	22
4.6.2 Consultants reports	23
4.7 Access	24
4.7.1 Line cutting	24
4.7.2 Vehicular tracks/costeans	24

	Page No.
5. WORK COMPLETED IN THE FINAL YEAR	25
5.1 Introduction	25
5.2 DDH MAC 43 - Geology	25
5.3 DDH MAC 43 - Geochemistry	27
5.4 DDH MAC 43 - Geophysics	28
5.4.1 Downhole EM	28
5.4.2 Magnetic susceptibility	29
6. ENVIRONMENTAL DISTURBANCE AND REHABILITATION	30
6.1 Introduction	30
6.2 To be left open	30
6.3 To be rehabilitated	30
7. CONCLUSIONS	32
8. EXPENDITURE	33
9. REFERENCES	34

## **1.0 SUMMARY**

This report summarises the results of exploration for volcanic-hosted base metal and gold mineralisation on EL 106/87, Lake Mackintosh, for the period February 1988 to February 1998. Results of some earlier exploration, from the period 1970-1988, are also included as background to the more recent exploration.

The majority of EL 106/87 is underlain by the Que-Hellyer Volcanics (QHV), which host the Que River (discovered in 1975), Hellyer (discovered in 1983) base metal deposits. The presence of this mineralisation has led to sustained and extensive exploration that has included:

- Nineteen diamond drill holes, for 13,460m of drilling.
- Heli-borne magnetics and radiometrics.
- Experimental seismic traverses.
- Research projects into the structure, stratigraphy and geochemistry of the Que-Hellyer Volcanics.
- Small scale frequency-domain EM surveys.
- A total of 244 line km of track cutting to allow access for:
  - \* Reading of 149.6 line km of ground time-domain EM.
  - \* Collection of 4057 'b' and 'c' horizon soil samples for analysis by total digest and 1031 samples for analysis by partial digest techniques.
  - \* Collection of 254 stream sediment samples (-80# and BCL).
  - \* Collection of 48 line km of ground magnetic data.
  - \* 1:2500 scale geological mapping and rock-chip sampling (218 samples).
  - \* Collection of gravity data from 530 infill stations.

During the final year of EL 106/87 work focussed on diamond drill testing at Mayday which comprises a structural / stratigraphic target associated with a partial digest soil anomaly. The targeted Que Hellyer Volcanics were not intersected and the hole was terminated in the Southwell Subgroup at 1504.8m. No source for the soil geochemical anomaly is evident.

## **2.0 INTRODUCTION**

The Lake Mackintosh Exploration Licence (EL106/87) was granted to Aberfoyle Resources Limited on 5 February 1988 under the provisions of the Hellyer Mine Agreement Ratification Act 1987. The licence comprised 135 sq. km. previously covered by EL's 2/70 (Mackintosh) and 15/73 (Hatfield) and encloses CML's 68M/84 (Que River) and 103M/87 (Hellyer) which have an area of 20.2 sq. km; Figure 1.

Under the terms of the act the licence was issued for ten years with mandatory partial relinquishments on the second and fifth anniversaries. The first partial relinquishment was effected in February 1990 when the licence was reduced from 135 to 95 sq. km. (McNeill 1990, TCR 90-3073; Fig. 2). The second partial relinquishment, from 95 to 54 sq. km, was effected in February 1992 (Wallace 1992, TCR 93-3410; Fig. 3). Current tenure and the location of a proposed retention licence over the Mt. Charter mineralisation are shown on Figure 4.

This report details exploration completed in the twelve months to 5 February 1998 and summarises previous exploration by referencing the relevant reports from the previous nine years.

As EL 2/70 (Mackintosh), granted on 1 January 1970, and EL 15/73 (Hatfield), granted on 5 May 1973, were never relinquished, as such, a comprehensive summary of exploration completed on these licences prior to 5 February 1988 has not been prepared. However, exploration of the "Mackintosh District" (EL's 2/70, 15/73, and 5/74) for the period 1969-December 1985 is summarised by Anon (1986; TCR 86-2521) and a list of reports relating to the two licences that were combined to form EL 106/87 are presented in Appendix I. Additionally some results from EL 15/73 and 2/70 are summarised in this report as background to exploration completed during the life of EL 106/87.




Figure 1

**Aberfoyle Resources Limited**  
EXPLORATION DIVISION

NORTH WEST TASMANIA

MACKINTOSH E.L.106/87

**ORIGINAL LICENCE BOUNDARY**

Compiled : AMcN

Drawn :

Traced : JLR

Checked : A McN

Plate No. : MAC 181

REVISIONS			
Init.	Date	Init.	Date

Location Code : K55/6/44

Scale : As shown

Date : February, 1988

1-5-1




Figure 2

**Aberfoyle Resources Limited**  
EXPLORATION DIVISION

NORTH WEST TASMANIA

MACKINTOSH E.L.106/87

**AREA OF FIRST RELINQUISHMENT**

Compiled : AMcN

Drawn :

Traced : RJE

Checked :

Plate No. : MAC 274

REVISIONS			
Init.	Date	Init.	Date

Location Code : K55/6/44

Scale : As shown

Date : December, 1989

1  
5  
1




Figure 3

**Aberfoyle Resources Limited**  
EXPLORATION DIVISION

NORTH WEST TASMANIA  
MACKINTOSH E.L.106/87

**AREA OF SECOND RELINQUISHMENT**

Compiled :	AMcN
Drawn :	
Traced :	JMS
Checked :	
Plate No. :	MAC 352

REVISIONS			
Init.	Date	Init.	Date
RJE	May 90		
JMS	6-4-93		

1511




Figure 4

ABERFOYLE RESOURCES LIMITED EXPLORATION DIVISION	
Date 23/5/1997 Author RdeB Office BURNE Plate MACA75 Scale 1:100000	North Western Tasmania MACKINTOSH EL106/87 AREA OF CURRENT TENURE
Projection: AMG Zone 55 (AGD 66)	


### **3.0 EXPLORATION PHILOSOPHY**

Exploration by Aberfoyle Resources Ltd. on EL 106/87 has been aimed at the discovery of a >10 m.t. polymetallic VHMS deposit to replace the Hellyer resource when it is exhausted in mid-2000. A secondary target has been volcanic-hosted or structurally related gold mineralisation.

After the discovery of Hellyer, in 1983, exploration was largely led by the application of ground TDEM techniques for direct target definition to depths of 200-300m. However, by 1988 TDEM surveys had largely covered the outcropping QHV and Que River Shale (QRS) and the only targets defined by these surveys were down-graded as artefacts or false anomalies (lithological conductors).

Exploration recorded in this report was largely designed to test deep targets (>300m and <1500m depth) in the QHV or shallower targets in the less well explored dominantly sedimentary and felsic volcanic sequences (the Southwell Subgroup) or Tertiary basalt covered areas north and east of Hellyer.

Since 1993 in particular, target generation has focussed on defining deep targets through integrating geological, geophysical and geochemical data and developing a three dimensional structural model of the QHV basin. This approach was based upon recognition of the close association of both Hellyer and Que River orebodies with district scale structures and the more general observation that VHMS deposits are always associated with co-active faults. Targets generated by this process and tested (usually by diamond drilling) have been described in the relevant annual reports. Several targets remain untested on both EL 106/87 and ML 103M/87. Those remaining on the EL are shown on Figure 5 and summarised in Appendix II.



**MACKINTOSH EL 106/87  
REMAINING IDENTIFIED  
STRUCTURAL /  
STRATIGRAPHIC  
TARGET AREA  
LOCATIONS**

☐ Identified  
Target Area  
(Targets on ML's not shown)





Figure 5



## **4.0 PREVIOUS EXPLORATION**

### **4.1 Introduction**

Exploration completed at each prospect on EL 106/87 is summarised on Table 1 (see Figure 6 for prospect locations). Note that the "Regional" prospect generally refers to data collection and compilation over the outcropping QHV on EL 106/87, and may include some data from the Hellyer and Que River CML's. Background information, including details of surveys completed on EL's 2/70 and 15/73, and a more detailed summary of results, divided by work type, is included in the following sections.

### **4.2 Geology**

Several generations of geological mapping has been completed over EL 106/87 by both Mineral Resources Tasmania and Aberfoyle Resources Ltd.

#### **4.2.1 Mineral Resources Tasmania**

Three generations of regional mapping have covered the area of EL106/87:

1. The Mackintosh 1: 63,360 sheet (Barton et al., 1966); little detail shown in volcanics on this sheet.
2. More detailed, 1: 36, 680, mapping by P.L.F. Collins that was incorporated into the explanatory notes for the Mackintosh sheet (Collins et al., 1981).
3. Mapping, compiled at 1:25,000, as part of the Mount Read Volcanics Project. Map sheets 1 (Komyshan, 1986), 2 (Corbett and McNeill, 1986), and 7 (Vicary and Pemberton, 1988) cover the area of EL 106/87. Explanatory notes for these map sheets have also been published as Corbett and Komyshan (1989), McNeill and Corbett (1989), and Pemberton et al. (1991).

#### **4.2.2 Aberfoyle Resources Mapping**

##### *Outcrop maps*

Outcrop mapping has been completed at scales of 1:1,000 (detailed costean mapping), 1:2,500 (standard outcrop mapping) and 1:10,000 (reconnaissance mapping). Prior to 1983 outcrop mapping was compiled on the 1:2,500 QR81 series of plans. However, with the discovery of the Hellyer deposit, and revisions in geological concepts and terminology, a new geological legend (plate MAC114) and map series (MAC89 plans; see plate MAC355 for plan locations) was commenced (Hespe, 1986a). The pre-1983 mapping was included on the MAC89 series



TABLE 1: SUMMARY OF WORK COMPLETED

REPORTING PERIOD/REF	PROSPECT	DETAILS
Feb. 1988- Feb. 1989. McNeill TCR 89-2948	Southwell River / Leven River	Mapping, stream sediment sampling (108), rock chip sampling (54), VLF EM (2 lines), SIROTEM & EM37 depth sounding, drilling (MAC20, MAC22), down hole EM37.
	North Hatfield	Mapping and rock chip sampling (8).
	Henty Fault Zone	Mapping, rock chip sampling (24), stream sediment sampling (59 BCL & -80#).
	South Hatfield	Mapping, BCL & -80# stream sediment sampling (55), infill soil sampling (92), rock chip sampling (15), infill ground magnetics, drilling (MAC16) & down hole SIROTEM.
	I Zone	Gridding (20 km), mapping, infill soil sampling (652), rock chip sampling (21), follow up UTEM (3.5 km).
	Mt. Charter	Mapping, reassy of DDH's MC1-4, rock chip sampling (41), infill soil sampling.
Feb. 1989- Apr. 1990. McNeill, Rand, Henham  TCR90-3128	Regional	Compilation of regional cross sections and level plans, lineament study, gridding (67.7 km covering EL106/87 and CML103M/87), C horizon soil sampling (2011 over new gridding), compilation and processing of all soil data, reassy of 522 core grinds from 26 holes for lithological discriminator elements.
	Baryte Creek	Mapping, gridding (6 km), follow up EM survey (MAX-MIN 5.1 line km) over 2 airborne EM anomalies
	Tailings Dam	Mapping
	Golden Triangle	Mapping, Rock chip sampling
	N.W. Mt. Charter	Costean channel sampling (19)
	Charter Dolerite	Gridding (3.8 km), UTEM survey (3.1 line km)
	Medway	Gridding (23.6 km), Mapping, UTEM survey (4 loops, 30.4 line km).
	Link Road	Channel sampling, Gridding (4 km), UTEM (2 lines of the north Hellyer survey on CML 103M/87)
	Southwell Valley	EM (MAX-MIN 3 lines) over airborne EM anomaly 4/75.
	Leven River	DHEM (EM37) completed on MAC22.
	North Hatfield	Mapping (50 sq km).
	Black Harry Road	Mapping, costeaning (2) and subsequent channel sampling. Drilling by DOM DDH BLHY-1 (MAC24), down hole EM37.
	I Zone	Mapping, follow up UTEM (1 loop, 3.8 line km)
	Mt. Charter	Mapping, costeaning and subsequent sampling, reprocessing soil data, rock chip and pit sampling, drilling (MAC23), down hole EM37.
	S.W. Mt. Charter	Drilling (MAC25), down hole EM (SIROTEM).
	Henty Fault Zone	Gridding (7.2 line km), mapping, rock chip sampling (7), reassy of earlier soils for Au, stream sediment sampling (32 BCL, -72# & -200#).
	South Mackintosh	Mapping, rock chip sampling (51), ground magnetics (10 line km)
May 1990- Apr. 1991. McNeill, Wallace TCR91-3268	Regional	Reprocessing of all soil geochemistry & expansion of lineament study to cover the imaged data.
	Southwell Valley	Gridding (14.05 km), Mapping, rock chip sampling (12), UTEM survey (1 loop, 3.6 line km), ground magnetics (13.1 line km).
	I Zone / South Que River	Mapping, rock chip sampling (21), drilling (MAC28).

REPORTING PERIOD/REF	PROSPECT	DETAILS
May 1990- Apr. 1991. (cont.)	Mt. Charter	Drilling (MAC26), down hole EM (Zonge GDP-16)
	Mt. Charter Dolerite	Drilling (MAC27), down hole EM (Zonge GDP-16)
	South Mackintosh	Mapping, review of UTEM data
Apr. 1991- Apr. 1992. Richardson TCR92-3355	Regional	Structural/stratigraphic reinterpretation of Que-Hellyer volcanics commenced by Etheridge & Henley.
	South Que River	Drilling (MAC29, MAC30), down hole EM (MAC28, MAC29).
Apr. 1992- Apr. 1993. Richardson TCR93-3441	Regional	Completion of Etheridge & Henley structural project, reinterpretation of all existing 1:2500 scale mapping, helicopter borne aeromagnetic and radiometric survey.
	South Que River	Down hole EM (MAC30), reinterpretation of all DHEM data for this area.
	Mt. Charter	Geological reinterpretation, drilling (MAC32, MAC33).
	Murchison Highway	Drilling (MAC34, MAC35).
Apr. 1993- Feb. 1994. Richardson. TCR94-3537	Regional	Interpretation of previous years helimag data, infill gravity survey (530 stations).
	South Que River	Resurvey MAC28 with down hole UTEM.
	Mt. Charter	Down hole UTEM (MAC33).
	Murchison Highway	Down hole UTEM (MAC35).
	South Mt. Charter	Drilling (MAC36).
	Baryte Creek	Drilling (MAC37).
Feb. 1994- Feb. 1995. Richardson TCR95-3719	Regional	Gridding commenced for EM and soil programmes north of Hellyer (70 line km to date).
	South Mt. Charter	Down hole EM (MAC36).
	Baryte Creek	Down hole EM (MAC37).
Feb. 1995- Feb. 1996 Richardson TCR96-3839	Regional	Complete gridding (124 km total), Zonge GDP-16 TDEM survey (20 loop, 103 line km) north of Hellyer.
	Mt. Charter	Drilling (MAC40), down hole EM (Zonge & Crone).
	Leven River	Ground EM survey, drilling (MAC41), down hole EM.
Feb. 1996- Feb. 1997. McNeill, Hicks TCR97-3991	Regional	Update of Etheridge & Henley structural study, mag sus readings on selected holes, AGSO seismic traverse north of Hellyer, TMI soil sampling programme (1031) & EM depth soundings north of Hellyer.
	Murrays Road	Soil sampling (271).
	Link Road	Gridding (2.75 km), Wacker soil sampling (87).
Feb. 1997- Feb. 1998 This report	Mayday (Southwell River)	Drilling (MAC43), down hole EM.

plans where no more recent mapping was available and is distinguished by italic lettering. 1:10,000 scale outcrop mapping is recorded on the MAC193 series of plans and covers areas peripheral to the QHV where gridding was not undertaken. Outcrop plans were continuously updated during the life of the licence. Outcrop descriptions are based on field descriptions which may be supported by analyses of rock-chip samples (see section 4.3.4) and petrographic description of thin sections (some 350 of which have been prepared; locations are available digitally).

#### *Interpretive maps*

Interpretive plans and cross sections have been compiled from all available data sources; outcrop mapping, airphoto interpretation, drilling, rock-chip and soil geochemistry, open file reports on adjacent tenements and magnetic and gravity surveys. Plans have been produced at scales of 1:2,500 (MAC148 plan series; now out of date), 1:10,000 (MAC161 series in McNeill and Wallace, 1991; plate MAC380 in Richardson, 1993) and 1:25,000 (plate MAC412 in Richardson, 1994). Regional cross sections (at 1:10,000 scale) were presented in McNeill (1989b; plates MAC97A-D) and were updated in McNeill et al. (1990; plates MAC269A-D). Interpretive level plans (plates MAC284 and 285), largely derived from the regional cross sections were also included in McNeill et al. (1990).


#### 4.2.3 Geological interpretation

The Current geological understanding of EL 106/87 is summarised on plate MAC 412 (Richardson, 1994) and on Figure 7. In the northern and eastern parts of the EL outcropping lithologies are the Tertiary flood basalts and Cambro-Ordovician to Devonian clastic and carbonate sediments that overly prospective Middle-Late Cambrian Mount Read Volcanics. The volcanics in this area comprise the Central Volcanic Complex overlain by the Mt. Charter Group (as defined by Corbett, 1992; Figure 5). The economically significant unit of the Mt Charter Group is the Que-Hellyer Volcanics (QHV) which host the Hellyer, Que River and Mt. Charter VHMS bodies. Published descriptions of the volcanology, stratigraphy and primary litho-geochemistry of the QHV include Waters and Wallace (1992), Corbett (1992) and Crawford et al (1992).

The QHV are a sequence of marine calc - alkaline mafic to felsic volcanics and volcanoclastics deposited / erupted into an extensional basin interpreted to develop as a result of movement on regional synvolcanic faults such as the Henty, Mt. Charter and Mt. Cripps Faults. The QHV can be broadly subdivided into a lower sequence of basalt and feldspar phyric andesite lava


Figure 7

# MACKINTOSH AREA STRATIGRAPHIC CORRELATION DIAGRAM



SOUTH

NORTH



and volcanoclastic (geochemical suite 1 of Crawford et al. (1992)) separated from an upper sequence of dominantly basaltic rocks (geochemical suite 3) by a complex interval known as the Mixed Sequence. The Mixed Sequence marks a relatively quiescent period dominated by polymict epiclastics and numerous small volume dacitic lava / breccia bodies. Thickness of the Mixed Sequence varies from a few centimetres to more than 300m whilst the total thickness of the QHV can vary from around 20m (in the NW) to more than 1.5 kilometres in the vicinity of the orebodies.

A strong Cambrian structural control is recognised for the formation and morphology of the QHV basin and associated VHMS mineralisation. A network of syndepositional NE (mine grid) trending normal faults linked by NW trending transfer faults has been interpreted from facies and thickness variations, magnetics, gravity etc. In addition, a major NNE half graben like structure extends at least from Que River to Hellyer, hosting the two orebodies and localising strong footwall alteration. This structural zone is interpreted to relate to oblique extensional reactivation of a deep tapping, basement structure.

Compressional structures are thought to largely reflect reactivation of these syndepositional structures during the ?Late Cambrian, Devonian and Mesozoic events.

Southwell Subgroup (SSG) rocks outcrop or lie beneath Tertiary basalt over much of the northern half of EL 106/87. A distinctive quartz phyric volcanoclastic near the base of the SSG has been correlated by McPhie and Allen (1992) with hangingwall volcanoclastics overlying the Rosebery orebody. One interpretation of this correlation is that Rosebery mineralisation may be younger than Hellyer and that potential exists for mineralisation near the base of the SSG north of Hellyer.

#### 4.2.4 Mineralisation / alteration

Although there is evidence of base metal mineralisation higher and lower in the QHV, at present all known major occurrences occur within the Mixed Sequence. VHMS mineralisation occurs where hydrothermal fluids have been focussed into dilatant areas near the intersection of synvolcanic faults, possibly during a regional stress pulse at Mixed Sequence time (Windh and Etheridge, 1992).

The three stratiform and stratabound VHMS bodies discovered to date are (from north to south):

Hellyer: Mineralisation is described by McArthur and Dronseika (1990) and McArthur (1996). Hydrothermal alteration of the footwall and hangingwall are described by Gemmell and Large (1992) and Jack (1989) respectively. The structure of the orebody has been discussed by Drown and Downs (1990) and Downs (1993).

Que River: Mineralisation is described in McArthur and Dronseika (1990) and McGoldrick and Large (1992). Structure is described in Large et al (1988) and Young (1980) and alteration in Whitford et al (1989) and Offler and Whitford (1992).

Mt Charter: The geology of the Mt. Charter mineralisation is described in Rand (1988).

### **4.3 Geochemistry**

#### **4.3.1 Stream sediments**

Two generations of stream sediment sampling have been completed over the area of EL 106/87:

##### *1. Pre-1983.*

An initial regional -20# stream sediment sampling program covering EL 2/70 (Krummei, 1972) located an anomaly that was later found to be adjacent to AEM anomaly 8, leading to the discovery of the Que River deposit (Webster and Skey, 1979). An orientation survey, covering three streams draining the Que River deposit, was completed (ibid) and the -40# fraction selected as being a suitable medium for an expanded survey which covered the area of outcropping QHV's on EL's 2/70 and 15/73 (Skey, 1975; Skey and Webster, 1976a and 1976b). Results are shown on plates DT60A and B (Skey and Webster 1976a). Some stream humus and water samples were also collected (ibid).

##### *2. Post-1988.*

Stream sediment sampling was as a major component, in conjunction with mapping and rock-chip sampling, of initial base metal and Au exploration in poorly-explored areas outside the QHV. The felsic volcanics and sediments north and east of Hellyer (Southwell River prospect), the Henty Fault Zone, and the Central Volcanic Complex (South Hatfield prospect) were all sampled (McNeill, 1989; McNeill et al., 1990). Sample locations and results are shown on plates MAC203 and 204 (in McNeill et al., 1990). Sample media were the -72#, -200#, and -7mm (used for bulk cyanide leach analysis) size fractions, although not all fractions were analysed at each site.

#### 4.3.2 Soils

##### *Total digest*

A total of 15, 577 nominal 'C' horizon soil samples were collected on EL's 15/73, 2/70 and 106/87. As these samples were collected over a long time period a range of collection and analytical procedures were used:

- Sample collection:** The majority of samples were nominal "C" horizon collected using a hand auger from < 1.0m depth (but up to 1.5m in some surveys). "A" horizon samples (e.g., the Hg orientation survey described in McNeill, 1989a) and "B" horizon samples (collected in conjunction with partial digest samples; see below and McNeill and Hicks, 1997) were collected as part of some surveys. Sampling density was variable, with sample spacings of 10m to 50m (but generally 25m), on 50m to 500m spaced lines.

- Analysis:** Prior to analysis the majority of soil samples were dried and sieved to -80#, exceptions being some early (pre 1975; e.g., Skey, 1975) samples which were sieved to -40#, and samples from the partial digest sampling (McNeill and Hicks, 1997) which were sieved to -6mm prior to analysis. Pre- 1985 soils were analysed for Cu, Pb, and Zn with scattered Fe, Ba, and As results. Post -1985 the majority of samples were analysed for Cu, Pb, Zn, Ba, As, Ag, Cr, Ti, and Zr. The latter three elements were analysed as they are relatively immobile and were used to differentiate basalts, andesites and dacites in areas of poor outcrop (Hespe, 1986; Ni was also used in some surveys, but as it provides no more information than Cr it was not analysed in more recent surveys). Ba, As, and Ag were analysed as pathfinder elements, as they were cheap to assay, gave good anomaly contrast and are relatively immobile in the secondary environment. Gold assays were done on specific prospects, i.e., Mt. Charter and South Hatfield (McNeill, 1989c), Boundary (McNeill, 1989b), Henty Fault Zone (McNeill et al, 1990) and Murrays Road (McNeill and Hicks, 1997). A mercury and antimony orientation survey was also completed (Hespe in McNeill, 1989b) and some samples were analysed for mercury in the Tailings dam area (McNeill, 1989b).

Problems with quality control in some surveys led to the inclusion of at least one standard sample (prepared by homogenising 20kg of "C" horizon soil material from EL's 15/73 and 2/70; see Appendix 4 of McNeill, 1989a for composition) with every batch of soil samples submitted for analysis since early 1986.

•**Results:** Data indicates widespread dispersion of base metals throughout the QHV. Zones anomalous in base metals occur throughout the outcropping QHV, both areally and stratigraphically. They indicate hydrothermal activity was continuous throughout development of the basin, continuously pervading superposed volcanic sequences as subsidence of the basin progressed. Highest values are found within the footwall alteration zones outcropping between the Que River and Hellyer mines. In contrast footwall alteration exposed in the sub parallel Amoeba zone is almost barren.

In broad terms:

Copper is elevated within outcropping footwall alteration but also defines outcropping basalt.

Lead anomalies define -

- footwall alteration between Hellyer and Que River and at Mt. Charter
- Mixed Sequence containing clastic base metal sulphides
- evolved Upper basalt containing veins and disseminations of sphalerite and galena  
E and NE of Mt Charter
- weakly mineralised footwall epiclastic breccias on top of the Lower basalt, 1.5 km south of Que River mine
- Upper basalt overlying a thin intersection of base metal sulphides at the Hellyer ore position, 500m south of Mt. Charter
- a poorly understood zone within Upper basalt in the core of a syncline north of the Que Fault but not associated with any visible mineralisation.

Zinc anomalies reflect the same features as lead but are often broader due to greater mobility.

Barium anomalies indicate hydrothermal alteration; particularly in the Mt. Charter area.

Arsenic also has a strong correlation with known areas of hydrothermal alteration whilst silver is only consistently anomalous in soils at Mt. Charter.

#### *Partial Digest*

Partial digest, or transported metal ion (TMI), soil sampling has been shown to detect blind mineralisation and alteration zones in the Que-Hellyer Volcanics. A total of 1031 samples from within EL 106/87 (over the Murrays Road, Medway and Leven River prospects) were collected as part of a larger geochemical sampling program covering CML 103M/87 and EL's 3/95 Hatfield River (McNeill, 1997) and 13/94 Mt. Cattley (Hicks, 1997). Samples were collected from the "B" horizon at a nominal 50 metres spacing on a 200-600 metre spaced grid,

from dominantly residual soils. Duplicate and standard samples were added to the sample set which was assayed for Cu, Pb, Zn, Ba, As, Cd, Co, Ni, and Sb by both partial and total digest methods. Results of this survey are discussed in McNeill and Hicks (1997). A single coherent multi-line Cu, Pb, Zn, and Cd anomaly, the Mayday anomaly, was selected for diamond drill testing (see section 5 for results).

#### *Wacker*

Two deep geochemical sampling (Wacker) programs were completed in areas of cover or poor outcrop;

The Cripps Fault prospect (McNeill, 1989a): Sampling in an area of deep, up to >29m thick, glacial cover was designed to provide information on the location of the Que-Cripps fault. Some geochemical anomalism (up to 925 ppm Zn, 6000 ppm Ba and 210 ppm As) is associated with the fault.

The Link Road prospect (McNeill and Hicks, 1997): Sampling around a zone of fuchsite alteration and barite veining (hangingwall style alteration?) in Southwell Subgroup volcanoclastics indicated that the alteration had a restricted distribution and was not associated with anomalous base metals.

#### 4.3.3 Drill core

Drill holes in the QHV have routinely been sampled over their entire length by continuous core grinds. Samples are selected on lithological, alteration or mineralisation boundaries, or over 10-15m intervals where lithologies are uniform. Short intervals (<1-2m) are often cut. Prior to 1986 samples were routinely analysed for Cu, Pb and Zn,  $\pm$  Ag and Au, however, as a result of the work of Hespe (1986), lithochemical (Cr, Ti and Zr) and pathfinder elements (Ba, As) indicator elements have been routinely analysed. Sample preparation and analytical methods are summarised in Hespe (1986). As with soil samples, a standard (prepared from drill core) has been routinely included in all batches of core-grinds submitted for analysis since 1985.

During the life of EL 106/87 many older drillholes (pre-1985) were re-ground and(or) existing pulps of grinds were re-analysed for lithochemical and pathfinder elements to complete the core geochemical dataset (see McNeill (1989) and McNeill et al. (1990) for details). The new lithochemical data was compared with lithological descriptions, from drill logs, and discrepancies noted. The differences between visual and chemical classifications were then resolved by re-description of thin sections and(or) re-examination of core (McNeill et al, 1990;

McNeill and Wallace, 1991) and any significant changes to the geological interpretation incorporated into the regional cross sections and interpreted geology plans.

Core grinds of any QHV intersected in drillholes more recent than MAC32 (and including MAC29) have been analysed for whole-rock, and sulphur, in addition to the suite discussed above. Whole-rock analyses are used to:

- Help define alteration zones. Studies in the Que-Hellyer Volcanics (Jack, 1989; Gemmill, 1990, McNeill et al., 1990, and Henley, in McNeill and Wallace, 1991) indicated several potential vectors to alteration and mineralisation, particularly the alteration index of Ishikawa et al. (1976) which indicates Ca and Na mobility.

- Aid in rock classification and correlation as discussed by Crawford (1991) and Crawford et al. (1992).

#### 4.3.4 Rock-chip / channel sampling

Rock chip and channel sampling have been carried out in conjunction with geological mapping throughout the life of EL's 2/70, 15/73 and 106/87. In total 638 rock chip samples and 695 channel samples have been collected. Channel samples, normally over 10m intervals, were often taken when new access tracks and costeans were cut through areas of geological interest. Samples were generally analysed for Cu, Pb, Zn, Ag, Au, and (post 1985) Ba, As, Cr, Zr and Ti. Results are included in relevant annual reports.

#### 4.3.5 Pb-isotopes

Pb-isotopes have been demonstrated to be useful discriminants between economic Cambrian massive sulphide systems (Que River, Hellyer and Rosebery) and Devonian vein systems (Gulson and Porritt, 1987). During the life of EL 106/87 two samples; from the Henty Fault Zone and Black Harrys Road, were analysed by SIROTOPE. Results are included in McNeill et al. (1990).

### 4.4 Geophysics

#### 4.4.1 Ground EM

##### *Time-domain*

Given that EM surveys were instrumental in the discovery of the Que River and Hellyer ore deposits (Webster and Skey, 1979; Eadie et al., 1985), ground EM became a key technique in the exploration of the Que-Hellyer Volcanics (QHV). As stated in section 3, by the time of

granting of EL 106/87 most of the outcropping QHV had been covered by fixed loop time-domain EM (see plates MAC113B and C in McNeill, 1989a for reading line locations). EM surveys completed in the reporting period were therefore designed to:

1. Complete coverage of potential buried extensions to the QHV (Hellyer north survey, in McNeill et al., 1990).
2. Infill holes in the previous coverage of the QHV (Charter Dolerite survey, in McNeill et al., 1990).
3. Test poorly understood responses located in previous surveys (I-Zone surveys, in McNeill, 1989a and McNeill et al., 1990).
4. Explore exposed felsic volcanic sequences of the Southwell Subgroup (Medway survey, in McNeill et al., 1990, and the Southwell valley survey, in McNeill and Wallace, 1991).
5. Explore beneath largely Tertiary Basalt covered areas for both Southwell Subgroup and Mt. Cripps Subgroup volcanics (Mackintosh North survey, in Richardson, 1996. Note that this survey also covered parts of ELs 3/95 Hatfield River [McNeill, 1997] and 13/94 Mt. Cattley [Hicks, 1997]).

Only one anomaly considered worthy of drill testing was located by these surveys; a response in the Mackintosh North area (Leven River prospect) between 17000N and 18000N and centred at 9900E. This response was tested by DDH MAC41 with negative results (Richardson, 1996; see section 4.5 for details). The area of EL106/87 is now considered to be adequately tested for conductive targets of greater than Que River size, to within approximately 150m of surface, where ground EM coverage is effective (i.e., outside the powerline corridor). Test work at Hellyer and Que River has demonstrated that downhole EM (DHEM) is effective in locating mineralisation at >150m from drillholes, provided loop designs allow effective coupling with the target (Eadie, 1987; Silic and Eadie, 1989). As a result DHEM became a routine procedure in drilling programs on EL 106/87. A range of systems have been used including SIROTEM, EM-37, UTEM, Crone PEM and the Zonge GDP-16. Results of surveys are summarised in section 4.5. No off-hole conductors attributable to massive sulphide mineralisation were detected and the only two non-lithology related anomalies (in MAC28 and MAC40) were instrumental effects.

In addition to its direct ore-finding role, ground time-domain EM has been used, in an "in-loop sounding" mode to infer two types of stratigraphic information:

1. To estimate the thickness of Tertiary basalt cover at Middlesex Road, prior to drilling MAC20 and 22 (McNeill, 1989), and at the Leven River prospect, prior to drilling MAC41 (Richardson, 1996).

2. To estimate the depth to top of the Que River Shale over the AGSO seismic line and at the Mayday structural / geochemical target area (McNeill and Hicks, 1997).

#### *Frequency-domain*

Frequency-domain EM techniques have not been widely applied on EL 106/87. However, limited Max-Min and VLF surveys were used a first pass technique over previously un-checked DIGHEM anomalies, from the 1975 (Webster, 1975) survey, at the Barite Creek and Southwell Valley prospects (McNeill et al., 1990; Appendix 5). None of the anomalies were considered worthy of follow-up with time-domain EM.

#### 4.4.2 Seismic

Three seismic surveys have been conducted:

1. A two line (10800N and 10950N) reflection survey conducted by the University of Tasmania over the immediate environs of the Hellyer orebody (Read, 1986 and 1989).

2. The results from lines 10800N and 10950N encouraged Aberfoyle to collect two further lines of reflection data; a regional section on 10600N from the Murchison highway to Hellyer, and a section on 11800N north of Hellyer. Survey specifications and results are included in McNeill (1989b; Appendix 7). Data quality was poor, due to ground roll and air blast effects. Results suggest that the Hellyer orebody would not have been located by seismic reflection alone, but that stratigraphic variations could be resolved.

3. As part of the TASGO project, AGSO completed a seismic reflection line along the Cradle Mt. Link Road. A segment of this line (6300E to 7300E) was shot with closer spaced (10m) stations than the regional traverse. An initial interpretation of the partially processed and un-migrated data indicated that the survey may have mapped the top of the Que-Hellyer Volcanics. Two further lines, totalling 3.0 km, with a north-south orientation, were collected on CML 103M/87 and EL 106/87 in April 1995. This survey was tied in to a deep drill hole (HL469) north of Hellyer to provide geological control. Survey specifications are included in McNeill and Hicks (1997). Bends in the traverse have made a comprehensive interpretation difficult. However, some conclusions are presented in McNeill and Hicks (1997) and have been published by Yeates et al. (1997).

#### 4.4.3 Magnetics

##### *Ground*

During the life of EL 106/87 a total of 48 line km of ground magnetic data was collected from the Mt. Charter dolerite and South Hatfield (McNeill, 1989a), the South Mackintosh (McNeill et al., 1990) and the Southwell River (McNeill and Wallace, 1991) prospects. Data was collected at 10m spacing using an OMNI EDA4 proton precession magnetometer. This data was then dumped to a porta-pak computer and combined with data from a second OMNI magnetometer used in base station mode. Corrected data were then combined with previous surveys (see McNeill, 1989b). However, problems with levelling these different surveys and with data quality, meant the data sets could not be integrated without producing severe striping in both N-S and E-W orientations, that masked any subtle structural trends that otherwise may be evident in the data.

Although the QHV are relatively magnetically quiet, the ground data did show some prominent geological features. Sufficient magnetic susceptibility contrast appeared to exist for geological features to be clearly defined if a high quality aeromagnetic survey were to be undertaken.

##### *Airborne*

During march 1993 Geotrex conducted a 1087 line km helicopter borne magnetic survey over EL 106/87 and CMLs 103M/87 and 68M/84. The survey was designed to confirm geological trends and more accurately locate structures inferred from geological and geochemical data. Survey specifications and the processing of results are summarised in Richardson (1993, and 1994).

The QHV are generally magnetically quiet with only a few weakly magnetic dacite bodies. Most sources and breaks are shallow features reflected in the high frequency part of the data. Therefore, spatially persistent breaks that are inferred to reflect faults are probably Devonian or younger in age. However, many shallow aeromagnetic breaks mark major faults with geological evidence such rapid stratigraphic thickness and facies changes or localisation of hydrothermal alteration, indicating they must have been active during the Cambrian. A general fall in magnetic intensity from south to north is interpreted to reflect progressive deepening of the magnetic Central Volcanic Complex to the north. The regional gradient is terminated on a

major NW trending structure through Hellyer that coincides approximately with a major regional gravity discontinuity of the same orientation.

#### *Magnetic susceptibility measurements*

Magnetic susceptibility data have been collected from drill core to assist in interpreting magnetic surveys, assist with stratigraphic subdivision, and to determine if magnetite is responsible for spurious EM responses in some drill holes. Results and techniques are discussed in McNeill (1989b; Appendix 12) and McNeill and Hicks (1997).

#### 4.4.4 Radiometrics

Radiometric data (total count, U, Th and K) were collected during the 1993 helicopter borne aeromagnetic survey (see Richardson (1993) for survey specifications). Results have proved to be of little use. Tertiary basalt, Cambrian dolerite and cultural features such as roads and powerlines are the most prominent features and are best seen in the total count data (Richardson, 1994).

#### 4.4.5 Gravity

Gravity measurements were seen as a method of supporting the syn-volcanic fault model derived from geological and magnetic data, and in conjunction with magnetics could be used in 2-D modelling to refine cross sections through the Que-Hellyer Volcanic basin.

Prior to 1993 existing gravity data was restricted to detailed surveys over the Hellyer and Que River orebodies (Leaman and Richardson, 1981; Hudspeth and Richardson, 1985), wide (1-2 km) spaced data collected as part of the Mount Read Volcanics project, and a close spaced traverse over the Mt. Charter Dolerite (Hespe, 1986). To complete coverage a total of 530 infill stations, on available access throughout the EL, were read during 1993/94. All stations were then surveyed and levelled and data reduction undertaken by Leaman Geophysics. The new data was then merged with the existing surveys on EL 106/87 and CMLs 68M/84 and 103M/87 (summarised above) and with open-file data from the adjacent ELs 39/85 (Placer; Leaman, 1988) and 37/89 (Pasmenco). Station locations, numbering and reduced gravity values are shown on Plates MAC413A, B, and C (Richardson, 1995).

#### **4.5 Diamond drilling**

Nineteen diamond drill holes, for a total of 13,460 m, were completed by Aberfoyle Resources on EL 106/87 (see Figure 8 for collar locations). In addition three holes were drilled on EL 106/87 by the Tasmanian Department of Mines.

The Department of Mines holes are:

**MAC24 (488.8m)** at Black Harrys Road. Details in McNeill et al. (1990) and Pemberton et al. (1991).

**SBDP10 (321m)** on Beecroft Road (Leven River prospect; Pemberton et al., 1991).

**SBDP6 (316m)** on Racecourse Road (North Hatfield prospect). Details in McNeill (1990) and Baillie and Green (1988).

The aims and results of holes drilled by Aberfoyle Resources are summarised below:

##### **MAC16 (367.4m) Boundary Prospect (South Hatfield) - (McNeill, 1989)**

Designed to test a zone of hangingwall style (K-feldspar - sericite - quartz) alteration associated with barite-pyrite veining in rhyolitic volcanics of the Central Volcanic Complex. Alteration and barite veining were intersected to approximately 12m depth. Native Cu, malachite and chalcocite were recorded deeper in the hole, and may be associated with a thick zone of intense quartz veining. Base metal values were low, maximum 0.2% Cu, 700 ppm Zn, and no off-hole EM conductors were located.

##### **MAC20 (397.5m) Middlesex Road (Leven River) - (McNeill, 1989)**

Designed to test for QHV beneath Tertiary basalt cover south west of the Pancontinental drilling on EL 14/85. Hole intersected volcanoclastics and sediments of the Southwell Subgroup beneath 51m of Tertiary basalt cover. No significant mineralisation, anomalous geochemistry or EM conductors were located. Trilobite fossils from limestone clasts from this hole have been described by Jago and McNeill (1997).

##### **MAC22 (424.3m) Middlesex Road (Leven River) - (McNeill, 1989)**

Following the failure of DDH MAC20 to intersect QHV correlates a step out of 900m to the north was made for MAC22. Assuming the southwesterly dips determined in MAC20 remained constant then this hole should have tested deeper in the stratigraphy. The hole intersected volcanoclastics, sediments and rhyolitic lava of the Southwell Subgroup beneath 144m of younger cover rocks. Only minor syngenetic sphalerite was intersected (18m @ 0.1%Zn, 0.07%Pb from 259-277.3m) and no off-hole conductors were detected. See Van




Figure 8

ABERFOYLE RESOURCES LIMITED EXPLORATION DIVISION	
Date 30/5/1997	North Western Tasmania <b>MACKINTOSH EL106/87</b> DRILL HOLE LOCATION GUIDE
Author: RdeB	
Office BURDIE	
Plate MAC477	
Scale 1:100000	Projection: AMG Zone 55 (AGD 66)

Kerkvoort (1995) and reports by Lees (in McNeill and Wallace, 1991; and Richardson, 1992) for detailed logs and discussion of the volcanology of this hole.

**MAC23 (280m) Mt. Charter - (McNeill et al., 1990)**

DDH MAC23 was designed to test variations in Au grade in the Mt. Charter alteration zone beneath the outcropping massive barite and most intense alteration. The hole failed to intersect the massive barite, the position of which is occupied by a polymict volcanoclastic, but stayed in stringer core type alteration for its entire length. The entire length of this hole was anomalous in Au ( $>0.1$  g/t) with best intersections of 6m@ 2.3 g/t Au and 12 g/t Ag (135-141m) and 6m@ 2.9 g/t Au, 2.8 g/t Ag (216-222m). Base metals were also anomalous with best intersections of 0.25% Cu, 1.2% Pb, and 2.4% Zn (62-64m) and 1m@ 1.9% Pb, 3.74% Zn (197-198m).

**MAC25 (530.3m) S.W. Mt. Charter - (McNeill et al., 1990)**

This hole was designed to test the Que River ore position, at the base of the Mt. Charter dacite, beneath effective UTEM and adjacent to the co-active Mt. Charter fault. A thicker than interpreted Hellyer basalt sequence and a shallower dip on the Mt. Charter fault, meant that the fault was intersected before the Que River ore position. Mineralisation was restricted to "inter-pillow" stringers of pyrite and sphalerite (1.89m@ 0.1% Pb, 2.6% Zn, 5 g/t Ag from 123.6-125.4m). No off-hole conductors were detected.

**MAC26 (850.7m) Mt. Charter - (McNeill and Wallace 1991)**

DDH MAC26 was designed to drill down the core of the steeply north plunging Mt Charter alteration system to test for 1. Au mineralisation and 2. Base metal mineralisation at the Que River ore position below effective ground EM. The hole intersected 447m of stringer core alteration before passing into stringer envelope alteration and a barren ore position at 691.7-731.4m. Intersections include 58.4m@ 0.7% Pb, 1.15% Zn, 32 g/t Ag, 1.75 g/t Au (0-58.4m) and 1.3m@ 3.8% Pb, 7.4% Zn, 29.5g/t Ag and 0.68 g/t Au (245.7-247m). No off-hole conductors were detected.

**MAC27 (883.8m) Charter Dolerite - (McNeill and Wallace 1991)**

This hole was designed to test the Que-Hellyer host horizon below effective EM, in an area of known hangingwall style alteration and unusual sphalerite-rich mineralisation adjacent to the co-active Mt. Charter Fault. A thicker Que River Shale and Hangingwall Basalt package coupled with fault complications meant that the hole failed to intersect the target ore-positions.

Intersections include: 85m@0.1% Pb, 0.3% Zn (249.8-335m) and 1.7m@ 1.23% Zn (759.3-761m). No off-hole conductors were detected. Mineralisation and alteration in this hole are described in detail by Hine (1995; see section 4.6.1).

**MAC28 (907.5m) South Que River - (McNeill and Wallace 1991)**

DDH MAC28 was designed to test the interpreted south plunging blind extension of the Que River alteration system beneath effective EM and to provide stratigraphic information in a poorly drilled area. The hole intersected a zone of stringer envelope (SEZ) type alteration below 716m, which although base metal-poor was anomalous in As (to 56 ppm), a feature typical of outcropping SEZ alteration (e.g., Amoeba Zone). A four loop DHEM survey located a complex off-hole conductor that was recommended for drilling.

**MAC29 (762m) South Que River - (Richardson 1992)**

This hole was designed to test the DHEM response above and to the east of DDH MAC28. A 100m wide fault bounded zone of SEZ alteration was intersected but no obvious source for the DHEM anomaly was seen. Drilling problems meant that the rod string was cemented in the hole from 463-709m and this resulted in the loss of critical DHEM data required to resolve the MAC28 anomaly.

**MAC30 (185m) South Que River - (Richardson 1992)**

**MAC30A (785.1m) South Que River - (Richardson 1992)**

Following the inconclusive test of the MAC28 conductor by MAC29 it was decided to step 100m south for a final attempt to test the target. No well defined alteration zone, ore position or source for the conductor was intersected. However, a response was still evident in the DHEM data. Experimentation with different DHEM systems led to the conclusion that the anomalous responses in MAC28 and MAC30 were due to a previously unrecognised "ferrite core" effect and thus an instrumental artefact resulting from the design of the Zonge SIROTEM and CRONE systems (Richardson, 1994; Appendix 1).

**MAC32 (278.4m) Mt. Charter - (Richardson 1993)**

This hole was designed to test the Que-Hellyer ore position, at the base of the Mt. Charter dacite, in the Mt. Charter alteration zone near the intersection of the interpreted co-active Murchison Highway and Barite Creek structures. The hole was abandoned after strong azimuthal deviation, a result of drilling at 45° to the cleaved Mt. Charter alteration zone. The

entire length of the hole was in the Mt. Charter alteration system and was anomalous in Au, with a best intersection of 32.2m@ 0.4 g/t Au (0-32.2m).

**MAC33 (910.7m) Mt. Charter - (Richardson 1993)**

This hole was designed to test the same target as MAC32, but was drilled from the opposite direction in an attempt to minimise deviation. A complete QHV sequence was intersected, from the Que River Shale to the Animal Creek Greywacke. Disseminated sphalerite (125m@ 0.2% Zn) was present in Hellyer basalt correlates but the hole did not intersect the Murchison Highway structure and therefore remained in the less prospective footwall block. However, a DHEM survey coupled with DHEM in DDH MAC26 indicate that no large conductive bodies are present in the target position.

**MAC34 (31m) Murchison Highway Zone - (Richardson 1993)**

**MAC35 (1189.6m) Murchison Highway Zone - (Richardson 1993)**

These holes were designed to test the Que-Hellyer ore position in the core of the Old Mill site syncline near the intersection of four interpreted syn-volcanic structures. MAC34 was abandoned due to strong azimuthal deviation but MAC35 was successful in testing the target zone. Although the hole intersected a thick polymict volcanoclastic package with minor lava, considered to be the Mixed Sequence, there was no mineralisation or significant footwall style alteration. Once again disseminated sphalerite mineralisation in pillow lava sequence equivalents yielded 71m@ 0.2% Zn. No off-hole conductors were detected. The mineralisation and alteration in DDH MAC35 are also described by Hine (1995).

**MAC36 (814.4m) Mt. Charter - (Richardson 1994)**

This Hole was designed to test the base of the Mt. Charter dacites overlying the interpreted Que-Hellyer structure, along strike from the Mt Charter alteration zone, on section 4500N. The hole intersected dacite then footwall andesites before passing into Animal Creek greywacke, a sequence similar to that intersected by MAC33 and MAC37 (see below). There was no obvious ore position at the base of the dacite but a 48m thick epiclastic dominated unit immediately above the greywacke is variably silica-sericite-pyrite (up to 5% Py) altered. Base metal values are generally low and no off-hole conductors were detected.

**MAC37 (835.2m) Mt. Charter - (Richardson 1994)**

This hole was designed to test polymict volcanoclastics at the base of a dacite wedge overlying the Barite Creek structure and associated with an As soil anomaly. A sequence of dacite and

silicic andesite with minor volcanoclastic (similar to lithologies in MAC33, 35 and 36) overlying the lower Basalt. No significant mineralisation or off-hole conductors were located. Silica-sericite-pyrite (up to 10-15% pyrite) alteration is locally intense, particularly from 270-300m but base metal values are low.

#### **MAC40 (1168.7m) Mt. Charter - (Richardson 1996)**

DDH MAC40 was designed to test two targets; the Mt. Charter position (?Hellyer ore position) down dip and plunge of the barite outcrops and the base of the Mt Charter dacite (the Que River ore position). The hole intersected these potential ore positions but both were unmineralised and the lack of strong alteration suggests that the Mt Charter alteration zone does not focus on the Murchison Highway structure on this section. As with many other holes in the Charter area disseminated sphalerite mineralisation on the Hellyer basalt yielded assays of 88.9m@ 0.22% Zn. No other significant assays were reported. An anomaly was recorded in the initial DHEM survey, but, was recognised as a "ferrite core" artefact by comparing data from air- and metal-cored probes.

#### **MAC41 (353.5m) Mackintosh North (Leven River) - (Richardson 1996)**

This hole was designed to test a TDEM anomaly, in an area of Tertiary Basalt cover, located during the Mackintosh North EM survey. The anomaly was interpreted to be flat lying, with a width of 200m, a depth to top of 250m, and an approximate strike length of 1500m, extending onto the adjacent EL 3/95 (McNeill, 1996). The anomaly was tested by a vertical hole which intersected Tertiary basalt (to 281m) overlying Tertiary sediments (including porous organic-rich material - the interpreted source of the anomaly) and calcareous sandstone and limestone (Ordovician Gordon Group?). No mineralisation was intersected.

## **4.6 Research**

### **4.6.1 Theses**

Post-graduate research projects into aspects of the geology, alteration, and mineralisation of EL 106/87 are:

**Rand (1988);** The Geology and Mineralisation of the Mount Charter Prospect. B.Sc. (Hons.) thesis, University of Tasmania, Hobart. Some of the results of this study were published in Huston et al. (1992).

**Hine (1995); Mafic -hosted zinc mineralisation, High Point, Western Tasmania. B.Sc. (Hons) thesis, University of Tasmania, Hobart.**

**Van Kerkvoort (1995); The geology of the Mt. Cattley-Two Hummocks area, Western Tasmania. B.Sc. (Hons) thesis. University of Tasmania, Hobart. This project largely dealt with EL 14/91 Mt. Tor, but includes drill logs and discussion of DDHs MAC-20 and 22 on EL 106/87.**

**Waters (1995); Volcanology and sedimentology of the host succession to VHMS style mineralisation within the Cambrian Que-Hellyer Volcanics, northwestern Tasmania. PhD. Thesis, Monash University, Melbourne. Progress reports were included in McNeill (1990, 1991) and Richardson (1992, 1993) and part of this study has been published as Waters and Wallace (1992).**

**Glenn Lees; The eruptive and depositional history of the Cambrian Upper Rhyolitic Sequence, Hellyer region, Mt. Read Volcanic belt, western Tasmania. This M.Sc. project at Monash University was not completed but progress reports are included in McNeill and Wallace (1991) and Richardson (1992).**

#### **4.6.2 Consultants reports**

**Structural study of the Hellyer Que-River area by A. Cooke, World geoscience Corp. Ltd. (in McNeill and Wallace, 1991).**

**Exploration research in the Mount Read Volcanics, Tasmania by R.J. Henley, Epithermex International. A two part report on structural models and lithogeochemistry (in McNeill and Wallace, 1991).**

**Structural setting of the Mackintosh block and its mineralisation by J. Windh and M. Etheridge, Etheridge and Henley, Geoscience consultants (in Richardson, 1993).**

**Review of drill targets and update of structural model of the Mackintosh Basin, NW Tasmania by R. Sliwa, Etheridge, Henley and Williams (in McNeill and Hicks, 1997).**

## 4.7 Access

### 4.7.1 Line cutting

During the life of EL 106/87 a total of 244 line km (including some cutting on CML 103M/87 but, excluding DHEM and ground EM loop lines) of gridding was completed. Gridding completed each year (since 1973) is shown on the MAC153 series of plans, at 1:10,000 scale. Gridding had largely been re-cutting or extending the Mackintosh grid ("mine grid"), initially established prior to the discovery of the Que River deposit. Mine grid has its origin at AMG coordinates 5 388 687.4 mN, 383 972.8 mE (mine grid 000 mN, 000 mE) and with mine grid north oriented at 22° 07' 22" east of AMG (i.e., approximately 10° east of magnetic north).

### 4.7.2 Vehicular tracks /costeans

Early exploration, during the lives of EL 2/70 and 15/73, led to the establishment of a network of 4WD access tracks and costeans over the area of EL 106/87 (these are shown on the MAC89 outcrop mapping series of plans; see reports in Appendix I for details). This access was used during the life of EL 106/87 and additional track construction and costeaning has been minimal. Many drillholes have been sited on, or within 50m, of existing access tracks.

Exceptions were:

- DDH's MAC-36, and 37 on the east flank of Mt. Charter required construction of 900m of access from the Barite Creek track (see plate MAC89-DD).
- DDH's MAC-28, 29 and 30 required construction of a 620m access track from the 6000N costean at I-Zone (see plate MAC89-BB).
- DDH's MAC-32 and 33, north and west of Mt. Charter respectively, required the construction of 150m long access tracks (see plate MAC89-CC).

Similarly only four new costeans have been dug; the 4350E and 4500E costeans at Mt. Charter (plate MAC89-CC McNeill et al., 1990) and the 1615E and 1645E costeans at Black Harrys Road (plate MAC89-3; McNeill et al., 1990). Rehabilitation of these tracks and costeans is discussed in section 6.0.

## **5.0 WORK COMPLETED IN THE FINAL YEAR**

### **5.1 Introduction**

Work during the current reporting period has focussed on drill testing of the Mayday target. During 1996 a partial digest soil geochemical survey was conducted over a corridor along strike from the Hellyer and Que River orebodies; from Hellyer mine, ten kilometres NNE to the boundary of EL 106/87. The aim of the survey was to detect massive sulphide mineralisation, overlain by barren hangingwall rocks below the penetration depth of surface EM. Results of this survey have been presented in McNeill and Hicks (1997). Only one area of relatively coherent anomalous partial digest geochemistry was detected. This coincided with a previously recognised structural / stratigraphic target, known as the Mayday target due to its proximity to the Mayday fault. An 1800 metre vertical hole was proposed to test the Hellyer ore position beneath the anomalous geochemistry.

DDH MAC 43 was collared on 5-3-97 and completed on 15-11-97 at 1504.8 metres. Collar location is at AMG 5399317.8N 396471.8E (Mine Grid 14558.3N 7577.7E).

### **5.2 DDH MAC 43 - Geology**

A detailed log and list of logging codes is attached as Appendix III, whilst a cross section is included as Plate MAC 479. A summary log is as follows:

0	-	40.4m	Sandstone and conglomerate - Denison Group
40.4	-	112.1	Polymict rhyolitic lapilli to breccia epiclastics - Mt. Cripps Subgroup
112.1	-	120.5	Mudstone - Mt. Cripps Subgroup (MCSG)
120.5	-	256.8	Quartz - feldspar crystal rich polymict lapilli to breccia epiclastics - MCSG
256.8	-	277.8	Mudstone - MCSG
277.8	-	321.8	Quartz rich volcanogenic sandstone - MCSG
321.8	-	329.1	Mudstone - MCSG
329.1	-	341.5	Quartz rich volcanogenic sandstone - MCSG
341.5	-	363.6	Polymict pumiceous breccia - Southwell Subgroup (SSG)?
363.6	-	523	Rhyolite lava and breccia - SSG
523	-	527.5	Fault Zone
527.5	-	538.6	Sericite altered volcanogenic sandstone - SSG

- 538.6 - 571.4 Grey micaceous siltstone with bands of siliciclastic granule conglomerate - SSG
- 571.4 - 577.6 Polymict pumiceous breccia - SSG
- 577.6 - 642.5 Grey micaceous siltstone with bands of siliciclastic granule conglomerate - SSG
- 642.5 - 699 Polymict ash to breccia pumiceous volcaniclastic - SSG
- 699 - 735.4 Ash volcaniclastic interbedded with siltstone - SSG
- 735.4 - 799 Polymict crystal rich volcaniclastic - SSG
- 799 - 962.2 Polymict pumiceous rhyolitic breccia to coarse lapilli and minor ash volcaniclastic - SSG
- 962.2 - 1043.7 Quartz - feldspar crystal volcaniclastic - SSG
- 1043.7 - 1069.9 Shale - SSG
- 1069.9 - 1084.3 Dacitic? Lava and ash volcaniclastic - SSG
- 1084.3 - 1096.6 Interbedded black shale and quartz phyric volcaniclastics - SSG
- 1096.6 - 1174.3 Black shale - SSG
- 1174.3 - 1185.1 Polymict rhyolitic ash to lapilli volcaniclastics - SSG
- 1185.1 - 1214.2 Black shale - SSG
- 1214.2 - 1307.2 Quartz - feldspar crystal volcaniclastic - SSG
- 1307.2 - 1502.9 Polymict pumiceous lapilli to breccia volcaniclastic - SSG
- 1502.9 - 1504.8 Dacite? Lava - SSG

MAC 43 was collared in Upper Owen sandstone and siliciclastic conglomerate. The entire hole was drilled in rhyolitic volcaniclastics with minor lava and sediments of the Mt Cripps and Southwell Subgroups (MCSG and SSG). The base of the Southwell Subgroup was not reached.

The sequence intersected by MAC43 bears only broad resemblance to that exposed in the Cradle Mountain Link road. On the link road the boundary between the MCSG and SSG is placed at the base of a sequence of siltstone and siliciclastic granule conglomerate that separates SSG felsic lava from a MCSG crystal tuff sequence. In MAC 43 the boundary is tentatively placed at the gradational contact between overlying quartz rich volcanogenic sandstone and underlying pumice bearing breccia at 341.5m. About 20m below this contact a flow banded quartz  $\pm$  feldspar phyric rhyolite lava was intersected. A felsic lava is also exposed in the Cradle Mountain Link road at the top of the SSG. In MAC 43 the felsic lava is underlain by 80m of siltstone and siliciclastic granule conglomerate which under the current

interpretation would presumably represent a very thin Murrays Road greywacke. Alternatively this interval of sediment may represent the basal unit of the MCSG.

The remainder of MAC43 intersected an alternating sequence of polymict pumiceous epiclastics and grey to black siltstone and shale, correlated with the SSG. Minor unfossiliferous? limestone clasts were recorded from some SSG breccias similar to MAC 20 and MAC 22 to the north.

Core to bedding angles in the sediments indicate stratigraphy to be shallow dipping above the rhyolite lava body but to steepen with depth until dips around 80° dominate. Core orientation was not possible until the hole flattened sufficiently to allow an orientation device to be used. Successful orientations are consistent with stratigraphy generally dipping to the ESE but below around 1100m becoming steeply overturned to the NW. Overturning is supported by a general mirror image repetition of lithological units around 1100m.

Given the interpretation that below 1100m MAC 43 was drilling overturned stratigraphy it was decided to stop and complete downhole EM in order to provide encouragement to continue drilling. As no DHEM response was obtained and no stratigraphic indication of proximity to the Que - Hellyer volcanics was evident it was decided to terminate the hole. The source of the partial digest soil anomaly therefore remains unresolved.

Trace disseminated pyrite occurred throughout the hole and only very weak local galena ± sphalerite ± chalcopyrite veining was intersected. Narrow intervals were reported from 733.2 - 733.3m, 905.8 - 905.9m, 1043.7 - 1067.9m, 1077 - 1079.6m. Minor massive pyrite and variably Si+Py±Se±Cl altered volcanic clasts, with 15-30% Py, were intersected between 646.85 and 681.6m. These are not known elsewhere in the SSG and their significance, other than indicating hydrothermal activity, is unclear.

No PVC casing was used for DHEM in MAC43 and the HW casing was left in the hole to allow for possible extension by future explorers.

### **5.3 DDH MAC 43 - Geochemistry**

Only local assays were carried out on MAC43. Core grinds were taken of weakly pyritic felsic volcaniclastic between 60 and 100m. Ten metre samples were assayed for whole rock and Zr,

Cr, Ba, As, Cu, Pb, Zn, Ag and Au. Results are included in Appendix IV. Base and precious metal values are uniformly low.

Core grind and split core samples were also taken over several intervals containing weak base metal veining between 733 and 1130m. These samples were analysed for Cu, Pb, Zn, Ag, Au, Ba and As. Sample intervals and results are included in Appendix IV. A summary of mineralised intervals is as follows:

- 733.2-733.3m, 0.1m@ 0.14% Cu, 1.76% Zn, 5.52% Pb, 27 g/t Ag. (strongly Gn+Sp veined shale).
- 1043.7-1050.3m, 6.6m@ 990 ppm Pb, 0.34% Zn, 6 g/t Ag (?Devonian veins in shale and volcanoclastics).
- 1075.3-1079.6m, 4.3m@ 0.27% Pb, 6 g/t Ag (?Devonian veins in volcanoclastic).
- 1106.1-1116.1m, 10m@ 500 ppm Pb, 0.12% Zn, 2 g/t Ag (?Devonian veins in shale).

In order to confirm the origin of lead in galena bearing veinlets, a sample from 733.2m was submitted to the CSIRO for lead isotope analysis. The sample returned a clear Hellyer signature. Results are included in Appendix V.

#### **5.4 DDH MAC 43 - Geophysics**

##### **5.4.1 Downhole EM**

At the time of casing off to NQ at around 500m it was proposed to carry out in-hole depth soundings. The aim was to refine the estimate of depth to the top of the Que River shale that had been obtained from surface soundings and was a basis for the expected depth to the Que Hellyer Volcanics in the target area. Equipment failures caused the sounding to be delayed until 570m. Unfortunately, after carrying out both downhole and surface tests (at 570m and 740m) it appears that no available downhole probe is capable of measuring, with suitable quality, the extremely low signal from deep, poorly conductive stratigraphy. Consequently, meaningful results could not be obtained and the original surface based estimate of 900-1500m to the top of the Que River Shale could not be improved.

A two loop downhole EM survey ( Zonge GDP-16) was undertaken in an open hole from 1504.8m back to the HQ casing at 525m. Loop locations and survey results are shown in

Appendix VI. No off-hole conductors were located and the only response is due to the shale unit intersected below 1050m.

#### 5.4.2 Magnetic Susceptibility

Magnetic susceptibility measurements of drill core from MAC 43 were taken at one metre intervals over the full length of the hole. Results are included in Appendix VII.

Mt. Cripps Subgroup volcanics are variably magnetic until a lithological boundary at 227.4m. From this contact to what is interpreted as the base of the Mt. Cripps Subgroup, susceptibilities are low. They remain low within the Southwell Subgroup except for two intervals of volcanoclastic (710.4 - 724.3 and 758.6 - 782m) which have distinctive pink and green mottled and banded alteration typical of the Tyndall Group. One other zone of moderately elevated values is associated with a red siliceous Fe rich interval of volcanoclastic breccia between 1442 and 1477m.

## **6.0 ENVIRONMENTAL DISTURBANCE AND REHABILITATION**

### **6.1 Introduction**

Exploration over the life of EL 106/87 and preceding EL's 2/70 and 15/73 resulted in the establishment of several kilometres of 4WD tracks and associated diamond drill pads. A survey of all drill sites and tracks was undertaken and a preliminary plan of proposed rehabilitation formulated. This was discussed with Mineral Resources Tasmania resulting in agreement on the final scope of work. Work to be completed over the summer of 1997 / 1998 is shown on Plate MAC 478 and summarised below.

### **6.2 To be left open**

The following tracks will not be rehabilitated as they are considered to provide key access into remote or prospective areas. Barriers will be constructed at their junction with the HEC transmission line to prevent casual access but will allow for re-opening by future explorers if required.

- The main track accessing Mt. Charter (excluding tracks and drill sites near the summit).
- The Barite Creek track (excluding the MAC36/37 access tracks and drill sites).
- The South Charter track.

### **6.3 To be rehabilitated**

Rehabilitation will be carried out using an excavator to drag stockpiled soil back onto tracks from the verges. Some older tracks have not been used for several years and consequently now have well established regrowth. These will not be disturbed. Several drill holes in the vicinity of Mt. Charter are making water. These will be re-entered and sealed prior to track and pad rehabilitation. The following tracks and drill sites will be rehabilitated:

- MAC43 drill site and access track (north of Plate MAC 478).
- MAC28/29/30 drill sites and access track from the HEC line to MAC30.
- MAC34/35 drill sites and access track from the HEC line to MAC35.
- The first 200m of the MC14 access track off the High Point track.
- The disused lower portion of the Mt. Charter track accessing MAC33.
- The top section of the Mt. Charter track accessing MAC23/26.
- MAC36/37 drill sites and access track back to the junction with the Barite Creek track.

- The disused portion of the South Charter track adjacent to the HEC line.
- MC17 drill site and access track off the South Charter track.

## **7.0 CONCLUSIONS**

The focus of the final years work on Mackintosh EL 106/87 was diamond drill testing of the highest ranked partial leach soil geochemical anomaly, identified during the 1996 TMI survey. This anomaly coincides with a structural / stratigraphic target identified in 1994. Targeted Que - Hellyer Volcanics were not reached as they occur at greater than anticipated depth. No source for the geochemical anomaly is evident.

EL 106/87 has been the focus of Aberfoyle's' Tasmanian exploration effort for the past ten years but no new massive sulphide mineralisation has been discovered since the Hellyer orebody in 1983. Many targets with varying geological, geochemical and geophysical support were drill tested during that time. Other, lower ranked targets remain to be tested.

**8.0 EXPENDITURE**

The following expenditure was incurred on EL 106/87 Lake Mackintosh for the period February 1997 to the time of writing (1<sup>st</sup> January 1998). Final expenditure will be forwarded to Mineral Resources Tasmania upon completion of rehabilitation work.

GEOLOGY	\$125,099
SURVEY	\$3,446
GEOPHYSICS	\$76,015
GEOCHEMISTRY	\$3,297
DIAMOND DRILLING	\$284,019
ACCESS	\$24,104
REHABILITATION	\$10,829
OTHER SERVICES	\$13,975
ADMINISTRATION	\$55,366
<b>TOTAL</b>	<b>\$596,150</b>

## 9.0 REFERENCES

- Anon, 1986. Summary of exploration in the Mackintosh District, 1969 to 1986. Unpub. Report to Director of Mines by Aberfoyle Resources Pty. Ltd. (TCR 86-2521)
- Baillie, P.W., and Green G.R., 1988. Completion report sub-basalt drilling project hole 6. Unpub. Rept. Dept. Mines Tasmania, 1988/06.
- Barton, C. M. et al., 1966. Geological Atlas 1 Mile Series. Zone 7; sheet 44 (8014N) Mackintosh. Department of Mines Tasmania.
- Barwick, D. G., 1991. Geology and Mineralisation of the Sock Creek and High Point Areas, Western Tasmania. Unpub. BSc (Hons) Thesis. University of Tasmania, Hobart. 107p.
- Bold, A., 1977. The Geology of the Que River Area West Tasmania. Unpub. BSc (Hons) Thesis. Monash University, Melbourne. 133p.
- Collins, P. L. F., 1981, The Mount Read Volcanics and Associated Rocks in the Pinnacles-Que River Area. In Collins, P. L. F., et al 1981, Geological Atlas 1 Mile Series. Sheet 44 (8014N). Mackintosh Explan. Rep. Department of Mines Tasmania, p117-144.
- Corbett, K. D., 1992. Stratigraphic-Volcanic Setting of Massive Sulphide Deposits in the Cambrian Mount Read volcanics, Tasmania. *Econ. Geol.* 87:564-586.
- Corbett, K. D. and Komysan, P., 1989. Geology of the Mount Charter Area. Geological Report, Mount Read Volcanic Project Tasm. 1. Tasmanian Department of Mines.
- Corbett, K. D. and McNeill, A. W., 1986. Geology of the Rosebery - Mount Block Area. Map 2. Mount Read Volcanics Project. Department of Mines Tasmania.
- Crawford, A.J., 1991. Petrology, geochemistry and correlation of lavas in Bulgobac River drillholes BRD-01, 02, 03 and 05, W. Tasmania. Unpub. Rept. to Placer Exploration (TCR91-3214B)
- Crawford, A.J., Corbett, K.D., and Everard, J.L., 1992. Geochemistry of the Cambrian Volcanic-Hosted Massive Sulfide-Rich Mount Read Volcanics, Tasmania, and Some Tectonic Implications: *Econ. Geol.*, 87:597-619.
- Downs, R.C., 1990, Geological mapping and logging at Hellyer: *Australas. Inst. Mining Metall. Northwest Queensland Branch, Mine Geologists' Conf. Papers, Mt. Isa, Australia, October 1990, p. 1-7.*
- Downs, R.C., 1993. Syn-depositional fault controls on the Hellyer volcanic-hosted massive sulphide deposit. Unpub M. *Econ. Geol. Thesis, University of Tasmania, Hobart, 63pp.*
- Drown, C.G., and Downs, R.C., 1990. Deformational style and strain partitioning at the Hellyer Volcanogenic massive sulphide deposit (Abstr.) in Gondwana: terranes and resources. 10<sup>th</sup> Australian Geological convention, Hobart, 176-177.
- Drown, C.G., and Richardson, S.M., 1990, Hellyer massive sulphide deposit: *Geol. Soc. Aust., Gondwana: Terranes and Resources, 10<sup>th</sup> Australian Geological Convention, Hobart, Australia, February 1990, Excursion Guide E1, 8 p.*
- Eadie, E.T., 1987. The downhole EM response of the Hellyer ore deposit. *Explor. Geophys.* 18: 255-264.

- Eadie, E.T., 1988, Discovery and geology of Que River and Hellyer, in Australas. Inst. Min. Metall. Annual Conference, July 1988.
- Eadie, E.T. and Silic, J., 1984, The application of geophysics in the discovery of the Hellyer ore deposit, in AMIRA Technical Meeting, Exploration, Minerals and Petroleum, Melbourne 1984, pp.11-14 (AMIRA Limited: Melbourne)
- Eadie E.T., Silic, J., and Jack, D.J., 1985. The application of geophysics to the discovery of the Hellyer ore deposit, Tasmania. Explor. Geophys. 16: 207-209.
- Gardner, D. N., 1986. S Lens Mineralisation at Que River. Unpub. BSc (Hons) Thesis. University of Tasmania, Hobart. 78p.
- Gemmell, J.B., 1990. Hellyer Stringer Zone Project. Progress report. Confidential report to Aberfoyle Resources Ltd. CODES, University of Tasmania.
- Gemmell, J.B., 1991, Evolution of the alteration zone and stringer system beneath the Hellyer volcanogenic massive sulfide deposit, Tasmania, Australia [abs.]: Meet. Geol. Association of Canada, Toronto, January 1991.
- Gemmell, J.B., and Large, R.R., 1990, Geologic evolution of the stringer zone underlying the Hellyer volcanogenic sulphide deposit, Tasmania: Pacific Rim Congress Proc., v. II, Gold Coast, Queensland, Australia, May 1990, p. 385-392.
- Gemmell, J. B., Large, R. R., 1992. Stringer System and Alteration Zones Underlying the Hellyer Volcanic Hosted Massive Sulphide Deposit, Tasmania, Australia: Econ.Geol., 87: 620-649.
- Gemmell, J.B., Large, R.R., and Barreiro, B., 1990,  $\delta^{34}\text{S}$  and Pb isotope study of the stringer zone beneath the Hellyer sulphide deposit, Tasmania [abs.]: Seventh International Conference on Geochronology and Isotope Geology, Canberra, Australia, September 1990.
- Gemmell, J.B., Large, R.R., McArthur, G.J., Drown, C.G., and Downs, R.C., 1990, Formation of the alteration pipe and stringer zone beneath the Hellyer massive sulphide deposit, Tasmania [abs.]: Geol. Soc. Aust., Gondwana: Terranes and Resources, 10<sup>th</sup> Australian Geological Convention, Hobart, Australia, February 1990, Abstracts No. 25, p. 8-9.
- Gulson, B.L. and Porritt, P.M., 1987. Base metal exploration in the Mount Read Volcanics, western Tasmania, Pt. II: Lead isotope signatures and genetic implications. Econ. Geol., 82:291-307.
- Hespe, A. M., 1986. Hatfield EL 15/73 Tasmania, Internal Technical Progress Report for the Period June, 1985 to May, 1986. (TCR 87-2663)
- Hicks, D.J., 1996. Exploration Licence 13/94, Mt. Cattley, Tasmania. Relinquishment Report. Aberfoyle Resources Ltd. Unpub. Report. (TCR 96-3925)
- Hine, R. D., 1995. Mafic -hosted zinc mineralisation, High Point, Western Tasmania. Unpubl. BSc (Hons) thesis, University of Tasmania, Hobart, 106p.
- Hudspeth, J.W., 1986, A gravity survey of the Hellyer deposit, IN Large, R.R., ed., The Mount Read Volcanics and associated ore deposits: Geol. Soc. Aust. Tas. Div., Hobart, p. 45-48.

- Hudspeth, J.W., and Richardson, R.G., 1985. A preliminary gravity survey at the Hellyer prospect. Tasmania Department of Mines Unpub. Rept. 1985/25.
- Huston, D.L., Bottrill, R.S., Creelman, R.A., Khin Zaw, Ramsden, T.R., Rand, S.W., Gemmell, J.B., Jablonski, W., Sie, S.H., and Large R.R., 1992. Geologic and geochemical controls on the mineralogy and grain size of gold-bearing phases, eastern Australian volcanic-hosted massive sulfide deposits. *Econ. Geol.* 87: 542-563.
- Ishikawa, S., Sawaguchi, T., Iwaya, S., and Horiuchi, M., 1976. Delineation of prospecting targets for Kuroko deposits based on modes of volcanism of underlying dacite and alteration halos. *Mining Geology*, 26:105-117 (in Japanese with english Abstr.).
- Jack, D. J., 1989. Hellyer Host Rock Alteration. Unpub. MSc. Thesis. University of Tasmania, 181p.
- Jack, D.J., 1990, Geochemistry and alteration of the hangingwall basalts to the Hellyer volcanogenic massive sulphide deposit, Tasmania [abs.]: *Geol. Soc. Aust., Gondwana: Terranes and Resources*, 10<sup>th</sup> Australian Geological Convention, Hobart, Australia, February 1990, Abstracts No. 25, p. 7.
- Komyshan, P., 1986, Geology of the Hellyer-Mt. Charter area, in *The Mount Read Volcanics and associated ore deposits* (Ed. Large, R.R.), pp.11-19 (*Geol. Soc. Aust., Tas. Div.: Hobart*).
- Jago, J.B., and McNeill, A.W., 1997. A late middle Cambrian shallow water trilobite fauna from the Mount Read Volcanics, Northwestern Tasmania. *Pap. Proc. Roy. Soc. Tasm.* 131: 85-90.
- Komyshan, P.K., 1986. Geology of the Mount Charter-Hellyer Area. Tasmanian Dept. Mines, Mt Read Volcanics Project. Map 1.
- Large, R.R., McGoldrick, P.J., Berry, R.F., and Young, C.H., 1988. A tightly folded, gold rich, massive sulfide deposit: Que River mine, Tasmania: *Econ. Geol.*, 83:681-693.
- Leaman, D., 1988. Que Road Gravity Survey. EL 39/85 Unpub. Report for CSR Ltd. (TCR 89-2909A).
- Leaman, D., and Richardson, R.G., 1981. Gravity survey of the Que River deposit, western Tasmania. Tasmania Department of Mines. Unpub. Rept. 1981/24.
- Mather, J., 1985. Time domain induced polarisation spectral analysis from the Que River-Hellyer area. Unpub. BSc (Hons) thesis, Geology Department, University of Tasmania.
- McArthur, G.J., 1986, The Hellyer massive sulphide deposit, IN Large, R.R., ed., *The Mount Read Volcanics and associated ore deposits: Geol. Soc. Aust. Tasm. Div., Hobart*, p. 11-19.
- McArthur, G.J., 1988, Using geology to control geostatistics in the Hellyer deposit: *Math. Geol.*, v. 20, no. 4 p. 343-366.
- McArthur, G.J., 1990, Metal and textural zonation in the Hellyer massive sulphide deposit [abs.]: *Geol. Soc. Aust., Gondwana: Terranes and Resources*, 10<sup>th</sup> Australian Geological Convention, Hobart, Australia, February 1990, Abstracts No. 25, p. 108-109.

- McArthur, G.J., and Dronseika, E.V., 1990, The Que River and Hellyer volcanogenic Ag-Pb-Zn sulphide deposits, IN Hughes, F.E., ed., *Geology of the mineral deposits of Australia and Papua New Guinea: Australas. Inst. Mining Metall. Monograph No. 14*, p. 1229-1239.
- McArthur, G.J., and Kuipers, A.J., 1990, Computer methods for mine geology at Hellyer: Australas. Inst. Mining Metall. Northwest Queensland Branch, mine Geologists' conf. Papers, Mt. Isa, Australia, October 1990, p. 49-55.
- McArthur, G. J., 1996. Textural Evolution of the Hellyer Massive Sulphide Deposit. Unpub. Phd. Thesis, University of Tasmania. 272p.
- McGoldrick, P.J., Large, R.R., 1992. Geologic and Geochemical Controls on Gold-Rich Stringer Mineralization in the Que River Deposit, Tasmania: *Econ. Geol.*, 87: 667-685.
- McGoldrick, P.J., Large, R.R., and Jenkins, G.W., 1990, The footwall precious metal zone at Que River: Transition from a VMS to an epithermal mineral assemblage (abs.): *Geol. Soc. Australia Abstracts*, v. 25, p. 293-294.
- McNeill, A. W., 1989a. Lake Mackintosh EL 106/87, Tasmania. Technical Progress Report for the Period February, 1988 to February, 1989. Report to Department of Mines, Tasmania by Aberfoyle Resources Ltd. (TCR 89-2948)
- McNeill, A. W., 1989b. Exploration Licence 2/70 Tasmania. Technical Progress Report for the period December 1985 to February 1988. Aberfoyle Resources Ltd., unpublished report. (TCR 89-3006)
- McNeill, A. W., 1990. Exploration Licence 106/87. Report on Areas to be relinquished February 1990. (TCR 90-3073)
- McNeill, A.W., 1996. Exploration Licence 3/95 Hatfield River, Tasmania. Final Report. Aberfoyle Resources Ltd. Unpub. Report. (TCR 96-3878)
- McNeill, A. W. and Corbett, K. D., 1989. Geology of the Tullah-Mount Black Area. Geological Report, Mount Read Volcanic Project., Tasm. 2. Tasmania Department of Mines.
- McNeill, A. W., and Hicks, D. J., 1997. Exploration Licence 106/87. Progress Report for the period February, 1996 to February, 1997. Report to Department of Mines, Tasmania by Aberfoyle Resources Ltd. (TCR 97-3991)
- McNeill, A. W., Rand, S.W., and Henham, R.J., 1990. Lake Mackintosh, Exploration Licence 106/87, Tasmania. Technical Progress Report for the Period February, 1989 to April, 1990. Report to Tasmania Department of Mines by Aberfoyle Resources Ltd. (TCR 90-3128)
- McNeill, A. W., and Wallace, D.B., 1991. Exploration Licence 106/87. Progress Report for the Period May, 1990 to April, 1991. Aberfoyle Resources Ltd. Unpub. Report. (TCR 91-3268)
- McPhie, J. and Allen, R.L., 1992. Facies Architecture of Mineralised Submarine Volcanic Sequences: Cambrian Mount Read Volcanics, Western Tasmania. *Econ. Geol.* 87:587-596.
- Odgers, P., 1985. A comparison between time domain and frequency domain induced polarisation in relation to stratigraphy in the northern part of the Que River area. Unpub. Honours thesis, University of Melbourne.

- Offler, R., and Whitford, D., 1992. Wall-rock alteration and metamorphism of a volcanogenic massive sulfide deposit at Que River, Tasmania: *Petrology and mineralogy: Econ. Geol.*, 87: 686-705.
- Pemberton, J., Vicary, M. J., and Corbett, K. D., 1991. *Geology of the Cradle Mountain Link Road - Mt Tor area. Geol. Rep. Mt. Read. Volc. Proj. Tasm.* 4
- Ramsden, A.R., Kinealy, K.M., Creelman, R.A., French, D.H., 1990, Precious and base-metal mineralogy of the Hellyer volcanogenic Ag-Pb-Zn sulphide deposit, northwest Tasmania - a study by electron microprobe, IN Gray, P.M.J., Bowyer, G.J., Castle, J.F., Vaughan, D.J., Warner, N.A., eds., *Sulphide deposits - their origin and processing: The Inst. Mining and Metall.*, p. 49-71.
- Rand, S. W., 1988. *The Geology and Mineralisation of the Mount Charter Prospect. Unpub. B.Sc. (Hons.) Thesis, University of Tasmania.* 140p.
- Read, J.J., 1986. *Shallow seismic reflection survey of the Hellyer orebody. Unpub. BSc(Hons.) thesis. Geology Department, University of Tasmania.*
- Read, J.J., 1989. *Seismic reflection investigations of the Hellyer orebody and Que-Hellyer Volcanics, north west Tasmania. Explor. Geophys.* 20:159-162.
- Richardson, S. M., 1992. *Exploration Licence 106/87. Progress Report for the Period April, 1991 to April, 1992. Aberfoyle Resources Ltd. Unpub. Report. (TCR 92-3355)*
- Richardson, S. M., 1993. *Exploration Licence 106/87. Progress report for the Period April, 1992 to April, 1993. Aberfoyle Resources Ltd. Unpub. Report. (TCR 93-3441)*
- Richardson, S. M., 1994. *Exploration Licence 106/87. Progress Report for the Period April 1993 to February 1994. Aberfoyle Resources Ltd. Unpub. Report. (TCR 94-3537)*
- Richardson, S. M., 1995. *Exploration Licence 106/87, Lake Mackintosh, Tasmania. Progress report for the period February 1994 to February 1995. Aberfoyle Resources Ltd. Technical report. (TCR 95-3719)*
- Richardson, S. M., 1996. *Exploration Licence 106/87, Lake Mackintosh, Tasmania. Progress Report for the Period February 1995 to February, 1996. Aberfoyle Resources Ltd. Technical report. (TCR 96-3839)*
- Staltari, G., 1986. *The Que River TEM case study. Exploration Geophysics* 17, 125-128.
- Scott, D., 1988. *Stratigraphic and Volcanogenic Analysis of the Hellyer Upper Rhyolite Sequence, Western Tasmania. Unpub. B.App.Sci. Report. University of Technology, Sydney.*
- Sharpe, R., 1991. *The Distribution, Mineralogy and Paragenesis of the Hellyer Baritic and Siliceous Caps. Unpub. BSc (Hons) Thesis. University of Tasmania, Hobart.* 114p.
- Silic, J., and Eadie, E.T., 1989. *DHEM: the Que-Hellyer Volcanics experience. Explor. Geophys.* 20: 65-70.
- Silic, J., Eadie, E.T., and Jack, D.J., 1985, *Application of time-domain electromagnetic methods in the discovery of the Hellyer deposit, Tasmania, Australia: SEG 55<sup>th</sup> Annual Int. Meet. & Exposition, 1985, Expanded Abstracts, p. 241-243.*
- Sinclair, B. J., 1994. *Geology and Geochemistry of the Que River Shale, Western Tasmania. Unpub. BSc (Hons) Thesis. University of Tasmania, Hobart.* 114p.

- Sise, J.R., and Jack, D.J., 1984, Exploration case history of the Hellyer deposit [abs.]: IN Baillie, P.W., and Collins, P.L.F., eds., mineral exploration and tectonic processes in Tasmania: Geol. Soc. Aust. Tasm. Div., Hobart, November, 1984.
- Skey, E.H., 1984, Dualism in the discovery of the Que River and Hellyer polymetallic sulphide deposits in western Tasmania [abs.], Geol. Soc. Aust., 12, pp.489-491.
- Skey, E.H., and Young, C.H., 1980, Que River Zn-Pb- deposit, Dundas trough, Tasmania: Jour. Geochem. Explor., v. 12, p. 284-290.
- Staff of Aberfoyle Resources, 1986, Geology and discovery of the Que River and Hellyer polymetallic sulphide ores, Tasmania. (DRAFT) (See no. 11).
- Staff of Aberfoyle Resources, 1989, Exploration case history of the discovery of the Hellyer polymetallic massive sulphide deposit: IN Burrett, C.F., and Martin, E.L., eds., Geology and Mineral Resources of Tasmania: Geol. Soc. Aust., p. 430-434.
- Staff of Aberfoyle Resources, 1990, Geology and discovery of the Que River and Hellyer polymetallic sulphide ores, Tasmania, IN Glasson, K.R., and Rattigan, J.H., eds., Geological aspects of the discovery of some important mineral deposits in Australia: Australas. Inst. Mining Metall. Monograph No. 17, p. 187-196.
- Sliwa, R., 1996. Review of drill targets and update of structural model of the Mackintosh Basin, NW Tasmania. Etheridge, Henley and Williams, Geoscience Consultants Unpub. Report to Aberfoyle Resources Ltd.
- Van Kerkvoort, G. A., 1995. The geology of the Mt. Cattley-Two Hummocks area, Western Tasmania. Unpub. BSc (Hons) thesis. University of Tasmania, Hobart. 144p.
- Vicary, M. J., and Pemberton, J., 1988, Geology of the Back Peak - Cradle Mountain Link Road Area. Map 7 Mount Read Volcanics Project. Department of Mines Tasmania.
- Wallace, D.B., 1984, Developments in the geological understanding of the Que River deposits: Geol. Soc. Australia-Dept. Mines Mineral Exploration and Tectonic Processes in Tasmania Symposium, Burnie, Tasmania, November 1984, p. 55-56.
- Wallace, D. B., 1993. Exploration Licence 106/87, Lake Mackintosh Partial Relinquishment Report for Second Statutory Relinquishment, February 1993. (TCR 93-3410)
- Wallace, D.B., and Green, G.R., 1982, Que river mine and aspects of the Mount Read Volcanics in the Pieman River area, in Geology mineralization, exploration: Western Tasmania. A symposium in honour of the late K.O. Reid. Queenstown, November 1982: Queenstown, Geol. Soc. Australia, Tasmanian Div., p. 52-66.
- Warren, W., 1989. Sulphides in the Hanging Wall Basalts of the Hellyer Massive Sulphide Deposit. Unpub. B.App.Sci. Thesis. University of Technology, Sydney. 100p.
- Waters, J., 1990, Styles of sedimentation and volcanism within the Que-Hellyer Volcanics [abs.]: Geol. Soc. Aust., Gondwana: Terranes and Resources, 10<sup>th</sup> Australian Geological convention, Hobart, Australia, February 1990, Abstracts No. 25, p. 32-33
- Waters, J. C., 1995. Volcanology and sedimentology of the host succession to VHMS mineralisation within the Cambrian Que-Hellyer Volcanics, northwestern Tasmania. Unpub. PhD thesis, Monash University, Melbourne. 303p.

- Waters, J.C., and Wallace, D.B., 1992. Volcanology and sedimentology of the host sequence to the Hellyer and Que River volcanic-hosted massive sulfide deposits, northwestern Tasmania. *Econ. Geol.* 87: 650-666.
- Webster, S. S., 1975. Airborne Geophysical Survey of an Area within EL 2/70, Mackintosh, EL 15/73 Hatfield and EL 5/74 Mayday. Unpub. Report to Cominco Exploration Pty Ltd. (TCR 75-1134)
- Webster, S.S., and Skey, E.H., 1979. Geophysical and Geochemical Case History of the Que River Deposit, Tasmania, in Hood P.J. ed. *Geophysics and Geochemistry in the Search for Metallic Ores. Geol. Surv. Canada. Econ. Geol. Rep.* 31:697-720.
- Whitford, D.J., and Craven, S.J., 1986, Strontium isotopic studies at Que River and Hellyer, IN Large, R.R., *The mount Read Volcanics and associated ore deposits: Geol. Soc. Aust. Tasm. Div., Hobart, p. 87-88.*
- Whitford, D.J., McPherson, W.P.A., and Wallace, D.B., 1989. *Geochemistry of the Host Rocks of the Volcanogenic Massive Sulphide Deposit at Que River, Tasmania. Econ. Geol.* 84:1-21.
- Windh, J., and Etheridge, M., 1992. *Structural Setting of the Mackintosh block and its Mineralisation. Etheridge and Henley, Geoscience Consultants Unpub. Report to Aberfoyle Resources Ltd.*
- Yeates, A.N., McNeill, A.W., Richardson, S., Barton, T.J., Drummond, B.J., Silic, J., and Richardson, R.G., 1997. High-resolution reflection seismic in the Hellyer ore environment. (Abstr.) *New developments in research for ore deposit exploration. Specialist group in Economic geology, GSA, .*
- Young, C.H., 1980. *Que River, a synclinally folded stratiform volcanogenic sulfide deposit: Aberfoyle Explor. Pty. Ltd., unpub. Rept. 23p.*
- Zaw, K., Gemmell, J.B., Large, R.R., Mernagh, T.P., and Ryan, C.G., 1994. Evolution of VHMS fluids in the footwall stringer zone, Hellyer deposit, western Tasmania: Constraints from fluid inclusion microthermometry and geochemistry: 12<sup>th</sup> Australian Geological Convention, Perth, Australia, *Geol. Soc. Aust. Abstr*, 37: 482.
- Zaw, K., Gemmell, J.B., Large, R.R., Mernagh, T.P. and Ryan, C.G., 1996, Evolution and source of ore fluids in the stringer system, Hellyer VHMS deposit, Tasmania, Australia: evidence of hydrothermal ore genesis, *Ore Geol. Reviews*, vol10, pp.251-278.
- Zaw, K., Gemmell, J. B., Hunns, S. R., Mernagh, T. P., Ryan, C. G., Large, R. R., Both, R. A., 1995. Composition of Fluid Inclusions from the Hellyer and Mt. Chalmers VHMS Deposits, Australia: Implications for Source of Ore-Forming Fluids. (Abstr.) PACRIM '95. p663-668.

**APPENDIX I**

## Appendix I.

### Chronological listing of open file reports for EL 2/70, Mackintosh, and 15/73, Hatfield.

#### EL 2/70 Mackintosh:

- Varley, R. J., 1971. EL 2/70, Report on the Mackintosh Area 1970-71. Report to Tasmania Department of Mines by Aberfoyle Management Pty Ltd. (TCR 71-737)
- Krummei, G., 1972, EL2/70 Mackintosh. Progress report on the Mackintosh area. Aberfoyle Management Pty. Ltd., unpublished report. (TCR 72-881)
- Slade, J. and Webster, S. S., 1972. Report on Combined Helicopter Magnetic and EM Survey in the Mackintosh River Area (EL 2/70) Tasmania, for Paringa Mining and Exploration ltd. (TCR 72-863)
- Skey, E. H., 1974. EL2/70 Mackintosh. Progress report on exploration. Cominco Exploration Pty. Ltd., unpublished report. (TCR 74-1020)
- Webster, S. S., 1975. Airborne Geophysical Survey of an Area within EL 2/70, Mackintosh, EL 15/73 Hatfield and EL 5/74 Mayday. Unpub. Report to Cominco Exploration Pty Ltd. (TCR 75-1134)
- Skey, E. H., 1975. EL2/70 Mackintosh. Report on exploration for six months ending June 1975. Cominco Exploration Pty. Ltd., unpublished report. (TCR 75-1125)
- Skey, E. H. and Webster, S. S., 1976. EL2/70 Mackintosh. Report on exploration for year ending June 1976. Cominco Exploration Pty. Ltd., unpublished report. (TCR 76-1167)
- Young, C. H., 1977. EL's 2/70 Mackintosh and 15/73 Hatfield. Report on exploration for year ending June 1977. Cominco Exploration Pty. Ltd., unpublished report. (TCR 77-1222)
- Young, C. H., 1978. Annual Report, EL 2/70 Mackintosh Tasmania, for twelve Months ending 1 June, 1978. Unpub. Report. Aberfoyle Resources Ltd. (TCR 78-1277)

Young, C. H., 1979. EL's 15/73 Hatfield and 2/70 Mackintosh. Report on exploration for year ending June 1979. Aberfoyle Exploration Pty. Ltd., unpublished report. (TCR 79-1360)

Young, C. H., 1980. EL's 15/73 Hatfield and 2/70 Mackintosh. Report on exploration for year ending June 1980. Aberfoyle Exploration Pty. Ltd., unpublished report. (TCR 80-1459)

Sise, J. R., 1981. EL's 15/73 Hatfield and 2/70 Mackintosh West. Report on exploration for year ending June, 1981. Aberfoyle Exploration Pty. Ltd., unpublished report. (TCR 81-1579)

Sise, J. R., 1981. Interim Report, EL 2/70 Mackintosh Tasmania for six Months Ending December 20, 1981. Report to Department of Mines, Tasmania by Aberfoyle Resources Ltd. (TCR 82-1806)

Sise, J. R., 1983. EL2/70 Mackintosh. Report on exploration for year ending December 1982. Aberfoyle Exploration Pty. Ltd., unpublished report. (TCR 83-1939)

Jack, D. J., 1984. EL2/70 Mackintosh. Report on exploration from January 1983 to April 1984. Aberfoyle Exploration Pty. Ltd., unpublished report. (TCR 84-2277)

Hespe, A. M., 1986, Mackintosh EL 2/70, Tasmania. Progress Report for the Period May, 1984 to December, 1985. Report to Department of Mines, Tasmania by Aberfoyle Resources Ltd. (TCR 86-2551)

McNeill, A. W., 1989. Exploration Licence 2/70 Tasmania. Technical Progress Report for the period December 1985 to February 1988. Aberfoyle Resources Ltd., unpublished report. (TCR 89-3006)

**EL 15/73 Hatfield:**

Skey, E. H., 1975. EL15/73 Hatfield. Report on exploration for six months ending June 1975. Cominco Exploration Pty. Ltd., unpublished report. (TCR 75-1148)

- Skey, E. H. and Webster, S. S., 1976. EL15/73 Hatfield. Report on exploration for year ending June 1976. Cominco Exploration Pty. Ltd., unpublished report. (TCR 76-1168)
- Young, C. H., 1977. EL's 15/73 Hatfield and 2/70 Mackintosh. Report on exploration for year ending June 1977. Cominco Exploration Pty. Ltd., unpublished report. (TCR 77-1222)
- Young, C. H., 1978. EL15/73 Hatfield. Report on exploration for year ending June 1978. Abminco NL, unpublished report. (TCR 78-11276)
- Young, C. H., 1979. EL's 15/73 Hatfield and 2/70 Mackintosh. Report on exploration for year ending June 1979. Aberfoyle Exploration Pty. Ltd., unpublished report. (TCR 79-1360)
- Young, C. H., 1980. EL's 15/73 Hatfield and 2/70 Mackintosh. Report on exploration for year ending June 1979. Aberfoyle Exploration Pty. Ltd., unpublished report. (TCR 80-1459)
- Sise, J. R., 1981. EL's 15/73 Hatfield and 2/70 Mackintosh Report on exploration for year ending June 1981. Aberfoyle Exploration Pty. Ltd., unpublished report. (TCR 81-1579)
- Sise, J. R., 1982. EL 15/73 Hatfield. Report on exploration for six months ending December 25, 1981. Aberfoyle Exploration Pty. Ltd., unpublished report. (TCR 82-1807)
- Sise, J. R., 1983. EL15/73 Hatfield. Report on exploration for year ended December 25, 1982. Aberfoyle Exploration Pty. Ltd., unpublished report. (TCR 83-1940)
- Hespe, A. M., 1984. Hatfield EL 15/73 Tasmania. Internal Technical Progress Report for the Period 26/12/82 - 27/5/84. Aberfoyle Resources Ltd. (TCR 85-2355)
- Hespe, A. M., 1984. Hatfield EL 15/73 Tasmania. Report on Exploration 28 May, 1984 - 25 December, 1984. Report to Tasmania Department of Mines by Aberfoyle Resources Ltd. (TCR 85-2429)
- Hespe, A. M., 1985. Hatfield EL 15/73 Tasmania. Internal Technical Progress Report for the Period 26 December, 1984 to 27 May, 1985. (TCR 86-2555)

Hespe, A. M., 1986. Hatfield EL 15/73 Tasmania, Internal Technical Progress Report for the Period June, 1985 to May, 1986. (TCR 87-2663)

McNeill, A. W., 1989. Exploration Licence 15/73, Tasmania. Technical Progress Report for the Period June, 1986 to February, 1988. Report to Department of Mines, Tasmania by Aberfoyle Resources Ltd. (TCR 89-3005)

## **APPENDIX II**

## Appendix II.

### Summary of untested identified structural / stratigraphic targets on EL 106/87

#### TARGET NAME: MAYDAY 1

##### TARGET BACKGROUND:

Lower Mt. Cripps Subgroup volcanogenic sandstones are restricted to S side of Mayday Fault. Mayday Fault is inferred Cambrian transfer which when extrapolated to the E also controls Back Peak Beds (Lower Mt. Read Volcs. ) and location of anomaly 13 and Anio Ck. gold mineralisation. Magnetic interpretation suggests Que-Hellyer Structure may extend to NE of Hellyer and cross Mayday transfer at Mayday 1.

**Structure:** Target lies at intersection of inferred Que-Hellyer Corridor and Mayday Fault transfer.

**Stratigraphy:** QHV at great depth

**Structural activity:** Localisation of units against Mayday Fault indicates Cambrian activity.

**Alteration and geochemistry:** Massive pyrite clasts at Murrays Road associated with minor fuchsite.

**Indicators:** Murrays road massive pyrite clasts at base of Mt. Cripps Subgroup.

**DRILLING:** MAC43 to 1505m failed to reach the QHV

#### TARGET NAME: QUE SOUTH

##### TARGET BACKGROUND:

In MAC29 and MAC30 Se+Si+Py altered clasts occur in epiclastics and breccias in the transition zone between Feldspar Phyrlic Andesite and Lower Basalt. These can be geochemically anomalous eg. MAC29 157.5-162.2 0.18% Zn.

**Structure:** Target defined by inferred Que - Hellyer Structure near its intersection with the Barite Creek Structure.

**Stratigraphy:** Target is at the Lower Basalt - Feldspar Phyrlic Andesite contact.

**Structural activity:** Rapid thinning of the Lower Basalt is inferred across the Que - Hellyer Structure toward the east. No direct evidence of activity.

**Alteration and geochemistry:** Anomalous base metals in soils and in drill core around LB / Afp contact. Hydrothermally altered clasts in epiclastics around this boundary.

**Indicators:** Anomalous geochemistry and altered clasts.

**DRILLING:** 1\* 800metre inclined hole required.

**TARGET NAME: AMOEBA ZONE****TARGET BACKGROUND:**

Altered dacite outcrops along Amoeba Zone. This hole proposes to test the base of the dacite where the alteration zone crosses a possible NW transfer ( Brambles Yard Structure). Only one very shallow hole (HAT3 ) has been drilled in this area.

**Structure:** Target defined by intersection of NNE Que-Hellyer style Amoeba Zone and NW Brambles Yard transfer.

**Stratigraphy:** Target is at base of dacite sequence / top of Feldspar Phyric Andesite.

**Structural activity:** Hydrothermal alteration indicates activity of Amoeba Zone. No other evidence known.

**Alteration and geochemistry:** Amoeba Zone alteration is pervasive sericite+silica+ pyrite. It is base metal poor but generally anomalous in arsenic.

**Indicators:** Amoeba Zone alteration.

**DRILLING:** 1\* 600 metre inclined hole required.

**TARGET NAME: MURCHISON HIGHWAY****TARGET BACKGROUND:**

This target has a setting analogous to Hellyer. Amoeba Zone alteration could be interpreted to plunge S along the axial plane of anticlinally folded Upper Basalt / dacite. This hole proposes to test the base of dacite in the core of the anticline as the Hellyer ore position is probably within surface EM range. DHEM from MAC 35 and Paminco DDH BHD3 restrict the size of target that can be accommodated.

**Structure:** Many inferred coactive structures appear to converge in this area. These are 1) Amoeba Zone

- 2) Que Fault
- 3) High Point

**Stratigraphy:** Target is at base of dacite sequence.

**Structural activity:** Upper Basalt appears to thicken in nose of anticline possibly reflecting thicker stratigraphy above an active structure, as at Hellyer.

**Alteration and geochemistry:** Minor lead anomaly in soil hangingwall to target.

**Indicators:** None known.

**DRILLING:** 1\* 900 metre inclined hole required.

**TARGET NAME: AMOEBA SOUTH****TARGET BACKGROUND:**

Altered dacite outcrops along Amoeba Zone. This hole proposes to test the base of the dacite beneath the best developed alteration and an associated gravity anomaly at the southern end of the outcropping alteration zone.

**Structure:** Target defined by NNE Que-Hellyer style Amoeba Zone.

**Stratigraphy:** Target is at base of dacite sequence / top of Feldspar Phyric Andesite.

**Structural activity:** Hydrothermal alteration indicates activity of Amoeba Zone. No other evidence known.

**Alteration and geochemistry:** Amoeba Zone alteration is pervasive sericite+silica+pyrite. It is base metal poor but generally anomalous in arsenic.

**Indicators:** Amoeba Zone alteration. Gravity anomaly.

**DRILLING:** 1\* 700 metre inclined hole required.

**TARGET NAME: BRONCO****TARGET BACKGROUND:**

This target is defined by the intersection of two inferred Cambrian structures interpreted from magnetics. The proposed hole would test the base of the Upper Basalt and underlying dacite adjacent to the intersection of the two structures. The target would be in the SW quadrant of the intersection where soil geochemistry is elevated and Cambrian subsidence is interpreted to have occurred .

**Structure:** Target defined by intersection of NNE High Point Structure (potentially analogous to Que-Hellyer Structure ) and NW Brambles Yard transfer.

**Stratigraphy:** Target is at base of Upper Basalt and at contact of dacite sequence / top of Feldspar Phyric Andesite.

**Structural activity:** Que River Shale thickens to west (across High Point Structure ? ) in holes on 7700N.

**Alteration and geochemistry:** No alteration known but moderate lithology normalised Pb soil anomaly adjacent to target in hangingwall Upper Rhyolite Sequence and Que River Shale.

**Indicators:** Hangingwall Pb soil anomaly.

**DRILLING:** 1\*800 metre inclined hole required.

**TARGET NAME: MAVERICK****TARGET BACKGROUND:**

Target is lower Upper Rhyolite Sequence (URS) and QHV along strike from Hellyer mine in an area of strong fuchsite alteration and minor barite veining.

**Structure:** Target not well defined by structures identified to date. Possible NE trending (Que Fault trend) magnetic break.

**Stratigraphy:** Target is basal Southwell Subgroup and / or Hellyer ore position.

**Structural activity:** None known.

**Alteration and geochemistry:** Strong pervasive fuchsite alteration of epiclastics well up in the SSG. Rhyolite lava adjacent to the altered epiclastics hosts massive barite veins.

**Indicators:** The hanging wall fuchsite alteration and barite veining. Wacker sampling indicated limited extent of alteration and associated weak Pb anomalism. TMI sampling indicated sporadic elevated Cu Zn and Pb values.

**TARGET NAME: MEDWAY****TARGET BACKGROUND:**

Area has been covered by EM and no conductors identified.

Outcrop is Upper Southwell Subgroup so attractive stratigraphy is inferred to lie at great depth. Several potentially prospective structures are inferred to intersect in this area and a deep magnetic source is interpreted to lie at depth.

**Structure:** NNE trending extension of Que-Hellyer Structure intersects a regional E-W gravity linear. A NNW fault also is interpreted to pass through the area.

**Stratigraphy:** Target would be Lower Southwell Subgroup ( Rosebery host equivalent?? ) or QHV but these are interpreted to lie at considerable depth.

**Structural activity:** No evidence known.

**Alteration and geochemistry:** Sporadic anomalous TMI Cu, Co, As geochemistry.

**Indicators:** None known.

**APPENDIX III**



# Aberfoyle Resources Limited

## HELLYER DIVISION

# DIAMOND DRILL LOG VALID CODES

SHEET 1 OF 2

## LITHOLOGY

### STRATIGRAPHY

Tertiary basalt & gravels	TER
Upper Rhyolitic Sequence	URS
Que River Shale	QRS
Pillow Lava Sequence	PLS
Hanging Wall Volcaniclastic Sequence	HVS
Hellyer Mineralized Sequence	HMS
Feldspar Phyrlic Andesite Sequence	FPS
Lower Epiclastic Sequence	LES
Stringer Enveloper Zone	SEZ
Stringer Zone	STZ

### ROCK TYPE

#### Composition Prefix

Barite	Ba
Basalt	B
Base Metal Sulphides	BMS
Chert	Ch
Dacite	D
Feldspar Phyrlic Andesite	FP
Glassy silica - colloform pyrite ore	GSP
Highly Altered (obliterating primary textures)	HA
Massive Chalcopyrite	MCPy
Massive Pyrite	MPy
Polymict	Y
Pumice	Pu
"Quellite"	Q
Rhyolite	R
Sandstone, Siltstone	Ss
Shale	Sh

#### Volcanic Texture Suffix

Ash volcaniclastic (0-2mm)	av
Lapilli volcaniclastic (2-64mm)	lv
fine (2-8mm)	flv
medium (8-32mm)	mlv
coarse (32-64mm)	clv
Breccia volcaniclastic (>64mm)	bv
Lava	l
Pillow Lava	pl
Variolitic pillow lava	vp

### COLOUR

#### PREFIX

D = Dark  
L = Light  
B = Bright

Black  
Brown  
Green  
Grey  
Orange  
Pink  
Polychrome  
Purple  
Red  
White  
Yellow

Bk  
Br  
Gn  
Gy  
Or  
Pk  
Pc  
Pp  
Rd  
Wh  
Yw

### VOLCANICLASTIC FRAGMENTS

#### Composition

As per rock type composition prefix codes

#### Sorting

Well sorted  
Moderately sorted  
Poorly sorted

W  
M  
P

#### Shape

Angular  
Sub-angular  
Sub-rounded  
Rounded  
Arcuate  
Eutaxitic  
Variable  
Elongate

A  
sA  
sR  
R  
Ar  
X  
V  
E

### VOLCANICLASTIC MATRIX

#### Composition

Crystal  
Glassy  
Lithic  
Vitric  
or any standard mineralogy code

X  
G  
L  
V

### LOWER CONTACT

#### Gradation

Sharp but not faulted  
Faulted  
Gradational up to 1m  
  1-10m  
  over 10m

S  
F  
G1  
G2  
G3

#### Style

Conformable  
Interfingering  
Irregular

C  
F  
I

### VESICLES

#### Concentration

Trace  
Weak  
Moderate  
Strong  
Extreme

1  
2  
3  
4  
5

#### Shape

Annular sub-vesicles  
Elongate  
Spherical  
Variable

A  
E  
S  
V



# Aberfoyle Resources Limited

## HELLYER DIVISION

655062

# DIAMOND DRILL LOG VALID CODES

SHEET 2 OF 2

### ALTERATION

#### Texture

Disseminated	D
Interpillow	I
Nodular	N
Pervasive	P
Pillow Margins	M
Pseudomorphous	O
Rim	R
Selected fragments	F
Stockwork	K
Structure controlled	S
Veins	V

#### Intensity

As per standard intensity coding	1
	2
	3
	4
	5

#### Mineralogy

Carbonate	CO
Chlorite	Cl
Epidote	Ep
Feldspar	Fd
Fuchsite	Fu
Illite	Il
K-Feldspar	Ab
Pyrite	Py
Sericite	Se
Silica	Si
Hematite	He
Tremolite	Tm

Code in order of predominance, use brackets for trace content.

### VEINING

#### Intensity

As per standard intensity coding	1
	2
	3
	4
	5

#### Mineralogy

As per mineralogy codes used for alteration and mineralisation, e.g.	CO
	Si
	Cl
	Ep

### MINERALISATION

#### Texture

Banded	Bn
Brecciated	Br
Colloform	Cm
Disseminated	Ds
Euhedral	Eu
Fragmental	Fr
Framboidal	Fb
Inter-pillow	IP
Joint Plane Coating	JP
Massive	Ma
Nodular	No
Pillow margin	Pm
Recrystallised	Rx
Reworked	Rk
Sheared	Sh
Shrinkage Shadows	Ss
Veins	Vn
Vein Selvedge	VS
Boxwork Veining	Bx
Fragment Rim	Rm

#### Mineralogy and Contents

Mineralogy prefix, percentage suffix (tr for trace), ! for emphasis

Arsenopyrite	As
Barite	Ba
Chalcopyrite	Cp
Galena	Gn
Pyrite	Py
Sphalerite	Sp
Gangue	Ga

### FAULTS

#### Gouge

Brecciated	Br
Cavernous	Cv
Puggy	Pu
Rehealed	Rh
Slickensided	Sl

Carbonate	CO
Chlorite	Cl
Mylonite	My
Quartz	Si
Rubble	Ru
Sericite	Se

### FOLIATION

#### Type

Bedding	Bd
Clast Alignment	CA
Cleavage	Cl
Flow Banding	FB
Shearing	Sh
Sulphide Banding	SB
Vesicle Alignment	VA

### WEATHERING

Type prefix, standard intensity coding suffix

Leached	L
Oxidised	O















































































**APPENDIX IV**

## ABERFOYLE EXPLORATION

DRILL CORE DATA LEDGER  
INTERVAL SAMPLES

LICENCE: EL108/87

TOTAL: 25 samples

## DRILLHOLE MAC-043

PROSPECT: MAY

FROM (m)	TO (m)	SAMPLE#	TYPE	COLLECTED	ORE SUITE								TRACE SUITE				CALCULATED		
					Cu ppm GA101	Pb ppm GA101	Zn ppm GA101	Ag ppm GA101	Au ppm GG309	Ba ppm GX401	As ppm GX401	Cr ppm GX401	Zr ppm GX401	Ti ppm GX401	P2O6 wt% OX408	Ti/Zr	P2O6/TiO2	AI	
60.0	70.0	828991	CGI	AMcN: 1/04/97	38	7	40	-2	-0.01	956	3	17	251	3597	0.09	0.15	84.3		
70.0	80.0	828992	CGI	AMcN: 1/04/97	27	14	52	-2	-0.01	1021	2	11	232	3117.4	0.08	0.15	75.7		
80.0	90.0	828993	CGI	AMcN: 1/04/97	18	17	47	-2	-0.01	895	5	5	231	3117.4	0.09	0.17	71.0		
90.0	100.0	828994	CGI	AMcN: 1/04/97	17	9	44	-2	-0.01	873	5	14	239	3716.9	0.16	0.26	82.8		
733.2	733.3	628774	SCI	AMcN: 1/10/97	1466	54300	18000	27	-0.01	1159	70								
733.3	735.4	628775	CGI	AMcN: 1/10/97	64	352	457	2	-0.01	937	26								
735.4	736.9	628776	SCI	AMcN: 1/10/97	229	34	75	-1	-0.01	504	16								
782.0	787.3	628777	CGI	AMcN: 1/10/97	25	-3	99	-1	-0.01	531	6								
787.3	792.4	628778	CGI	AMcN: 1/10/97	78	137	290	-1	-0.01	862	78								
792.4	807.0	628779	CGI	AMcN: 1/10/97	46	136	166	6	-0.01	1316	46								
951.7	954.5	628780	CGI	AMcN: 1/10/97	22	-3	48	-1	-0.01	5662	4								
954.5	958.7	628781	CGI	AMcN: 1/10/97	7	3	44	-1	-0.01	6602	36								
958.7	962.2	628782	CGI	AMcN: 1/10/97	39	3	80	-1	-0.01	13000	6								
1033.7	1043.7	628784	CGI	AMcN: 1/10/97	35	26	51	1	-0.01	1649	8								
1043.7	1050.3	628785	CGI	AMcN: 1/10/97	125	986	3440	3	-0.01	1329	40								
1050.3	1060.3	628786	CGI	AMcN: 1/10/97	41	131	244	1	-0.01	1575	42								
1060.3	1068.7	628787	CGI	AMcN: 1/10/97	37	71	175	1	-0.01	1337	32								
1068.7	1069.9	628788	SCI	AMcN: 1/10/97	44	37	141	2	-0.01	715	28								
1069.9	1075.3	628789	CGI	AMcN: 1/10/97	21	57	134	1	-0.01	1308	12								
1075.3	1079.6	628790	CGI	AMcN: 1/10/97	54	2756	113	6	-0.01	1466	-2								
1079.6	1084.3	628791	CGI	AMcN: 1/10/97	33	43	122	1	-0.01	1374	6								
1084.3	1096.1	628792	CGI	AMcN: 1/10/97	41	171	464	1	-0.01	1279	26								
1096.1	1106.1	628793	CGI	AMcN: 1/10/97	38	115	188	1	-0.01	1490	44								
1106.1	1116.1	628794	CGI	AMcN: 1/10/97	65	498	1173	2	-0.01	1421	36								
1116.1	1128.3	628795	CGI	AMcN: 1/10/97	39	57	252	-1	-0.01	1151	40								

## DRILLHOLE MAC-043

PROSPECT: MAY

## WHOLEROCK

FROM (m)	TO (m)	SAMPLE#	SiO2 wt%	TiO2 wt%	Al2O3 wt%	Fe2O3 wt%	MnO wt%	MgO wt%	CaO wt%	Na2O wt%	K2O wt%	S wt%	LOI wt%	TOTAL wt%
			OX408	OX408	OX408	OX408	OX408	OX408	OX408	OX408	OX408	OX408	OM615	OX408
60.0	70.0	828991	73.60	0.60	13.30	2.83	0.03	0.51	0.20	1.00	5.95		2.25	100.40
70.0	80.0	828992	71.30	0.52	12.90	4.25	0.14	0.58	0.45	1.49	5.47		2.64	99.70
80.0	90.0	828993	72.90	0.52	12.60	3.86	0.08	0.60	0.60	1.66	4.92		2.37	100.10
90.0	100.0	828994	71.60	0.62	13.00	4.11	0.03	0.78	0.53	0.77	5.48		2.70	99.80
733.2	733.3	628774												
733.3	735.4	628775												
735.4	736.9	628776												
782.0	787.3	628777												
787.3	792.4	628778												
792.4	807.0	628779												
951.7	954.5	628780												
954.5	958.7	628781												
958.7	962.2	628782												
1033.7	1043.7	628784												
1043.7	1050.3	628785												
1050.3	1060.3	628786												
1060.3	1068.7	628787												
1068.7	1069.9	628788												
1069.9	1075.3	628789												
1075.3	1079.6	628790												
1079.6	1084.3	628791												
1084.3	1096.1	628792												
1096.1	1106.1	628793												
1106.1	1116.1	628794												
1116.1	1128.3	628795												





Phone (0041) 316837

14 Thirkell St. COOEE TAS 7320

Fax (0041) 318890

## ANALYTICAL REPORT No.

ABE201.60.12990

THIS REPORT MUST BE READ IN CONJUNCTION WITH THE ACCOMPANYING ANALYTICAL DATA

INVOICE TO:

Aberfoyle Resources Limited  
 Exploration Division  
 PO Box 952  
 BURNIE TAS 7320

ORDER No.

PROJECT

11967

DATE RECEIVED

RESULTS REQUIRED

04/04/97

ASAP

No. OF PAGES OF RESULTS

DATE REPORTED

No. OF COPIES

TOTAL No. OF SAMPLES

3

29/05/97

1

5

SAMPLE NUMBERS

SAMPLE DESCRIPTION

ELEMENT/METHOD

828991/5

PU Prep :

Cu,Pb,Zn,Ag/GA101

828991/5

PU Prep :

Wt,Au,Au(R)/GG309

828991/5

PU Prep :

Ba,As,Cr,Zr/GX401

828991/5

PU Prep :

AL2O3,SiO2,TiO2,Fe2O3,MnO,CaO,K2

RESULTS TO

Richard de Bonford  
 Aberfoyle Resources Limited  
 Exploration Division  
 PO Box 952  
 BURNIE TAS 7320

RESULTS TO

RESULTS TO

REMARKS

AUTHORISED OFFICER



## ANALYTICAL DATA

SAMPLE PREFIX

REPORT No.

REPORT DATE

CLIENT ORDER No.

PAGE

SAMPLE PREFIX		REPORT No.				REPORT DATE		CLIENT ORDER No.		PAGE	
		ABE201.60.12990				29/05/97		11967		1 OF 3	
ETHOD	SAMPLE No.	Cu	Pb	Zn	Ag	Au	Au(R)	Ba	As	Cr	
		GA101	GA101	GA101	GA101	GG309	GG309	GX401	GX401	GX401	
1	828991	38	7	40	<2	<0.01	-	956	3	17	
2	828992	27	14	52	<2	<0.01	-	1021	2	11	
3	828993	18	17	47	<2	<0.01	-	895	5	5	
4	828994	17	9	44	<2	<0.01	<0.01	873	5	14	
5	828995 *	124	180	2269	2	<0.01	-	1043	18	891	
6											
7											
8											
9											
10											
11											
12											
13											
14											
15											
17											
18											
19											
20											
21											
22											
23											
24	DETECTION	5	4	5	2	0.01	0.01	10	2	5	
25	UNITS	ppm	ppm	ppm	ppm	ppm	ppm	ppm	ppm	ppm	

sults in ppm unless otherwise specified  
 -- = element not determined

IS = insufficient sample  
 SNR = sample not received

AUTHORISED  
 OFFICER



**ANALYTICAL DATA**

SAMPLE PREFIX

REPORT No.

REPORT DATE

CLIENT ORDER No.

PAGE

ABE201.60.12990

29/05/97

11967

2 OF 3

	SAMPLE No.	Zr	Ti	TiO2	AL2O3	SiO2	Fe2O3	MnO	CaO	K2O
METHOD		GX401	OX408							
1	828991	251	0.36	0.60	13.30	73.60	2.83	0.03	0.20	5.95
2	828992	232	0.31	0.52	12.90	71.30	4.25	0.14	0.45	5.47
3	828993	231	0.31	0.52	12.60	72.90	3.86	0.08	0.60	4.92
4	828994	239	0.37	0.62	13.00	71.60	4.11	0.03	0.53	5.48
5	828995 *	82	0.27	0.45	15.10	51.30	9.93	0.85	3.89	1.19
6										
7										
8										
9										
10										
11										
12										
13										
14										
15										
17										
18										
19										
20										
21										
22										
23										
24	DETECTION	5	0.01	0.01	0.05	0.05	0.01	0.01	0.01	0.01
25	UNITS	ppm	%	%	%	%	%	%	%	%

sults in ppm unless otherwise specified  
 -- element not determined

IS = insufficient sample  
 SNR = sample not received

AUTHORISED OFFICER



**ANALYTICAL DATA**

SAMPLE PREFIX

REPORT No.

REPORT DATE

CLIENT ORDER No.

PAGE

ABE201.60.12990

29/05/97

11967

3 OF 3

	SAMPLE No.	MgO	P2O5	Na2O	LOI	Total			
METHOD		OX408	OX408	OX408	OX408	OX408			
1	828991	0.51	0.086	1.00	2.25	100.40			
2	828992	0.58	0.080	1.49	2.64	99.70			
3	828993	0.60	0.093	1.66	2.37	100.10			
4	828994	0.78	0.157	0.77	2.70	99.80			
5	828995 *	7.47	0.182	3.48	5.44	99.60			
6									
7									
8									
9									
10									
11									
12									
13									
14									
15									
17									
18									
19									
20									
21									
22									
23									
24	DETECTION	0.01	0.005	0.05	0.01	0.01			
25	UNITS	%	%	%	%	%			

Results in ppm unless otherwise specified  
 - = element not determined

IS = insufficient sample  
 SNR = sample not received

AUTHORISED OFFICER





Our reference : BU013747  
 Your reference : 18050  
 Project code : Core Grind & Split Core  
 Date received : 26/09/97  
 Date reported : 26/10/97

**Analabs Pty. Ltd.**  
 ACN 004 591 664  
 14 Thirkell St, Burnie  
 Tasmania 7320  
 Telephone : (004) 31 6837  
 Facsimile : (004) 31 8890

Richard De Bomford  
 Services Officer  
 Exploration Division  
 Aberfoyle Resources Limited  
 Exploration Division  
 P.O. Box 952  
 BURNIE  
 TAS 7320

Number of pages of results : 2  
 Number of Samples : 22  
 First Sample : 628774  
 Last Sample : 628795

Invoice to:  
 Richard De Bomford  
 Services Officer  
 Exploration Division  
 Aberfoyle Resources Limited  
 Exploration Division  
 P.O. Box 952  
 BURNIE  
 TAS 7320

Electronic Data Transmission :  
 Modem Y 26/10/97  
 Facsimile / /  
 Disk Report Y / /

Preliminary Reports :  
 02/10/97 Report  
 09/10/97 Report  
 10/10/97 Report  
 10/10/97 Report  
 16/10/97 Report  
 17/10/97 Report  
 17/10/97 Report  
 17/10/97 Report  
 17/10/97 Report  
 27/10/97 Report

Results to:

Results to:

Remarks :

Authorised by .....  
 On behalf of:

Richard Newman  
 Laboratory Manager

The results in the following analytical report pertain to the samples provided to this laboratory for preparation and/or analysis as requested by the client.







## ANALYSIS DESCRIPTION

Job number : BU013747 Order number : 18050

-----  
Scheme code : S005 - Jaw Crushing to nominal 6mm to 12mm  
-----

Sample preparation. Jaw crushing to nominal 6mm to 12mm.

-----  
Scheme code : S020 - Dry, Ringmill < 500g  
-----

Sample preparation. Dry, Ringmill. < 500g.

-----  
Scheme code : G101 - Perchloric acid digest, Geochemical samples  
-----

Perchloric acid digest, (HClO<sub>4</sub>), Geochemical samples

-----  
Scheme code : A101 - AAS analysis  
-----

AAS analysis of sample after G101 digest.

-----  
Scheme code : F630 - 30g fire assay, Lead collection, AAS  
-----

Fire assay, Lead collection, Aqua Regia digest, AAS, 30g sample.

-----  
Scheme code : X401 - Pressed powder, XRF, Trace determination  
-----

Pressed powder, XRF, Trace determination.

-----  
Scheme code : X406 - Glass fusion, XRF, High level concentrations  
-----

Glass fusion, XRF, High level concentrations, Ilmenite, Rutile, Zircon, Baryte.

-----  
Scheme code : A105 - AAS analysis  
-----

AAS analysis of sample after G105 digest.

## ANALYSIS DESCRIPTION

655115



Job number

: BU013747

Order number

A: 1800 ALAB S

-----  
Scheme code : A120 - AAS analysis  
-----

AAS analysis of sample after G120 digest.

PROJECT <b>MAC 43</b>	BSS SIEVE SIZE CODE - MESH NUMBER A 200 D 80 G 30 B 150 E 60 H 20 C 100 F 40 T = TOTAL	SAMPLE TYPE CODE <input type="checkbox"/> OXIDIZED PRODUCTS O <input type="checkbox"/> FRESH ROCK R <input type="checkbox"/> STREAM SEDIMENTS S	<input type="checkbox"/> WEATHERED BEDROCK W <input type="checkbox"/> SURFACE TRANSPORTED T <input type="checkbox"/> RESIDUAL SOIL E <input type="checkbox"/> MINE DUMP M	CARD PUNCH PRINT YES <input type="checkbox"/> NO <input type="checkbox"/>	VERIFY YES <input type="checkbox"/> NO <input type="checkbox"/>	DATE <b>8-5-97</b>	SHEET <b>1 of 1</b>
--------------------------	--	--	--	--	--	-----------------------	------------------------

EASTINGS							NORTHINGS							SAMPLE NUMBER		DEPTH IN CMS		SIZE FRACTION		Sample Type	METAL VALUES PPM																				GEOLOGICAL LOG																																						
1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21	22	23	24	25	26	27	28	29	30	31	32	33	34	35	36	37	38	39	40	41	42	43	44	45	46	47	48	49	50	51	52	53	54	55	56	57	58	59	60	61	62	63	64	65	66	67	68	69	70	71	72	73	74	75	76	77	78	79	80
MAC43							596.7m							622974																											MURRAYS ROAD GREYNACK																																						
MAC 43							545.5m							622975																											OR BASAL MCSG.??																																						
395300							5398625							622976																											XRF Ba, Th, Nb, La, Ce, P2O5, Zr, TiO2, Y, V, Cr.																																						
395320							5398625							622977																											Basal Mt. Cripps Subgroup from cutting on Cradle Mt. Link Road																																						



Phone (004) 316837

14 Thirkell St. COOEE TAS 7320

Fax (004) 318890

**ANALYTICAL REPORT No.**

ABE201.60.13110

THIS REPORT MUST BE READ IN CONJUNCTION WITH THE ACCOMPANYING ANALYTICAL DATA

INVOICE TO:

Aberfoyle Resources Limited  
 Exploration Division  
 PO Box 952  
 BURNIE TAS 7320

ORDER No.

PROJECT

11987

DATE RECEIVED

RESULTS REQUIRED

08/05/97

ASAP

No. OF PAGES OF RESULTS

DATE REPORTED

No. OF COPIES

TOTAL No. OF SAMPLES

2

20/05/97

1

2

SAMPLE NUMBERS

SAMPLE DESCRIPTION

ELEMENT/METHOD

22974/5

CO Prod : 8P033

Ba,Th,Nb,La,Ce,P,Zr,Ti,Y,V,Cr/GX

RESULTS TO

Richard de Bonford  
 Aberfoyle Resources Limited  
 Exploration Division  
 PO Box 952  
 BURNIE TAS 7320

RESULTS TO

RESULTS TO

REMARKS

AUTHORISED OFFICER

A N A L A B S



## ANALYTICAL DATA

SAMPLE PREFIX

REPORT No.

REPORT DATE

CLIENT ORDER No.

PAGE

		ABE201.60.13110				20/05/97		11987		1 OF 2	
	SAMPLE No.	V	Cr	Y	Zr	Nb	Ba	La	Ce	Th	
ETHOD		GX401	GX401	GX401	GX401	GX401	GX401	GX401	GX401	GX401	
1	622974	164	537	21	152	8	261	22	18	4	
2	622975	117	233	29	197	15	397	29	64	14	
3											
4											
5											
6											
7											
8											
9											
10											
11											
12											
13											
14											
15											
16											
17											
18											
19											
20											
21											
22											
23											
24	DETECTION	5	5	3	5	3	10	10	10	4	
25	UNITS	ppm	ppm	ppm	ppm	ppm	ppm	ppm	ppm	ppm	

Results in ppm unless otherwise specified  
 -- = element not determined

IS = insufficient sample  
 SNR = sample not received

AUTHORISED  
 OFFICER



**ANALYTICAL DATA**

SAMPLE PREFIX

REPORT No.

REPORT DATE

CLIENT ORDER No.

PAGE

		ABE201.60.13110		20/05/97		11987		2 OF 2	
	SAMPLE No.	P205	TiO2						
ETHOD		GX401	GX401						
1	622974	687	5086						
2	622975	756	6571						
3									
4									
5									
6									
7									
8									
9									
10									
11									
12									
13									
14									
15									
16									
17									
18									
19									
20									
21									
22									
23									
24	DETECTION	30	50						
25	UNITS	ppm	ppm						

results in ppm unless otherwise specified  
 -- = element not determined

IS = insufficient sample  
 SNR = sample not received

AUTHORISED OFFICER



Phone (004) 316837

14 Thirkell St. COOEE TAS 7320

Fax (004) 318890

## ANALYTICAL REPORT No.

ABE201.60.13148

THIS REPORT MUST BE READ IN CONJUNCTION WITH THE ACCOMPANYING ANALYTICAL DATA

INVOICE TO:

Aberfoyle Resources Limited  
 Exploration Division  
 PO Box 952  
 BURNIE TAS 7320

ORDER No.

PROJECT

11993

DATE RECEIVED

RESULTS REQUIRED

19/05/97

ASAF

No. OF PAGES OF RESULTS

DATE REPORTED

No. OF COPIES

TOTAL No. OF SAMPLES

2

29/05/97

1

2

SAMPLE NUMBERS

SAMPLE DESCRIPTION

ELEMENT/METHOD

522976/7

CO Prep :

Ba,Th,Nb,La,Ce,P,Zr,Ti,V,Cr/BX

RESULTS TO


Richard de Bonford  
 Aberfoyle Resources Limited  
 Exploration Division  
 PO Box 952  
 BURNIE TAS 7320

RESULTS TO

RESULTS TO

REMARKS

AUTHORISED OFFICER



**ANALYTICAL DATA**

SAMPLE PREFIX

REPORT No.

REPORT DATE

CLIENT ORDER No.

PAGE

SAMPLE PREFIX		REPORT No.				REPORT DATE		CLIENT ORDER No.		PAGE	
		ABE201.60.13148				29/05/97		11993		1 OF 2	
	SAMPLE No.	Ba	Th	Nb	La	Ce	P205	Zr	TiO2	Y	
ETHOD		GX401	GX401	GX401	GX401	GX401	GX401	GX401	GX401	GX401	
1	622976	847	19	18	65	84	824.8	250	6190.3	40	
2	622977	151	7	8	31	44	1076.9	205	6807.5	29	
3											
4											
5											
6											
7											
8											
9											
10											
11											
12											
13											
14											
15											
16											
17											
18											
19											
20											
21											
22											
23											
24	DETECTION	10	4	3	10	10	0.1	5	0.1	3	
25	UNITS	ppm	ppm	ppm	ppm	ppm	ppm	ppm	ppm	ppm	

Results in ppm unless otherwise specified  
 -- = element not determined

IS = insufficient sample  
 SNR = sample not received

AUTHORISED OFFICER



**ANALYTICAL DATA**

SAMPLE PREFIX

REPORT No.

REPORT DATE

CLIENT ORDER No.

PAGE

ABE201.60.13148

29/05/97

11993

2 OF 2

	SAMPLE No.	V	Cr						
METHOD		GX401	GX401						
1	622976	106	136						
2	622977	107	626						
3									
4									
5									
6									
7									
8									
9									
10									
11									
12									
13									
14									
15									
16									
17									
18									
19									
20									
21									
22									
23									
24	DETECTION	5	5						
25	UNITS	ppm	ppm						

Results in ppm unless otherwise specified  
 -- = element not determined

IS = insufficient sample  
 SNR = sample not received

AUTHORISED OFFICER

**APPENDIX V**

**Aberfoyle Resources Ltd**  
ACN 004 664 108

Exploration Division

39 River Road  
Burnie  
Tasmania 7320  
Australia

Telephone (03) 6431 6333  
Facsimile (03) 6431 6896  
email bur@aberfoyle.oz.au

PO Box 952  
Burnie Tasmania 7320  
Australia

9706amn1

**A B E R F O Y L E**

6 June, 1997

Dr Graham Carr  
CSIRO  
Division of Exploration & Mining  
Institute of Minerals, Energy & Construction  
P O Box 136  
NORTH RYDE NSW 2113

Dear Graham/Geoff

As discussed in our conversation of today, please find enclosed a single sample (628773) for urgent Pb-isotope analysis.

This sample comprises galena from a zone of galena-sphalerite-chalcopryrite veining in DDH MAC-43 (733m depth) currently being drilled north of the Hellyer mine. The veins are hosted by volcanoclastic of the Southwell sub-group and we are interested to know if the mineralisation has a Cambrian or Devonian Pb-isotope signature, and if it is Cambrian, is it similar to Que river, Rosebery or Hellyer?

Could you please fax results as soon as they are available?

Yours faithfully  
**ABERFOYLE RESOURCES LIMITED**



**A W McNeill  
GEOLOGIST**

*P.S. Geoff, there was less Galena in the vein than I expected.  
I hope the enclosed sample is sufficient.*

655125


**CSIRO**  
AUSTRALIA
**EXPLORATION  
AND MINING**

 AUSTRALIAN LEADERSHIP IN  
 GLOBAL EXPLORATION AND MINING

 North Ryde Laboratories  
 Riverside Corporate Park  
 West Entrance, 51 Delhi Road  
 PO Box 136  
 North Ryde NSW 2113  
 Australia  
 Telephone 61 2 9887 8666  
 Facsimile 61 2 9887 9821

**FACSIMILE MESSAGE**

Date 13-6-97

 Routine  Urgent  Confidential

No. of Pages 3

To	Andrew McNeill
Company	Aberfoyle Resources
Facsimile No.	03 64316896
Copy	

From	Geoff Denton	
Phone No.	02 9490 8879	Facsimile No. 02 9490 8921
Subject	Pb Isotope Results	

Andrew

Not much interpretation required here.

	$^{206}\text{Pb}/^{204}\text{Pb}$	$^{207}\text{Pb}/^{204}\text{Pb}$	$^{208}\text{Pb}/^{204}\text{Pb}$
1	18.367	15.617	38.224
2	18.377	15.629	38.252


We carried out the analysis in duplicate, and as you can see the Pb isotopic composition of the sample is homogeneous with the Hellyer signature.

Regards

  
 Geoff Denton


Mit Read

655126





655127

13-JUN-1997 15:35  
CSIRO DEM & DPR NTH RYDE  
61 2 94908921 P.003/003




**APPENDIX VI**


Scale 1:10,000




655130



Aberfoyle Resources Ltd  
Datafile: loop1.av  
LOOP: 1  
LINE: 43.00N  
Date Plotted: 1/12/97  
Horiz scale 1: 4848.5  
  
MAC 43 DHM  
  
LOOP 1  
HZ COMPONENT  
ZONGE GOP16 16HZ  
READ/PLOTTED GCC  
ABERFOYLE RESOURCES LTD



655131



Program: 130121  
Aberfoyle Resources Ltd  
Datafile: loop2.av  
LOOP: 2  
LINE: 43.00N  
Date Plotted: 1/12/97  
Horiz scale 1: 4848.5

MAC 43 DHEM

LOOP 2

HZ COMPONENT

ZONGE GOP16 16HZ

READ/PLOTTED GCC

ABERFOYLE RESOURCES LTD

5 cm

**APPENDIX VII**

## MAGNETIC SUSCEPTIBILITY

MAC-043

DEPTH (m)	MAGSUS (SI)								
30	14	116	120	202	1081	288	0	374	0
31	7	117	63	203	1179	289	0	375	8
32	8	118	41	204	1415	290	0	376	0
33	8	119	0	205	1767	291	0	377	0
34	28	120	0	206	1103	292	0	378	0
35	6	121	16	207	1403	293	0	379	0
36	1	122	58	208	1250	294	0	380	0
37	13	123	5	209	1170	295	0	381	0
38	5	124	0	210	1190	296	0	382	0
39	35	125	0	211	1330	297	6	383	0
40	57	126	114	212	1231	298	27	384	0
41	100	127	1264	213	1615	299	2	385	0
42	20	128	1041	214	1145	300	0	386	0
43	46	129	1099	215	960	301	4	387	0
44	5	130	861	216	1360	302	0	388	0
45	42	131	670	217	543	303	0	389	0
46	4	132	751	218	739	304	7	390	0
47	124	133	1087	219	0	305	0	391	0
48	144	134	1918	220	120	306	0	392	0
49	88	135	1301	221	519	307	0	393	0
50	86	136	751	222	357	308	0	394	0
51	47	137	1236	223	570	309	0	395	0
52	218	138	1164	224	246	310	0	396	0
53	39	139	1134	225	35	311	20	397	0
54	223	140	636	226	118	312	3	398	0
55	642	141	956	227	113	313	1	399	0
56	392	142	565	228	57	314	0	400	0
57	100	143	65	229	257	315	0	401	4
58	376	144	1020	230	112	316	0	402	8
59	122	145	647	231	1	317	0	403	2
60	101	146	653	232	50	318	0	404	0
61	70	147	797	233	22	319	12	405	0
62	11	148	579	234	3	320	12	406	0
63	11	149	661	235	0	321	1	407	0
64	31	150	709	236	43	322	0	408	0
65	2	151	568	237	32	323	0	409	12
66	3	152	1123	238	0	324	0	410	0
67	3	153	701	239	0	325	0	411	2
68	3	154	539	240	9	326	0	412	0
69	12	155	348	241	0	327	0	413	4
70	27	156	7	242	0	328	0	414	1
71	12	157	0	243	0	329	0	415	5
72	148	158	0	244	0	330	0	416	11
73	297	159	34	245	0	331	0	417	3
74	3	160	12	246	0	332	11	418	10
75	3	161	0	247	0	333	3	419	4
76	14	162	104	248	0	334	1	420	1
77	11	163	22	249	0	335	0	421	0
78	0	164	0	250	0	336	0	422	3
79	0	165	4	251	0	337	0	423	1
80	0	166	0	252	0	338	4	424	1
81	46	167	296	253	1	339	0	425	1
82	11	168	3	254	27	340	0	426	4
83	143	169	3	255	4	341	0	427	1
84	25	170	613	256	5	342	0	428	6
85	19	171	886	257	0	343	0	429	0
86	19	172	655	258	0	344	0	430	4
87	11	173	12	259	0	345	0	431	2
88	12	174	63	260	0	346	0	432	5
89	12	175	0	261	0	347	0	433	1
90	16	176	5	262	0	348	0	434	8
91	12	177	12	263	0	349	0	435	3
92	31	178	2	264	0	350	0	436	62
93	13	179	67	265	0	351	0	437	4
94	3	180	58	266	0	352	0	438	0
95	12	181	105	267	0	353	0	439	0
96	14	182	4	268	0	354	0	440	0
97	37	183	627	269	4	355	0	441	0
98	13	184	721	270	0	356	0	442	0
99	21	185	716	271	9	357	0	443	0
100	13	186	597	272	0	358	7	444	0
101	8	187	49	273	0	359	1	445	6
102	0	188	0	274	0	360	0	446	0
103	55	189	0	275	0	361	0	447	0
104	18	190	5	276	0	362	0	448	3
105	690	191	358	277	0	363	0	449	0
106	728	192	404	278	0	364	0	450	14
107	343	193	947	279	18	365	0	451	1
108	1124	194	891	280	6	366	0	452	0
109	737	195	346	281	0	367	0	453	0
110	913	196	386	282	0	368	0	454	0
111	17	197	4	283	0	369	0	455	0
112	208	198	88	284	0	370	0	456	0
113	5	199	12	285	0	371	0	457	0
114	72	200	87	286	0	372	0	458	0
115	41	201	224	287	0	373	0	459	0

## MAGNETIC SUSCEPTIBILITY

MAC-043

DEPTH (m)	MAGSUS (SI)								
460	13	546	13	632	5	718	3866	804	0
461	1	547	3	633	2	719	2675	805	0
462	0	548	3	634	0	720	3298	806	0
463	0	549	4	635	0	721	3108	807	0
464	4	550	8	636	0	722	676	808	0
465	5	551	5	637	1	723	46	809	0
466	0	552	7	638	0	724	32	810	0
467	0	553	18	639	0	725	4	811	0
468	17	554	8	640	9	726	1	812	0
469	1	555	4	641	2	727	2	813	0
470	0	556	1	642	0	728	6	814	0
471	6	557	14	643	0	729	5	815	0
472	10	558	4	644	0	730	86	816	6
473	10	559	4	645	0	731	15	817	5
474	0	560	5	646	14	732	14	818	5
475	2	561	19	647	1	733	8	819	5
476	0	562	22	648	0	734	18	820	6
477	0	563	4	649	0	735	22	821	4
478	0	564	5	650	3	736	35	822	2
479	7	565	7	651	7	737	20	823	7
480	1	566	9	652	5	738	57	824	1
481	0	567	3	653	12	739	25	825	2
482	0	568	13	654	3	740	9	826	4
483	0	569	3	655	4	741	5	827	3
484	0	570	37	656	2	742	14	828	1
485	0	571	8	657	0	743	3	829	2
486	0	572	55	658	0	744	1	830	2
487	0	573	8	659	6	745	0	831	3
488	0	574	8	660	1	746	4	832	1
489	0	575	5	661	5	747	20	833	2
490	0	576	0	662	4	748	43	834	0
491	3	577	0	663	4	749	13	835	11
492	0	578	0	664	4	750	13	836	2
493	12	579	0	665	0	751	20	837	0
494	3	580	0	666	0	752	13	838	0
495	1	581	0	667	4	753	8	839	7
496	0	582	5	668	1	754	5	840	2
497	0	583	0	669	2	755	13	841	0
498	0	584	0	670	1	756	6	842	5
499	2	585	0	671	0	757	16	843	2
500	0	586	0	672	52	758	11	844	14
501	2	587	0	673	7	759	33	845	2
502	2	588	0	674	27	760	18	846	3
503	7	589	0	675	7	761	192	847	0
504	8	590	0	676	3	762	140	848	0
505	3	591	0	677	2	763	264	849	0
506	0	592	8	678	3	764	716	850	0
507	0	593	3	679	12	765	865	851	0
508	2	594	0	680	7	766	1147	852	0
509	1	595	5	681	2	767	690	853	0
510	0	596	1	682	1	768	50	854	0
511	0	597	11	683	0	769	38	855	0
512	2	598	5	684	0	770	659	856	0
513	1	599	1	685	0	771	421	857	0
514	4	600	4	686	0	772	309	858	0
515	0	601	9	687	0	773	32	859	0
516	0	602	8	688	2	774	95	860	0
517	0	603	3	689	0	775	79	861	0
518	0	604	3	690	0	776	83	862	0
519	4	605	1	691	0	777	394	863	0
520	2	606	10	692	0	778	236	864	0
521	2	607	7	693	0	779	7	865	0
522	0	608	8	694	0	780	4	866	14
523	0	609	5	695	0	781	5	867	10
524	0	610	3	696	0	782	5	868	3
525	0	611	5	697	0	783	6	869	6
526	0	612	3	698	14	784	7	870	11
527	0	613	2	699	3	785	1	871	6
528	0	614	1	700	47	786	0	872	7
529	0	615	0	701	6	787	0	873	17
530	0	616	0	702	3	788	0	874	2
531	0	617	0	703	2	789	1	875	2
532	0	618	1	704	3	790	1	876	12
533	0	619	1	705	3	791	1	877	66
534	0	620	1	706	0	792	1	878	7
535	0	621	0	707	4	793	1	879	6
536	0	622	0	708	26	794	12	880	4
537	0	623	0	709	5	795	2	881	3
538	0	624	0	710	7	796	3	882	8
539	8	625	0	711	2545	797	4	883	3
540	7	626	1	712	3853	798	5	884	1
541	8	627	1	713	3547	799	0	885	1
542	0	628	2	714	4054	800	0	886	2
543	0	629	1	715	5374	801	0	887	3
544	4	630	10	716	5401	802	0	888	3
545	7	631	5	717	3404	803	0	889	2

## MAGNETIC SUSCEPTIBILITY


MAC-043

DEPTH (m)	MAGSUS (SI)								
890	1	976	0	1062	0				
891	0	977	0	1063	3				
892	0	978	0	1064	0				
893	0	979	0	1065	0				
894	4	980	0	1066	0				
895	0	981	1	1067	2				
896	0	982	1	1068	0				
897	0	983	3	1069	0				
898	0	984	1	1070	0				
899	0	985	0	1071	0				
900	0	986	3	1072	0				
901	0	987	1	1073	0				
902	0	988	1	1074	0				
903	0	989	0	1075	0				
904	0	990	0	1076	0				
905	0	991	0	1077	10				
906	0	992	0	1078	3				
907	0	993	0	1079	0				
908	0	994	0	1080	0				
909	0	995	0	1081	0				
910	5	996	0	1082	0				
911	1	997	0	1083	0				
912	1	998	0	1084	0				
913	0	999	0	1085	0				
914	0	1000	0	1086	0				
915	0	1001	9	1087	0				
916	0	1002	3	1088	0				
917	0	1003	0	1089	0				
918	4	1004	1	1090	0				
919	2	1005	3	1091	6				
920	0	1006	5	1092	1				
921	1	1007	1	1093	0				
922	10	1008	10	1094	0				
923	12	1009	12	1095	0				
924	4	1010	8	1096	0				
925	2	1011	3	1097	0				
926	1	1012	1	1098	0				
927	0	1013	0	1099	0				
928	0	1014	0	1100	0				
929	0	1015	0	1101	1				
930	8	1016	0	1102	1				
931	8	1017	10	1103	0				
932	0	1018	0	1104	0				
933	0	1019	4	1105	2				
934	0	1020	2	1106	11				
935	0	1021	1	1107	5				
936	1	1022	0	1108	1				
937	0	1023	1	1109	0				
938	0	1024	0	1110	8				
939	0	1025	0	1111	2				
940	0	1026	11	1112	2				
941	0	1027	0	1113	0				
942	0	1028	0	1114	0				
943	8	1029	0	1115	0				
944	0	1030	2	1116	0				
945	2	1031	1	1117	7				
946	6	1032	7	1118	1				
947	1	1033	1	1119	13				
948	1	1034	0	1120	4				
949	0	1035	0	1121	0				
950	0	1036	7	1122	0				
951	0	1037	1	1123	0				
952	0	1038	0	1124	0				
953	0	1039	0	1125	0				
954	0	1040	0	1126	0				
955	0	1041	5	1127	1				
956	0	1042	0	1128	5				
957	0	1043	5	1129	0				
958	0	1044	0	1130	0				
959	0	1045	0	1131	0				
960	0	1046	0	1132	0				
961	0	1047	0	1133	0				
962	0	1048	0	1134	0				
963	0	1049	8	1135	1				
964	0	1050	0	1136	5				
965	0	1051	4	1137	0				
966	0	1052	0	1138	0				
967	0	1053	0	1139	0				
968	0	1054	0	1140	0				
969	0	1055	66	1141	0				
970	0	1056	1	1142	0				
971	0	1057	0	1143	0				
972	0	1058	0	1144	0				
973	0	1059	0	1145	0				
974	0	1060	2	1146	0				
975	0	1061	0						

## MAGNETIC SUSCEPTIBILITY

MAC-043A

DEPTH (m)	MAGSUS (SI)								
1130	1	1216	1	1302	0	1388	0	1474	6
1131	2	1217	1	1303	0	1389	0	1475	778
1132	1	1218	3	1304	0	1390	0	1476	67
1133	2	1219	1	1305	7	1391	0	1477	9
1134	1	1220	1	1306	8	1392	0	1478	3
1135	3	1221	1	1307	10	1393	8	1479	26
1136	5	1222	3	1308	5	1394	1	1480	38
1137	2	1223	3	1309	47	1395	1	1481	19
1138	7	1224	19	1310	12	1396	0	1482	5
1139	5	1225	5	1311	12	1397	0	1483	5
1140	5	1226	26	1312	3	1398	0	1484	5
1141	2	1227	4	1313	18	1399	0	1485	0
1142	4	1228	1	1314	7	1400	0	1486	0
1143	4	1229	0	1315	2	1401	4	1487	0
1144	3	1230	4	1316	5	1402	3	1488	13
1145	3	1231	3	1317	2	1403	3	1489	6
1146	1	1232	5	1318	5	1404	5	1490	0
1147	2	1233	4	1319	1	1405	0	1491	0
1148	5	1234	2	1320	0	1406	41	1492	0
1149	2	1235	10	1321	0	1407	9	1493	0
1150	6	1236	9	1322	0	1408	7	1494	0
1151	5	1237	9	1323	0	1409	2	1495	0
1152	7	1238	5	1324	6	1410	1	1496	0
1153	5	1239	10	1325	17	1411	4	1497	0
1154	5	1240	4	1326	10	1412	1	1498	0
1155	7	1241	12	1327	6	1413	0	1499	0
1156	4	1242	16	1328	4	1414	0	1500	0
1157	5	1243	37	1329	9	1415	0	1501	0
1158	3	1244	16	1330	7	1416	0	1502	0
1159	3	1245	22	1331	6	1417	3	1503	0
1160	3	1246	36	1332	5	1418	0	1504	0
1161	4	1247	56	1333	2	1419	0		
1162	6	1248	23	1334	1	1420	0		
1163	4	1249	8	1335	0	1421	0		
1164	4	1250	14	1336	0	1422	0		
1165	11	1251	5	1337	1	1423	0		
1166	4	1252	2	1338	1	1424	0		
1167	10	1253	5	1339	5	1425	0		
1168	9	1254	2	1340	5	1426	0		
1169	5	1255	5	1341	0	1427	0		
1170	4	1256	3	1342	27	1428	0		
1171	3	1257	4	1343	8	1429	0		
1172	7	1258	2	1344	5	1430	0		
1173	2	1259	4	1345	2	1431	1		
1174	15	1260	3	1346	1	1432	0		
1175	7	1261	1	1347	1	1433	0		
1176	5	1262	3	1348	1	1434	0		
1177	13	1263	4	1349	2	1435	0		
1178	17	1264	9	1350	4	1436	0		
1179	7	1265	6	1351	13	1437	0		
1180	4	1266	4	1352	3	1438	0		
1181	11	1267	10	1353	2	1439	0		
1182	35	1268	6	1354	2	1440	0		
1183	9	1269	14	1355	1	1441	0		
1184	21	1270	14	1356	1	1442	0		
1185	6	1271	13	1357	4	1443	0		
1186	14	1272	11	1358	4	1444	0		
1187	13	1273	5	1359	4	1445	0		
1188	20	1274	11	1360	1	1446	0		
1189	13	1275	10	1361	0	1447	0		
1190	8	1276	6	1362	12	1448	0		
1191	14	1277	5	1363	4	1449	5		
1192	7	1278	2	1364	8	1450	85		
1193	19	1279	1	1365	7	1451	108		
1194	24	1280	2	1366	6	1452	71		
1195	8	1281	2	1367	3	1453	14		
1196	6	1282	8	1368	19	1454	80		
1197	6	1283	6	1369	5	1455	88		
1198	6	1284	3	1370	2	1456	17		
1199	6	1285	2	1371	12	1457	236		
1200	2	1286	9	1372	2	1458	260		
1201	2	1287	6	1373	4	1459	102		
1202	1	1288	4	1374	7	1460	25		
1203	1	1289	20	1375	1	1461	8		
1204	0	1290	18	1376	1	1462	192		
1205	1	1291	17	1377	4	1463	24		
1206	3	1292	12	1378	3	1464	46		
1207	1	1293	8	1379	4	1465	103		
1208	1	1294	4	1380	0	1466	9		
1209	12	1295	6	1381	0	1467	53		
1210	5	1296	2	1382	0	1468	13		
1211	2	1297	0	1383	0	1469	1109		
1212	2	1298	4	1384	0	1470	346		
1213	2	1299	7	1385	0	1471	391		
1214	0	1300	4	1386	0	1472	227		
1215	1	1301	0	1387	0	1473	231		




- LEGEND FOR TRACKS**
- Existing
  - Previously Rehabilitated
  - To Be Rehabilitated
  - Established Regrowth - Not To Be Disturbed
  - To Be Left Open For Future Access
  - Planned Barrier For Above

**98-4116**  
 RELINQUISHMENT REPORT-EL 106/87  
 LAKE MACKINTOSH-ABERFOYLE  
 A MCNEILL



<b>Aberfoyle Resources Limited</b> EXPLORATION DIVISION North Western Tasmania SOUTH MACKINTOSH / MT. CHARTER AREA REHABILITATION PLAN		Compiled: RAB/000 Drawn: Traced: Date: November 07 Plan No: MAC/08
REVISIONS No.   Date   Init.   Description	SCALE 1 : 10,000	



98-4116

RELINQUISHMENT REPORT-EL 106/87  
LAKE MACKINTOSH-ABERFOYLE  
A MCNEILL

For geological legend see Appendix I



**Aberfoyle Resources Limited**

EXPLORATION DIVISION

65513S

REVISIONS			
Init.	Date	Init.	Date


NORTH WEST TASMANIA  
MACKINTOSH - E.L. 106/87  
**CROSS SECTION 14 500N**  
**DRILLHOLE MAC43 (A)**

Compiled : S.R.  
Drawn: G.M.B  
Traced :  
Checked : S.R.  
Plate No. : MAC481A

Location Code :

Scale : 1:1000

Date : 07.01.98



98-4116

RELINQUISHMENT REPORT-EL 106/87  
LAKE MACKINTOSH-ABERFOYLE  
A MCNEILL

For geological legend see Appendix I



**Aberfoyle Resources Limited**

EXPLORATION DIVISION

655139

REVISIONS			
Init.	Date	Init.	Date

NORTH WEST TASMANIA  
MACKINTOSH - E.L. 106/87  
**CROSS SECTION 14 500N**  
**DRILLHOLE MAC43 (A)**

Compiled : S.R.  
Drawn: G.M.B  
Traced :  
Checked : S.R.

Location Code :

Scale : 1:1000

Date : 07.01.98

Plate No. - MAC481B